



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>C12N 15/00</b>		<b>A2</b>	(11) International Publication Number: <b>WO 99/58658</b>	
		(43) International Publication Date: 18 November 1999 (18.11.99)		
(21) International Application Number: PCT/US99/10646		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).		
(22) International Filing Date: 13 May 1999 (13.05.99)		<p><b>Published</b></p> <p><i>Without international search report and to be republished upon receipt of that report.</i></p>		
(30) Priority Data:				
09/078,904 60/085,751	13 May 1998 (13.05.98) 15 May 1998 (15.05.98)			US US
(71) Applicant: EPIMMUNE, INC. [US/US]; Suite 100, 5820 Nancy Ridge Drive, San Diego, CA 92121 (US).				
(72) Inventors: PIKES, John, D.; 6474 Lipmann Street, San Diego, CA 92122 (US). HERMANSON, Gary, G.; 3159 Via de Caballo, Encinitas, CA 92024 (US). SETTE, Alessandro; 5551 Linda Rosa Avenue, La Jolla, CA 92037 (US). ISHIOKA, Glenn, Y.; Apartment J5, 725 South Nardo Avenue, Solana Beach, CA 92075 (US). LIVINGSTON, Brian; 13555 Chaco Court, San Diego, CA 92129 (US). CHESNUT, Robert, W.; 1473 Kings Cross Drive, Cardiff-by-the-Sea, CA 92007 (US).				
(74) Agents: PARENT, Annette, S. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111-3834 (US).				
(54) Title: EXPRESSION VECTORS FOR STIMULATING AN IMMUNE RESPONSE AND METHODS OF USING THE SAME				
(57) Abstract				
<p>The present invention relates to nucleic acid vaccines encoding multiple CTL and HTL epitopes and MHC targeting sequences.</p>				

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NZ	New Zealand	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	PL	Poland	ZW	Zimbabwe
CI	Cote d'Ivoire	KP	Democratic People's Republic of Korea	PT	Portugal		
CM	Cameroon	CR	Republic of Costa Rica	RO	Romania		
CN	China	CZ	Czech Republic	RU	Russian Federation		
CU	Cuba	LC	Liechtenstein	SD	Sudan		
DE	Germany	LI	Lithuania	SE	Sweden		
DK	Denmark	LK	Sri Lanka	SG	Singapore		
EE	Estonia	LR	Liberia				

## EXPRESSION VECTORS FOR STIMULATING AN IMMUNE RESPONSE AND METHODS OF USING THE SAME

### CROSS-REFERENCES TO RELATED APPLICATIONS

5 This application claims the benefit of 09/078,904, filed May 13, 1998, and  
60/085,751, filed May 15, 1998, both herein incorporated by reference in their entirety.

### STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

10 This invention was made with government support under NIH Grant No.  
AI-42699-01, NIH Grant No. AI38584-03, and NIH Contract No. N01-AI-45241. The  
Government has certain rights in this invention.

### FIELD OF THE INVENTION

15 The present invention relates to nucleic acid vaccines encoding multiple  
CTL and HTL epitopes and MHC targeting sequences.

### BACKGROUND OF THE INVENTION

20 Vaccines are of fundamental importance in modern medicine and have  
been highly effective in combating certain human diseases. However, despite the  
successful implementation of vaccination programs that have greatly limited or virtually  
eliminated several debilitating human diseases, there are a number of diseases that affect  
millions worldwide for which effective vaccines have not been developed.

Major advances in the field of immunology have led to a greater  
25 understanding of the mechanisms involved in the immune response and have provided  
insights into developing new vaccine strategies (Kuby, *Immunology*, 443-457 (3rd ed.,  
1997), which is incorporated herein by reference). These new vaccine strategies have  
taken advantage of knowledge gained regarding the mechanisms by which foreign  
material, termed antigen, is recognized by the immune system and eliminated from the  
30 organism. An effective vaccine is one that elicits an immune response to an antigen of  
interest.

Specialized cells of the immune system are responsible for the protective  
activity required to combat diseases. An immune response involves two major groups of  
cells, lymphocytes, or white blood cells, and antigen-presenting cells. The purpose of

these immune response cells is to recognize foreign material, such as an infectious organism or a cancer cell, and remove that foreign material from the organism.

Two major types of lymphocytes mediate different aspects of the immune response. B cells display on their cell surface specialized proteins, called antibodies, that bind specifically to foreign material, called antigens. Effector B cells produce soluble forms of the antibody, which circulate throughout the body and function to eliminate antigen from the organism. This branch of the immune system is known as the humoral branch. Memory B cells function to recognize the antigen in future encounters by continuing to express the membrane-bound form of the antibody.

A second major type of lymphocyte is the T cell. T cells also have on their cell surface specialized proteins that recognize antigen but, in contrast to B cells, require that the antigen be bound to a specialized membrane protein complex, the major histocompatibility complex (MHC), on the surface of an antigen-presenting cell. Two major classes of T cells, termed helper T lymphocytes ("HTL") and cytotoxic T lymphocytes ("CTL"), are often distinguished based on the presence of either CD4 or CD8 protein, respectively, on the cell surface. This branch of the immune system is known as the cell-mediated branch.

The second major class of immune response cells are cells that function in antigen presentation by processing antigen for binding to MHC molecules expressed in the antigen presenting cells. The processed antigen bound to MHC molecules is transferred to the surface of the cell, where the antigen-MHC complex is available to bind to T cells.

MHC molecules can be divided into MHC class I and class II molecules and are recognized by the two classes of T cells. Nearly all cells express MHC class I molecules, which function to present antigen to cytotoxic T lymphocytes. Cytotoxic T lymphocytes typically recognize antigen bound to MHC class I. A subset of cells called antigen-presenting cells express MHC class II molecules. Helper T lymphocytes typically recognize antigen bound to MHC class II molecules. Antigen-presenting cells include dendritic cells, macrophages, B cells, fibroblasts, glial cells, pancreatic beta cells, thymic epithelial cells, thyroid epithelial cells and vascular endothelial cells. These antigen-presenting cells generally express both MHC class I and class II molecules. Also, B cells function as both antibody-producing and antigen-presenting cells.

Once a helper T lymphocyte recognizes an antigen-MHC class II complex on the surface of an antigen-presenting cell, the helper T lymphocyte becomes activated

and produces growth factors that activate a variety of cells involved in the immune response, including B cells and cytotoxic T lymphocytes. For example, under the influence of growth factors expressed by activated helper T lymphocytes, a cytotoxic T lymphocyte that recognizes an antigen-MHC class I complex becomes activated. CTLs monitor and eliminate cells that display antigen specifically recognized by the CTL, such as infected cells or tumor cells. Thus, activation of helper T lymphocytes stimulates the activation of both the humoral and cell-mediated branches of the immune system.

An important aspect of the immune response, in particular as it relates to vaccine efficacy, is the manner in which antigen is processed so that it can be recognized by the specialized cells of the immune system. Distinct antigen processing and presentation pathways are utilized. The one is a cytosolic pathway, which results in the antigen being bound to MHC class I molecules. An alternative pathway is an endoplasmic reticulum pathway, which bypasses the cytosol. Another is an endocytic pathway, which results in the antigen being bound to MHC class II molecules. Thus, the cell surface presentation of a particular antigen by a MHC class II or class I molecule to a helper T lymphocyte or a cytotoxic T lymphocyte, respectively, is dependent on the processing pathway for that antigen.

The cytosolic pathway processes endogenous antigens that are expressed inside the cell. The antigen is degraded by a specialized protease complex in the cytosol of the cell, and the resulting antigen peptides are transported into the endoplasmic reticulum, an organelle that processes cell surface molecules. In the endoplasmic reticulum, the antigen peptides bind to MHC class I molecules, which are then transported to the cell surface for presentation to cytotoxic T lymphocytes of the immune system.

Antigens that exist outside the cell are processed by the endocytic pathway. Such antigens are taken into the cell by endocytosis, which brings the antigens into specialized vesicles called endosomes and subsequently to specialized vesicles called lysosomes, where the antigen is degraded by proteases into antigen peptides that bind to MHC class II molecules. The antigen peptide-MHC class II molecule complex is then transported to the cell surface for presentation to helper T lymphocytes of the immune system.

A variety of factors must be considered in the development of an effective vaccine. For example, the extent of activation of either the humoral or cell-mediated branch of the immune system can determine the effectiveness of a vaccine against a

particular disease. Furthermore, the development of immunologic memory by inducing memory-cell formation can be important for an effective vaccine against a particular disease (Kuby, *supra*). For example, protection from infectious diseases caused by pathogens with short incubation periods, such as influenza virus, requires high levels of neutralizing antibody generated by the humoral branch because disease symptoms are already underway before memory cells are activated. Alternatively, protection from infectious diseases caused by pathogens with long incubation periods, such as polio virus, does not require neutralizing antibodies at the time of infection but instead requires memory B cells that can generate neutralizing antibodies to combat the pathogen before it is able to infect target tissues. Therefore, the effectiveness of a vaccine at preventing or ameliorating the symptoms of a particular disease depends on the type of immune response generated by the vaccine.

Many traditional vaccines have relied on intact pathogens such as attenuated or inactivated viruses or bacteria to elicit an immune response. However, these traditional vaccines have advantages and disadvantages, including reversion of an attenuated pathogen to a virulent form. The problem of reversion of an attenuated vaccine has been addressed by the use of molecules of the pathogen rather than the whole pathogen. For example, immunization approaches have begun to incorporate recombinant vector vaccines and synthetic peptide vaccines (Kuby, *supra*). Recently, DNA vaccines have also been used (Donnelly *et al.*, *Annu. Rev. Immunol.* 15:617-648 (1997), which is incorporated herein by reference). The use of molecules of a pathogen provides safe vaccines that circumvent the potential for reversion to a virulent form of the vaccine.

The targeting of antigens to MHC class II molecules to activate helper T lymphocytes has been described using lysosomal targeting sequences, which direct antigens to lysosomes, where the antigen is digested by lysosomal proteases into antigen peptides that bind to MHC class II molecules (U.S. Patent No. 5,633,234; Thomson *et al.*, *J. Virol.* 72:2246-2252 (1998)). It would be advantageous to develop vaccines that deliver multiple antigens while exploiting the safety provided by administering individual epitopes of a pathogen rather than a whole organism. In particular, it would be advantageous to develop vaccines that effectively target antigens to MHC class II molecules for activation of helper T lymphocytes.

Several studies also point to the crucial role of cytotoxic T cells in both production and eradication of infectious diseases and cancer by the immune system

(Byrne *et al.*, *J. Immunol.* 51:682 (1984); McMichael *et al.*, *N. Engl. J. Med.* 309:13 (1983)). Recombinant protein vaccines do not reliably induce CTL responses, and the use of otherwise immunogenic vaccines consisting of attenuated pathogens in humans is hampered, in the case of several important diseases, by overriding safety concerns. In the case of diseases such as HIV, HBV, HCV, and malaria, it appears desirable not only to induce a vigorous CTL response, but also to focus the response against highly conserved epitopes in order to prevent escape by mutation and overcome variable vaccine efficacy against different isolates of the target pathogen.

- Induction of a broad response directed simultaneously against multiple epitopes also appears to be crucial for development of efficacious vaccines. HIV infection is perhaps the best example where an infected host may benefit from a multispecific response. Rapid progression of HIV infection has been reported in cases where a narrowly focused CTL response is induced whereas nonprogressors tend to show a broader specificity of CTLs (Goulder *et al.*, *Nat. Med.* 3:212 (1997); Borrow *et al.*, *Nat. Med.* 3:205 (1997)). The highly variable nature of HIV CTL epitopes resulting from a highly mutating genome and selection by CTL responses directed against only a single or few epitopes also supports the need for broad epitope CTL responses (McMichael *et al.*, *Annu. Rev. Immunol.* 15:271 (1997)).

- One potential approach to induce multispecific responses against conserved epitopes is immunization with a minigene plasmid encoding the epitopes in a string-of-beads fashion. Induction of CTL, HTL, and B cell responses in mice by minigene plasmids have been described by several laboratories using constructs encoding as many as 11 epitopes (An *et al.*, *J. Virol.* 71:2292 (1997); Thomson *et al.*, *J. Immunol.* 157:822 (1996); Whitton *et al.*, *J. Virol.* 67:348 (1993); Hanke *et al.*, *Vaccine* 16:426 (1998); Vitiello *et al.*, *Eur. J. Immunol.* 27:671-678 (1997)). Minigenes have been delivered *in vivo* by infection with recombinant adenovirus or vaccinia, or by injection of purified DNA via the intramuscular or intradermal route (Thomson *et al.*, *J. Immunol.* 160:1717 (1998); Toes *et al.*, *Proc. Natl. Acad. Sci. USA* 94:14660 (1997)).

- Successful development of minigene DNA vaccines for human use will require addressing certain fundamental questions dealing with epitope MHC affinity, optimization of constructs for maximum *in vivo* immunogenicity, and development of assays for testing *in vivo* potency of multi-epitope minigene constructs. Regarding MHC binding affinity of epitopes, it is not currently known whether both high and low affinity epitopes can be included within a single minigene construct, and what ranges of peptide

affinity are permissible for CTL induction *in vivo*. This is especially important because dominant epitopes can vary in their affinity and because it might be important to be able to deliver mixtures of dominant and subdominant epitopes that are characterized by high and low MHC binding affinities.

- 5           With respect to minigene construct optimization for maximum immunogenicity *in vivo*, conflicting data exists regarding whether the exact position of the epitopes in a given construct or the presence of flanking regions, helper T cell epitopes, and signal sequences might be crucial for CTL induction (Del Val *et al.*, *Cell* 66:1145 (1991); Bergmann *et al.*, *J. Virol.* 68:5306 (1994); Thomson *et al.*, *Proc. Natl. Acad. Sci. USA* 92:5845 (1995); Shirai *et al.*, *J. Infect. Dis.* 173:24 (1996); Rahemtulla *et al.*, *Nature* 353:180 (1991); Jennings *et al.*, *Cell. Immunol.* 133:234 (1991); Anderson *et al.*, *J. Exp. Med.* 174:489 (1991); Uger *et al.*, *J. Immunol.* 158:685 (1997)). Finally, regarding development of assays that allow testing of human vaccine candidates, it should be noted that, to date, all *in vivo* immunogenicity data of multi-epitope minigene plasmids
- 10           have been performed with murine class I MHC-restricted epitopes. It would be advantageous to be able to test the *in vivo* immunogenicity of minigenes containing human CTL epitopes in a convenient animal model system.

- 15           Thus, there exists a need to develop methods to effectively deliver a variety of HTL (helper T lymphocyte) and CTL (cytotoxic T lymphocyte) antigens to stimulate an immune response. The present invention satisfies this need and provides related advantages as well.
- 20

#### SUMMARY OF THE INVENTION

- The invention therefore provides expression vectors encoding two or more
- 25   HTL epitopes fused to a MHC class II targeting sequence, as well as expression vectors encoding a CTL epitope and a universal HTL epitope fused to an MHC class I targeting sequence. The HTL epitope can be a universal HTL epitope (also referred to as a universal MHC class II epitope). The invention also provides expression vectors encoding two or more HTL epitopes fused to a MHC class II targeting sequence and
- 30   encoding one or more CTL epitopes. The invention additionally provides methods of stimulating an immune response by administering an expression vector of the invention *in vivo*, as well as methods of assaying the human immunogenicity of a human T cell peptide epitope *in vivo* in a non-human mammal.



In one aspect, the present invention provides an expression vector comprising a promoter operably linked to a first nucleotide sequence encoding a major histocompatibility (MHC) targeting sequence fused to a second nucleotide sequence encoding two or more heterologous peptide epitopes, wherein the heterologous peptide epitopes comprise two HTL peptide epitopes or a CTL peptide epitope and a universal HTL peptide epitope.

In another aspect, the present invention provides a method of inducing an immune response *in vivo* comprising administering to a mammalian subject an expression vector comprising a promoter operably linked to a first nucleotide sequence encoding a major histocompatibility (MHC) targeting sequence fused to a second nucleotide sequence encoding two or more heterologous peptide epitopes, wherein the heterologous peptide epitopes comprise two HTL peptide epitopes or a CTL peptide epitope and a universal HTL peptide epitope.

In another aspect, the present invention provides a method of inducing an immune response *in vivo* comprising administering to a mammalian subject an expression vector comprising a promoter operably linked to a first nucleotide sequence encoding a major histocompatibility (MHC) targeting sequence fused to a second nucleotide sequence encoding a heterologous human HTL peptide epitope.

In another aspect, the present invention provides a method of assaying the human immunogenicity of a human T cell peptide epitope *in vivo* in a non-human mammal, comprising the step of administering to the non-human mammal an expression vector comprising a promoter operably linked to a first nucleotide sequence encoding a heterologous human CTL or HTL peptide epitope.

In one embodiment, the heterologous peptide epitopes comprise two or more heterologous HTL peptide epitopes. In another embodiment, the heterologous peptide epitopes comprise a CTL peptide epitope and a universal HTL peptide epitope. In another embodiment, the heterologous peptide epitopes further comprise one to two or more heterologous CTL peptide epitopes. In another embodiment, the expression vector comprises both HTL and CTL peptide epitopes.

In one embodiment, one of the HTL peptide epitopes is a universal HTL epitope. In another embodiment, the universal HTL epitope is a pan DR epitope. In another embodiment, the pan DR epitope has the sequence  
AlaLysPheValAlaAlaTrpThrLeuLysAlaAlaAla (SEQ ID NO:38).

In one embodiment, the peptide epitopes are hepatitis B virus epitopes, hepatitis C virus epitopes, human immunodeficiency virus epitopes, human papilloma virus epitopes, MAGE epitopes, PSA epitopes, PSM epitopes, PAP epitopes, p53 epitopes, CEA epitopes, Her2/neu epitopes, or *Plasmodium* epitopes. In another embodiment, the peptide epitopes each have a sequence selected from the group consisting of the peptides depicted in Tables 1-8. In another embodiment, at least one of the peptide epitopes is an analog of a peptide depicted in Tables 1-8.

In one embodiment, the MHC targeting sequence comprises a region of a polypeptide selected from the group consisting of the Ii protein, LAMP-I, HLS-DM, HLA-DO, H2-DO, influenza matrix protein, hepatitis B surface antigen, hepatitis B virus core antigen, Ty particle, Ig- $\alpha$  protein, Ig- $\beta$  protein, and Ig kappa chain signal sequence.

In one embodiment, the expression vector further comprises a second promoter sequence operably linked to a third nucleotide sequence encoding one or more heterologous HTL or CTL peptide epitopes. In another embodiment, the CTL peptide epitope comprises a structural motif for an HLA supertype, whereby the peptide CTL epitope binds to two or more members of the supertype with an affinity of greater than 500 nM. In another embodiment, the CTL peptide epitopes have structural motifs that provide binding affinity for more than one HLA allele supertype.

In one embodiment, the non-human mammal is a transgenic mouse that expresses a human HLA allele. In another embodiment, the human HLA allele is selected from the group consisting of A11 and A2.1. In another embodiment, the non-human mammal is a macaque that expresses a human HLA allele.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide and amino acid sequences (SEQ ID NOS:1 and 2, respectively) of the IiPADRE construct encoding a fusion of the murine Ii gene with a pan DR epitope sequence substituted for the CLIP sequence of the Ii protein.

Figure 2 shows the nucleotide and amino acid sequences (SEQ ID NOS:3 and 4, respectively) of the I80T construct encoding a fusion of the cytoplasmic domain, the transmembrane domain and part of the luminal domain of the Ii protein fused to multiple MHC class II epitopes.

Figure 3 shows the nucleotide and amino acid sequences (SEQ ID NOS:5 and 6, respectively) of the IiThfull construct encoding a fusion of the cytoplasmic domain, transmembrane domain and a portion of the luminal domain of the Ii protein

fused to multiple T helper epitopes and amino acid residues 101 to 215 of the Ii protein, which encodes the trimerization region of the Ii protein.

Figure 4 shows the nucleotide and amino acid sequences (SEQ ID NOS:7 and 8, respectively) of the KappaLAMP-Th construct encoding a fusion of the murine immunoglobulin kappa signal sequence fused to multiple T helper epitopes and the transmembrane and cytoplasmic domains of LAMP-1.

Figure 5 shows the nucleotide and amino acid sequences (SEQ ID NOS:9 and 10, respectively) of the H2M-Th construct encoding a fusion of the signal sequence of H2-M fused to multiple MHC class II epitopes and the transmembrane and cytoplasmic domains of H2-M.

Figure 6 shows the nucleotide and amino acid sequences (SEQ ID NOS:11 and 12, respectively) of the H2O-Th construct encoding a fusion of the signal sequence of H2-DO fused to multiple MHC class II epitopes and the transmembrane and cytoplasmic domains of H2-DO.

Figure 7 shows the nucleotide and amino acid sequences (SEQ ID NOS:13 and 14, respectively) of the PADRE-Influenza matrix construct encoding a fusion of a pan DR epitope sequence fused to the amino-terminus of influenza matrix protein.

Figure 8 shows the nucleotide and amino acid sequences (SEQ ID NOS:15 and 16, respectively) of the PADRE-HBV-s construct encoding a fusion of a pan DR epitope sequence fused to the amino-terminus of hepatitis B virus surface antigen.

Figure 9 shows the nucleotide and amino acid sequences (SEQ ID NOS:17 and 18, respectively) of the Ig-alphaTh construct encoding a fusion of the signal sequence of the Ig- $\alpha$  protein fused to multiple MHC class II epitopes and the transmembrane and cytoplasmic domains of the Ig- $\alpha$  protein.

Figure 10 shows the nucleotide and amino acid sequences (SEQ ID NOS:19 and 20, respectively) of the Ig-betaTh construct encoding a fusion of the signal sequence of the Ig- $\beta$  protein fused to multiple MHC class II epitopes and the transmembrane and cytoplasmic domains of the Ig- $\beta$  protein.

Figure 11 shows the nucleotide and amino acid sequences (SEQ ID NOS:21 and 22, respectively) of the SigTh construct encoding a fusion of the signal sequence of the kappa immunoglobulin fused to multiple MHC class II epitopes.

Figure 12 shows the nucleotide and amino acid sequences (SEQ ID NOS:23 and 24, respectively) of human HLA-DR, the invariant chain (Ii) protein.

Figure 13 shows the nucleotide and amino acid sequences (SEQ ID NOS:25 and 26, respectively) of human lysosomal membrane glycoprotein-1 (LAMP-1).

Figure 14 shows the nucleotide and amino acid sequences (SEQ ID NOS:27 and 28, respectively) of human HLA-DMB.

5 Figure 15 shows the nucleotide and amino acid sequences (SEQ ID NOS:29 and 30, respectively) of human HLA-DO beta.

Figure 16 shows the nucleotide and amino acid sequences (SEQ ID NOS:31 and 32, respectively) of the human MB-1 Ig- $\alpha$ .

10 Figure 17 shows the nucleotide and amino acid sequences (SEQ ID NOS:33 and 34, respectively) of human Ig- $\beta$  protein.

Figure 18 shows a schematic diagram depicting the method of generating some of the constructs encoding a MHC class II targeting sequence fused to multiple MHC class II epitopes.

15 Figure 19 shows the nucleotide sequence of the vector pEP2 (SEQ ID NO:35).

Figure 20 shows the nucleotide sequence of the vector pMIN.0 (SEQ ID NO:36).

Figure 21 shows the nucleotide sequence of the vector pMIN.1 (SEQ ID NO:37).

20 Figure 22. Representative CTL responses in HLA-A2.1/K<sup>b</sup>-H-2<sup>bbs</sup> mice immunized with pMin.1 DNA. Splenocytes from primed animals were cultured in triplicate flasks and stimulated twice *in vitro* with each peptide epitope. Cytotoxicity of each culture was assayed in a <sup>51</sup>Cr release assay against Jurkat-A2.1/K<sup>b</sup> target cells in the presence (filled symbols, solid lines) or absence (open symbols, dotted lines) of peptide.

25 Each symbol represents the response of a single culture.

Figure 23. Presentation of viral epitopes to specific CTLs by Jurkat-A2.1/K<sup>b</sup> tumor cells transfected with DNA minigene. Two constructs were used for transfection, pMin.1 and pMin.2-GFP. pMin.2-GFP-transfected targets cells were sorted by FACS and the population used in this experiment contained 60% fluorescent cells.

30 CTL stimulation was measured by quantitating the amount of IFN- $\gamma$  release (A, B) or by lysis of <sup>51</sup>Cr-labeled target cells (C, D, hatched bars). CTLs were stimulated with transfected cells (A, C) or with parental Jurkat-A2.1/K<sup>b</sup> cells in the presence of 1  $\mu$ g/ml peptide (B, D). Levels of IFN- $\gamma$  release and cytotoxicity for the different CTL lines in the absence of epitope ranged from 72-126 pg/ml and 2-6% respectively.

Figure 24. Summary of modified minigene constructs used to address variables critical for *in vivo* immunogenicity. The following modifications were incorporated into the prototype pMin.1 construct; A, deletion of PADRE HTL epitope; B, incorporation of the native HBV Pol 551 epitope that contains an alanine in position 9; C, deletion of the Ig kappa signal sequence; and D, switching position of the HBV Env 335 and HBV Pol 455 epitopes.

Figure 25. Examination of variables that may influence pMin.1 immunogenicity. *In vivo* CTL-inducing activity of pMin.1 is compared to modified constructs. For ease of comparison, the CTL response induced by each of the modified DNA minigene constructs (shaded bars) is compared separately in each of the four panels to the response induced by the prototype pMin.1 construct (solid bars). The geometric mean response of CTL-positive cultures from two to five independent experiments are shown. Numbers shown with each bar indicate the number of positive cultures/total number tested for that particular epitope. The ratio of positive cultures/total tested for the pMin.1 group is shown in panel A and is the same for the remaining Figure panels (see Example V, Materials and Methods, *in vitro* CTL cultures, for the definition of a positive CTL culture). Theradigm responses were obtained by immunizing animals with the lipopeptide and stimulating and testing splenocyte cultures with the HBV Core 18-27 peptide.

#### DEFINITIONS

An "HTL" peptide epitope or an "MHC II epitope" is an MHC class II restricted epitope, i.e., one that is bound by an MHC class II molecule.

A "CTL" peptide epitope or an "MHC I epitope" is an MHC class I restricted epitope, i.e., one that is bound by an MHC class I molecule.

An "MHC targeting sequence" refers to a peptide sequence that targets a polypeptide, e.g., comprising a peptide epitope, to a cytosolic pathway (e.g., an MHC class I antigen processing pathway), an endoplasmic reticulum pathway, or an endocytic pathway (e.g., an MHC class II antigen processing pathway).

The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a

coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature, e.g., a fusion polypeptide comprising subsequence from different polypeptides, peptide epitopes from the same polypeptide that are not naturally in an adjacent position, or repeats of a single peptide epitope.

- As used herein, the term "universal MHC class II epitope" or a "universal HTL epitope" refers to a MHC class II peptide epitope that binds to gene products of multiple MHC class II alleles. For example, the DR, DP and DQ alleles are human MHC class II alleles. Generally, a unique set of peptides binds to a particular gene product of a MHC class II allele. In contrast, a universal MHC class II epitope is able to bind to gene products of multiple MHC class II alleles. A universal MHC class II epitope binds to 2 or more MHC class II alleles, generally 3 or more MHC class II alleles, and particularly 5 or more MHC class II alleles. Thus, the presence of a universal MHC class II epitope in an expression vector is advantageous in that it functions to increase the number of allelic MHC class II molecules that can bind to the peptide and, consequently, the number of Helper T lymphocytes that are activated.

- Universal MHC class II epitopes are well known in the art and include, for example, epitopes such as the "pan DR epitopes," also referred to as "PADRE" (Alexander *et al.*, *Immunity* 1:751-761 (1994); WO 95/07707, USSN 60/036,713, USSN 60/037,432, PCT/US98/01373, 09/009,953, and USSN 60/087,192 each of which is incorporated herein by reference). A "pan DR binding peptide" or a "PADRE" peptide of the invention is a peptide capable of binding at least about 7 different DR molecules, preferably 7 of the 12 most common DR molecules, most preferably 9 of the 12 most common DR molecules (DR1, 2w2b, 2w2a, 3, 4w4, 4w14, 5, 7, 52a, 52b, 52c, and 53), or alternatively, 50% of a panel of DR molecules representative of greater than or equal to 75% of the human population, preferably greater than or equal to 80% of the human population. Pan DR epitopes can bind to a number of DR alleles and are strongly immunogenic for T cells. For example, pan DR epitopes were found to be more effective at inducing an immune response than natural MHC class II epitopes (Alexander, *supra*).
- An example of a PADRE epitope is the peptide AlaLysPheValAlaAlaTrpThrLeuLysAlaAlaAla (SEQ ID NO:38) (for additional examples of PADRE epitopes, see Table 8 of TTC docket No. 018623-006221, filed May 12, 1999, USSN \_\_\_\_\_, herein incorporated by reference in its entirety).

With regard to a particular amino acid sequence, an "epitope" is a set of amino acid residues which is involved in recognition by a particular immunoglobulin, or in the context of T cells, those residues necessary for recognition by T cell receptor proteins and/or Major Histocompatibility Complex (MHC) receptors. In an immune system setting, *in vivo* or *in vitro*, an epitope is the collective features of a molecule, such as primary, secondary and tertiary peptide structure, and charge, that together form a site recognized by an immunoglobulin, T cell receptor or HLA molecule. Throughout this disclosure epitope and peptide are often used interchangeably. It is to be appreciated, however, that isolated or purified protein or peptide molecules larger than and comprising an epitope of the invention are still within the bounds of the invention.

As used herein, "high affinity" with respect to HLA class I molecules is defined as binding with an IC<sub>50</sub> (or K<sub>D</sub>) of less than 50 nM. "Intermediate affinity" is binding with an IC<sub>50</sub> (or K<sub>D</sub>) of between about 50 and about 500 nM. "High affinity" with respect to binding to HLA class II molecules is defined as binding with an K<sub>D</sub> of less than 100 nM. "Intermediate affinity" is binding with a K<sub>D</sub> of between about 100 and about 1000 nM. Assays for determining binding are described in detail, e.g., in PCT publications WO 94/20127 and WO 94/03205. Alternatively, binding is expressed relative to a reference peptide. As a particular assay becomes more, or less, sensitive, the IC<sub>50</sub>s of the peptides tested may change somewhat. However, the binding relative to the reference peptide will not significantly change. For example, in an assay run under conditions such that the IC<sub>50</sub> of the reference peptide increases 10-fold, the IC<sub>50</sub> values of the test peptides will also shift approximately 10-fold. Therefore, to avoid ambiguities, the assessment of whether a peptide is a good, intermediate, weak, or negative binder is generally based on its IC<sub>50</sub>, relative to the IC<sub>50</sub> of a standard peptide.

Throughout this disclosure, results are expressed in terms of "IC<sub>50</sub>s." IC<sub>50</sub> is the concentration of peptide in a binding assay at which 50% inhibition of binding of a reference peptide is observed. Given the conditions in which the assays are run (i.e., limiting HLA proteins and labeled peptide concentrations), these values approximate K<sub>D</sub> values. It should be noted that IC<sub>50</sub> values can change, often dramatically, if the assay conditions are varied, and depending on the particular reagents used (e.g., HLA preparation, etc.). For example, excessive concentrations of HLA molecules will increase the apparent measured IC<sub>50</sub> of a given ligand.

The terms "identical" or percent "identity," in the context of two or more peptide sequences, refer to two or more sequences or subsequences that are the same or

have a specified percentage of amino acid residues that are the same, when compared and aligned for maximum correspondence over a comparison window, as measured using a sequence comparison algorithms using default program parameters or by manual alignment and visual inspection.

- 5           The phrases "isolated" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany the material as it is found in its native state. Thus, isolated peptides in accordance with the invention preferably do not contain materials normally associated with the peptides in their in situ environment.

- 10           "Major histocompatibility complex" or "MHC" is a cluster of genes that plays a role in control of the cellular interactions responsible for physiologic immune responses. In humans, the MHC complex is also known as the HLA complex. For a detailed description of the MHC and HLA complexes, see Paul, *Fundamental Immunology* (3rd ed. 1993).

- 15           "Human leukocyte antigen" or "HLA" is a human class I or class II major histocompatibility complex (MHC) protein (see, e.g., Stites, et al., *Immunology*, (8th ed., 1994).

- An "HLA supertype or family", as used herein, describes sets of HLA molecules grouped on the basis of shared peptide-binding specificities. HLA class I  
20           molecules that share somewhat similar binding affinity for peptides bearing certain amino acid motifs are grouped into HLA superfamily, HLA supertype family, HLA family, and HLA xx-like supertype molecules (where xx denotes a particular HLA type), are synonyms.

- The term "motif" refers to the pattern of residues in a peptide of defined  
25           length, usually a peptide of from about 8 to about 13 amino acids for a class I HLA motif and from about 6 to about 25 amino acids for a class II HLA motif, which is recognized by a particular HLA molecule. Peptide motifs are typically different for each protein encoded by each human HLA allele and differ in the pattern of the primary and secondary anchor residues.

- 30           A "supermotif" is a peptide binding specificity shared by HLA molecules encoded by two or more HLA alleles. Thus, a preferably is recognized with high or intermediate affinity (as defined herein) by two or more HLA antigens.

          "Cross-reactive binding" indicates that a peptide is bound by more than one HLA molecule; a synonym is degenerate binding.



The term "peptide" is used interchangeably with "oligopeptide" in the present specification to designate a series of residues, typically L-amino acids, connected one to the other, typically by peptide bonds between the  $\alpha$ -amino and carboxyl groups of adjacent amino acids. The preferred CTL-inducing oligopeptides of the invention are 13  
5 residues or less in length and usually consist of between about 8 and about 11 residues, preferably 9 or 10 residues. The preferred HTL-inducing oligopeptides are less than about 50 residues in length and usually consist of between about 6 and about 30 residues, more usually between about 12 and 25, and often between about 15 and 20 residues.

An "immunogenic peptide" or "peptide epitope" is a peptide which  
10 comprises an allele-specific motif or supermotif such that the peptide will bind an HLA molecule and induce a CTL and/or HTL response. Thus, immunogenic peptides of the invention are capable of binding to an appropriate HLA molecule and thereafter inducing a cytotoxic T cell response, or a helper T cell response, to the antigen from which the immunogenic peptide is derived.

15 A "protective immune response" refers to a CTL and/or an HTL response to an antigen derived from an infectious agent or a tumor antigen, which prevents or at least partially arrests disease symptoms or progression. The immune response may also include an antibody response which has been facilitated by the stimulation of helper T cells.

20 The term "residue" refers to an amino acid or amino acid mimetic incorporated into an oligopeptide by an amide bond or amide bond mimetic.

"Synthetic peptide" refers to a peptide that is not naturally occurring, but is man-made using such methods as chemical synthesis or recombinant DNA technology.

The nomenclature used to describe peptide compounds follows the  
25 conventional practice wherein the amino group is presented to the left (the N-terminus) and the carboxyl group to the right (the C-terminus) of each amino acid residue. When amino acid residue positions are referred to in a peptide epitope they are numbered in an amino to carboxyl direction with position one being the position closest to the amino terminal end of the epitope, or the peptide or protein of which it may be a part. In the  
30 formulae representing selected specific embodiments of the present invention, the amino- and carboxyl-terminal groups, although not specifically shown, are in the form they would assume at physiologic pH values, unless otherwise specified. In the amino acid structure formulae, each residue is generally represented by standard three letter or single letter designations. The L-form of an amino acid residue is represented by a capital single

letter or a capital first letter of a three-letter symbol, and the D-form for those amino acids having D-forms is represented by a lower case single letter or a lower case three letter symbol. Glycine has no asymmetric carbon atom and is simply referred to as "Gly" or G.

- As used herein, the term "expression vector" is intended to refer to a
- 5 nucleic acid molecule capable of expressing an antigen of interest such as a MHC class I or class II epitope in an appropriate target cell. An expression vector can be, for example, a plasmid or virus, including DNA or RNA viruses. The expression vector contains such a promoter element to express an antigen of interest in the appropriate cell or tissue in order to stimulate a desired immune response.

10

#### DETAILED DESCRIPTION OF THE INVENTION

- Cytotoxic T lymphocytes (CTLs) and helper T lymphocytes (HTLs) are critical for immunity against infectious pathogens; such as viruses, bacteria, and protozoa; tumor cells; autoimmune diseases and the like. The present invention provides
- 15 minigenes that encode peptide epitopes which induce a CTL and/or HTL response. The minigenes of the invention also include an MHC targeting sequence. A variety of minigenes encoding different epitopes can be tested for immunogenicity using an HLA transgenic mouse. The epitopes are typically a combination of at least two or more HTL epitopes, or a CTL epitope plus a universal HTL epitope, and optionally include additional
- 20 HTL and/or CTL epitopes. Two, three, four, five, six, seven, eight, nine, ten, twenty, thirty, forty or about fifty different epitopes, either HTL and/or CTL, can be included in the minigene, along with the MHC targeting sequence. The epitopes can have different HLA restriction. Epitopes to be tested include those derived from viruses such as HIV, HBV, HCV, HSV, CMV, HPV, and HTLV; cancer antigens such as p53, Her2/Neu,
- 25 MAGE, PSA, human papilloma virus, and CEA; parasites such as *Trypanosoma*, *Plasmodium*, *Leishmania*, *Giardia*, *Entamoeba*; autoimmune diseases such as rheumatoid arthritis, myesthenia gravis, and lupus erythematosus; fungi such as *Aspergillus* and *Candida*; and bacteria such as *Escherichia coli*, *Staphylococci*, *Chlamydia*, *Mycobacteria*, *Streptococci*, and *Pseudomonas*. The epitopes to be encoded by the minigene are selected
- 30 and tested using the methods described in published PCT applications WO 93/07421, WO 94/02353, WO 95/01000, WO 97/04451, and WO 97/05348, herein incorporated by reference.

### HTL and CTL Epitopes

The expression vectors of the invention encode one or more MHC class II and/or class I epitopes and an MHC targeting sequence. Multiple MHC class II or class I epitopes present in an expression vector can be derived from the same antigen, or the

5 MHC epitopes can be derived from different antigens. For example, an expression vector can contain one or more MHC epitopes that can be derived from two different antigens of the same virus or from two different antigens of different viruses. Furthermore, any MHC epitope can be used in the expression vectors of the invention. For example, any

10 single MHC epitope or a combination of the MHC epitopes shown in Tables 1 to 8 can be used in the expression vectors of the invention. Other peptide epitopes can be selected by one of skill in the art, e.g., by using a computer to select epitopes that contain HLA allele-specific motifs or supermotifs. The expression vectors of the invention can also encode one or more universal MHC class II epitopes, e.g., PADRE (*see, e.g.*, SEQ ID NO:38 and Table 8 of TTC docket No. 018623-006221, filed May 12, 1999, USSN

15 \_\_\_\_\_).

Universal MHC class II epitopes can be advantageously combined with other MHC class I and class II epitopes to increase the number of cells that are activated in response to a given antigen and provide broader population coverage of MHC-reactive alleles. Thus, the expression vectors of the invention can encode MHC epitopes specific

20 for an antigen, universal MHC class II epitopes, or a combination of specific MHC epitopes and at least one universal MHC class II epitope.

MHC class I epitopes are generally about 5 to 15 amino acids in length, in particular about 8 to 11 amino acids in length. MHC class II epitopes are generally about 10 to 25 amino acids in length, in particular about 13 to 21 amino acids in length. A

25 MHC class I or II epitope can be derived from any desired antigen of interest. The antigen of interest can be a viral antigen, surface receptor, tumor antigen, oncogene, enzyme, or any pathogen, cell or molecule for which an immune response is desired. Epitopes can be selected based on their ability to bind one or multiple HLA alleles, and can also be selected using the "analog" technique described below.

30

### Targeting Sequences

The expression vectors of the invention encode one or more MHC epitopes operably linked to a MHC targeting sequence. The use of a MHC targeting sequence enhances the immune response to an antigen, relative to delivery of antigen alone, by

directing the peptide epitope to the site of MHC molecule assembly and transport to the cell surface, thereby providing an increased number of MHC molecule-peptide epitope complexes available for binding to and activation of T cells.

- MHC class I targeting sequences are used in the present invention, e.g.,
- 5 those sequences that target an MHC class I epitope peptide to a cytosolic pathway or to the endoplasmic reticulum (*see, e.g., Rammensee et al., Immunogenetics* 41:178-228 (1995)). For example, the cytosolic pathway processes endogenous antigens that are expressed inside the cell. Although not wishing to be bound by any particular theory, cytosolic proteins are thought to be at least partially degraded by an endopeptidase
- 10 activity of a proteasome and then transported to the endoplasmic reticulum by the TAP molecule (transporter associated with processing). In the endoplasmic reticulum, the antigen binds to MHC class I molecules. Endoplasmic reticulum signal sequences bypass the cytosolic processing pathway and directly target endogenous antigens to the endoplasmic reticulum, where proteolytic degradation into peptide fragments occurs.
- 15 Such MHC class I targeting sequences are well known in the art, and include, e.g., signal sequences such as those from Ig kappa, tissue plasminogen activator or insulin. A preferred signal peptide is the human Ig kappa chain sequence. Endoplasmic reticulum signal sequences can also be used to target MHC class II epitopes to the endoplasmic reticulum, the site of MHC class I molecule assembly.

- 20 MHC class II targeting sequences are also used in the invention, e.g., those that target a peptide to the endocytic pathway. These targeting sequences typically direct extracellular antigens to enter the endocytic pathway, which results in the antigen being transferred to the lysosomal compartment where the antigen is proteolytically cleaved into antigen peptides for binding to MHC class II molecules. As with the normal
- 25 processing of exogenous antigen, a sequence that directs a MHC class II epitope to the endosomes of the endocytic pathway and/or subsequently to lysosomes, where the MHC class II epitope can bind to a MHC class II molecule, is a MHC class II targeting sequence. For example, group of MHC class II targeting sequences useful in the invention are lysosomal targeting sequences, which localize polypeptides to lysosomes.
- 30 Since MHC class II molecules typically bind to antigen peptides derived from proteolytic processing of endocytosed antigens in lysosomes, a lysosomal targeting sequence can function as a MHC class II targeting sequence. Lysosomal targeting sequences are well known in the art and include sequences found in the lysosomal proteins LAMP-1 and

LAMP-2 as described by August *et al.* (U.S. Patent No. 5,633,234, issued May 27, 1997), which is incorporated herein by reference.

Other lysosomal proteins that contain lysosomal targeting sequences include HLA-DM. HLA-DM is an endosomal/lysosomal protein that functions in  
5 facilitating binding of antigen peptides to MHC class II molecules. Since it is located in the lysosome, HLA-DM has a lysosomal targeting sequence that can function as a MHC class II molecule targeting sequence (Copier *et al.*, *J. Immunol.* 157:1017-1027 (1996), which is incorporated herein by reference).

The resident lysosomal protein HLA-DO can also function as a lysosomal  
10 targeting sequence. In contrast to the above described resident lysosomal proteins LAMP-1 and HLA-DM, which encode specific Tyr-containing motifs that target proteins to lysosomes, HLA-DO is targeted to lysosomes by association with HLA-DM (Liljedahl *et al.*, *EMBO J.* 15:4817-4824 (1996)), which is incorporated herein by reference. Therefore, the sequences of HLA-DO that cause association with HLA-DM and,  
15 consequently, translocation of HLA-DO to lysosomes can be used as MHC class II targeting sequences. Similarly, the murine homolog of HLA-DO, H2-DO, can be used to derive a MHC class II targeting sequence. A MHC class II epitope can be fused to HLA-DO or H2-DO and targeted to lysosomes.

In another example, the cytoplasmic domains of B cell receptor subunits  
20 Ig- $\alpha$  and Ig- $\beta$  mediate antigen internalization and increase the efficiency of antigen presentation (Bonnerot *et al.*, *Immunity* 3:335-347 (1995)), which is incorporated herein by reference. Therefore, the cytoplasmic domains of the Ig- $\alpha$  and Ig- $\beta$  proteins can function as MHC class II targeting sequences that target a MHC class II epitope to the endocytic pathway for processing and binding to MHC class II molecules.

Another example of a MHC class II targeting sequence that directs MHC  
25 class II epitopes to the endocytic pathway is a sequence that directs polypeptides to be secreted, where the polypeptide can enter the endosomal pathway. These MHC class II targeting sequences that direct polypeptides to be secreted mimic the normal pathway by which exogenous, extracellular antigens are processed into peptides that bind to MHC  
30 class II molecules. Any signal sequence that functions to direct a polypeptide through the endoplasmic reticulum and ultimately to be secreted can function as a MHC class II targeting sequence so long as the secreted polypeptide can enter the endosomal/lysosomal pathway and be cleaved into peptides that can bind to MHC class II molecules. An

example of such a fusion is shown in Figure 11, where the signal sequence of kappa immunoglobulin is fused to multiple MHC class II epitopes.

In another example, the Ii protein binds to MHC class II molecules in the endoplasmic reticulum, where it functions to prevent peptides present in the endoplasmic reticulum from binding to the MHC class II molecules. Therefore, fusion of a MHC class II epitope to the Ii protein targets the MHC class II epitope to the endoplasmic reticulum and a MHC class II molecule. For example, the CLIP sequence of the Ii protein can be removed and replaced with a MHC class II epitope sequence so that the MHC class II epitope is directed to the endoplasmic reticulum, where the epitope binds to a MHC class II molecule.

In some cases, antigens themselves can serve as MHC class II or I targeting sequences and can be fused to a universal MHC class II epitope to stimulate an immune response. Although cytoplasmic viral antigens are generally processed and presented as complexes with MHC class I molecules, long-lived cytoplasmic proteins such as the influenza matrix protein can enter the MHC class II molecule processing pathway (Guéguen & Long, *Proc. Natl. Acad. Sci. USA* 93:14692-14697 (1996)), which is incorporated herein by reference. Therefore, long-lived cytoplasmic proteins can function as a MHC class II targeting sequence. For example, an expression vector encoding influenza matrix protein fused to a universal MHC class II epitope can be advantageously used to target influenza antigen and the universal MHC class II epitope to the MHC class II pathway for stimulating an immune response to influenza.

Other examples of antigens functioning as MHC class II targeting sequences include polypeptides that spontaneously form particles. The polypeptides are secreted from the cell that produces them and spontaneously form particles, which are taken up into an antigen-presenting cell by endocytosis such as receptor-mediated endocytosis or are engulfed by phagocytosis. The particles are proteolytically cleaved into antigen peptides after entering the endosomal/lysosomal pathway.

One such polypeptide that spontaneously forms particles is HBV surface antigen (HBV-S) (Diminsky *et al.*, *Vaccine* 15:637-647 (1997); Le Borgne *et al.*, *Virology* 240:304-315 (1998)), each of which is incorporated herein by reference. Another polypeptide that spontaneously forms particles is HBV core antigen (Kühröber *et al.*, *International Immunol.* 9:1203-1212 (1997)), which is incorporated herein by reference. Still another polypeptide that spontaneously forms particles is the yeast Ty protein (Weber *et al.*, *Vaccine* 13:831-834 (1995)), which is incorporated herein by

reference. For example, an expression vector containing HBV-S antigen fused to a universal MHC class II epitope can be advantageously used to target HBV-S antigen and the universal MHC class II epitope to the MHC class II pathway for stimulating an immune response to HBV.

5

### Binding Affinity of Peptide Epitopes for HLA Molecules

The large degree of HLA polymorphism is an important factor to be taken into account with the epitope-based approach to vaccine development. To address this factor, epitope selection encompassing identification of peptides capable of binding at high or intermediate affinity to multiple HLA molecules is preferably utilized, most preferably these epitopes bind at high or intermediate affinity to two or more allele specific HLA molecules.

CTL-inducing peptides of interest for vaccine compositions preferably include those that have a binding affinity for class I HLA molecules of less than 500 nM. HTL-inducing peptides preferably include those that have a binding affinity for class II HLA molecules of less than 1000 nM. For example, peptide binding is assessed by testing the capacity of a candidate peptide to bind to a purified HLA molecule *in vitro*. Peptides exhibiting high or intermediate affinity are then considered for further analysis. Selected peptides are tested on other members of the supertype family. In preferred embodiments, peptides that exhibit cross-reactive binding are then used in vaccines or in cellular screening analyses.

Higher HLA binding affinity is typically correlated with greater immunogenicity. Greater immunogenicity can be manifested in several different ways. Immunogenicity corresponds to whether an immune response is elicited at all, and to the vigor of any particular response, as well as to the extent of a population in which a response is elicited. For example, a peptide might elicit an immune response in a diverse array of the population, yet in no instance produce a vigorous response. In accordance with these principles, close to 90% of high binding peptides have been found to be immunogenic, as contrasted with about 50% of the peptides which bind with intermediate affinity. Moreover, higher binding affinity peptides leads to more vigorous immunogenic responses. As a result, less peptide is required to elicit a similar biological effect if a high affinity binding peptide is used. Thus, in preferred embodiments of the invention, high binding epitopes are particularly useful.

- The relationship between binding affinity for HLA class I molecules and immunogenicity of discrete peptide epitopes on bound antigens has been determined for the first time in the art by the present inventors. The correlation between binding affinity and immunogenicity was analyzed in two different experimental approaches (Sette *et al.*, *J. Immunol.* 153:5586-5592 (1994)). In the first approach, the immunogenicity of potential epitopes ranging in HLA binding affinity over a 10,000-fold range was analyzed in HLA-A\*0201 transgenic mice. In the second approach, the antigenicity of approximately 100 different hepatitis B virus (HBV)-derived potential epitopes, all carrying A\*0201 binding motifs, was assessed by using PBL (peripheral blood lymphocytes) from acute hepatitis patients. Pursuant to these approaches, it was determined that an affinity threshold of approximately 500 nM (preferably 50 nM or less) determines the capacity of a peptide epitope to elicit a CTL response. These data are true for class I binding affinity measurements for naturally processed peptides and for synthesized T cell epitopes. These data also indicate the important role of determinant selection in the shaping of T cell responses (*see, e.g., Schaeffer et al. Proc. Natl. Acad. Sci. USA* 86:4649-4653, 1989).

- An affinity threshold associated with immunogenicity in the context of HLA class II DR molecules has also been delineated (*see, e.g., Southwood et al. J. Immunology* 160:3363-3373 (1998), and USSN 60/087192, filed 5/29/98). In order to define a biologically significant threshold of DR binding affinity, a database of the binding affinities of 32 DR-restricted epitopes for their restricting element (i.e., the HLA molecule that binds the motif) was compiled. In approximately half of the cases (15 of 32 epitopes), DR restriction was associated with high binding affinities, i.e. binding affinities of less than 100 nM. In the other half of the cases (16 of 32), DR restriction was associated with intermediate affinity (binding affinities in the 100-1000 nM range). In only one of 32 cases was DR restriction associated with an IC50 of 1000 nM or greater. Thus, 1000 nM can be defined as an affinity threshold associated with immunogenicity in the context of DR molecules.

### 30 Peptide Epitope Binding Motifs and Supermotifs

In the past few years evidence has accumulated to demonstrate that a large fraction of HLA class I and class II molecules can be classified into a relatively few supertypes, each characterized by largely overlapping peptide binding repertoires, and consensus structures of the main peptide binding pockets.



For HLA molecule pocket analyses, the residues comprising the B and F pockets of HLA class I molecules as described in crystallographic studies were analyzed (Guo *et al.*, *Nature* 360:364 (1992); Saper *et al.*, *J. Mol. Biol.* 219:277 (1991); Madden *et al.*, *Cell* 75:693 (1993); Parham *et al.*, *Immunol. Rev.* 143:141 (1995)). In these analyses, residues 9, 45, 63, 66, 67, 70, and 99 were considered to make up the B pocket; and the B pocket was deemed to determine the specificity for the amino acid residue in the second position of peptide ligands. Similarly, residues 77, 80, 81, and 116 were considered to determine the specificity of the F pocket; the F pocket was deemed to determine the specificity for the C-terminal residue of a peptide ligand bound by the HLA class I molecule.

Through the study of single amino acid substituted antigen analogs and the sequencing of endogenously bound, naturally processed peptides, critical residues required for allele-specific binding to HLA molecules have been identified. The presence of these residues correlates with binding affinity for HLA molecules. The identification of motifs and/or supermotifs that correlate with high and intermediate affinity binding is an important issue with respect to the identification of immunogenic peptide epitopes for the inclusion in a vaccine. Kast *et al.* (*J. Immunol.* 152:3904-3912 (1994)) have shown that motif-bearing peptides account for 90% of the epitopes that bind to allele-specific HLA class I molecules. In this study all possible peptides of 9 amino acids in length and overlapping by eight amino acids (240 peptides), which cover the entire sequence of the E6 and E7 proteins of human papillomavirus type 16, were evaluated for binding to five allele-specific HLA molecules that are expressed at high frequency among different ethnic groups. This unbiased set of peptides allowed an evaluation of the predictive value of HLA class I motifs. From the set of 240 peptides, 22 peptides were identified that bound to an allele-specific HLA molecules with high or intermediate affinity. Of these 22 peptides, 20, (i.e., 91%), were motif-bearing. Thus, this study demonstrates the value of motifs for the identification of peptide epitopes for inclusion in a vaccine: application of motif-based identification techniques eliminates screening of 90% of the potential epitopes in a target antigen protein sequence.

Peptides of the present invention may also include epitopes that bind to MHC class II DR molecules. There is a significant difference between class I and class II HLA molecules. This difference corresponds to the fact that, although a stringent size restriction and motif position relative to the binding pocket exists for peptides that bind to class I molecules, a greater degree of heterogeneity in both size and binding frame

position of the motif, relative to the N and C termini of the peptide, exists for class II peptide ligands.

This increased heterogeneity of HLA class II peptide ligands is due to the structure of the binding groove of the HLA class II molecule which, unlike its class I counterpart, is open at both ends. Crystallographic analysis of HLA class II DRB\*0101-peptide complexes showed that the residues occupying position 1 and position 6 of peptides complexed with DRB\*0101 engage two complementary pockets on the DRBa\*0101 molecules, with the P1 position corresponding to the most crucial anchor residue and the deepest hydrophobic pocket (see, e.g., Madden, *Ann. Rev. Immunol.* 13:587 (1995)). Other studies have also pointed to the P6 position as a crucial anchor residue for binding to various other DR molecules.

Thus, peptides of the present invention are identified by any one of several HLA class I or II -specific amino acid motifs (see, e.g., Tables I-III of USSN 09/226,775, and 09/239,043, herein incorporated by reference in their entirety). If the presence of the motif corresponds to the ability to bind several allele-specific HLA antigens it is referred to as a supermotif. The allele-specific HLA molecules that bind to peptides that possess a particular amino acid supermotif are collectively referred to as an HLA "supertype."

#### Immune Response-Stimulating Peptide Analogs

In general, CTL and HTL responses are not directed against all possible epitopes. Rather, they are restricted to a few "immunodominant" determinants (Zinkernagel *et al.*, *Adv. Immunol.* 27:5159 (1979); Bennink *et al.*, *J. Exp. Med.* 168:1935-1939 (1988); Rawle *et al.*, *J. Immunol.* 146:3977-3984 (1991)). It has been recognized that immunodominance (Benacerraf *et al.*, *Science* 175:273-279 (1972)) could be explained by either the ability of a given epitope to selectively bind a particular HLA protein (determinant selection theory) (Vitiello *et al.*, *J. Immunol.* 131:1635 (1983)); Rosenthal *et al.*, *Nature* 267:156-158 (1977)), or being selectively recognized by the existing TCR (T cell receptor) specificity (repertoire theory) (Klein, *Immunology, The Science of Self on self Discrimination*, pp. 270-310 (1982)). It has been demonstrated that additional factors, mostly linked to processing events, can also play a key role in dictating, beyond strict immunogenicity, which of the many potential determinants will be presented as immunodominant (Sercarz *et al.*, *Annu. Rev. Immunol.* 11:729-766 (1993)).

The concept of dominance and subdominance is relevant to immunotherapy of both infectious diseases and cancer. For example, in the course of chronic viral disease, recruitment of subdominant epitopes can be important for successful clearance of the infection, especially if dominant CTL or HTL specificities have been inactivated by functional tolerance, suppression, mutation of viruses and other mechanisms (Franco *et al.*, *Curr. Opin. Immunol.* 7:524-531 (1995)). In the case of cancer and tumor antigens, CTLs recognizing at least some of the highest binding affinity peptides might be functionally inactivated. Lower binding affinity peptides are preferentially recognized at these times, and may therefore be preferred in therapeutic or prophylactic anti-cancer vaccines.

In particular, it has been noted that a significant number of epitopes derived from known non-viral tumor associated antigens (TAA) bind HLA class I with intermediate affinity (IC50 in the 50-500 nM range). For example, it has been found that 8 of 15 known TAA peptides recognized by tumor infiltrating lymphocytes (TIL) or CTL bound in the 50-500 nM range. (These data are in contrast with estimates that 90% of known viral antigens were bound by HLA class I molecules with IC50 of 50 nM or less, while only approximately 10% bound in the 50-500 nM range (Sette *et al.*, *J. Immunol.*, 153:558-5592 (1994)). In the cancer setting this phenomenon is probably due to elimination, or functional inhibition of the CTL recognizing several of the highest binding peptides, presumably because of T cell tolerization events.

Without intending to be bound by theory, it is believed that because T cells to dominant epitopes may have been clonally deleted, selecting subdominant epitopes may allow extant T cells to be recruited, which will then lead to a therapeutic or prophylactic response. However, the binding of HLA molecules to subdominant epitopes is often less vigorous than to dominant ones. Accordingly, there is a need to be able to modulate the binding affinity of particular immunogenic epitopes for one or more HLA molecules, and thereby to modulate the immune response elicited by the peptide, for example to prepare analog peptides which elicit a more vigorous response. This ability would greatly enhance the usefulness of peptide-based vaccines and therapeutic agents.

Thus, although peptides with suitable cross-reactivity among all alleles of a superfamily are identified by the screening procedures described above, cross-reactivity is not always as complete as possible, and in certain cases procedures to further increase cross-reactivity of peptides can be useful; moreover, such procedures can also be used to modify other properties of the peptides such as binding affinity or peptide stability.

Having established the general rules that govern cross-reactivity of peptides for HLA alleles within a given motif or supermotif, modification (i.e., analoging) of the structure of peptides of particular interest in order to achieve broader (or otherwise modified) HLA binding capacity can be performed. More specifically, peptides which exhibit the  
5 broadest cross-reactivity patterns, can be produced in accordance with the teachings herein. The present concepts related to analog generation are set forth in greater detail in co-pending USSN 09/226,775.

In brief, the strategy employed utilizes the motifs or supermotifs which correlate with binding to certain HLA class I and II molecules. The motifs or supermotifs  
10 are defined by having primary anchors, and in many cases secondary anchors (see Tables I-III of USSN 09/226,775). Analog peptides can be created by substituting amino acids residues at primary anchor, secondary anchor, or at primary and secondary anchor positions. Generally, analogs are made for peptides that already bear a motif or supermotif. Preferred secondary anchor residues of supermotifs and motifs that have  
15 been defined for HLA class I and class II binding peptides are shown in Tables II and III, respectively, of USSN 09/226,775.

For a number of the motifs or supermotifs in accordance with the invention, residues are defined which are deleterious to binding to allele-specific HLA molecules or members of HLA supertypes that bind to the respective motif or supermotif  
20 (see Tables II and III of USSN 09/226,775). Accordingly, removal of such residues that are detrimental to binding can be performed in accordance with the methods described therein. For example, in the case of the A3 supertype, when all peptides that have such deleterious residues are removed from the population of analyzed peptides, the incidence of cross-reactivity increases from 22% to 37% (L. Sidney *et al.*, *Hum. Immunol.* 45:79  
25 (1996)). Thus, one strategy to improve the cross-reactivity of peptides within a given supermotif is simply to delete one or more of the deleterious residues present within a peptide and substitute a small "neutral" residue such as Ala (that may not influence T cell recognition of the peptide). An enhanced likelihood of cross-reactivity is expected if, together with elimination of detrimental residues within a peptide, "preferred" residues  
30 associated with high affinity binding to an allele-specific HLA molecule or to multiple HLA molecules within a superfamily are inserted.

To ensure that an analog peptide, when used as a vaccine, actually elicits a CTL response to the native epitope *in vivo* (or, in the case of class II epitopes, a failure to elicit helper T cells that cross-react with the wild type peptides), the analog peptide may

be used to immunize T cells *in vitro* from individuals of the appropriate HLA allele. Thereafter, the immunized cells' capacity to induce lysis of wild type peptide sensitized target cells is evaluated. In both class I and class II systems it will be desirable to use as targets, cells that have been either infected or transfected with the appropriate genes to establish whether endogenously produced antigen is also recognized by the relevant T cells.

Another embodiment of the invention is to create analogs of weak binding peptides, to thereby ensure adequate numbers of cross-reactive cellular binders. Class I peptides exhibiting binding affinities of 500-50000 nM, and carrying an acceptable but suboptimal primary anchor residue at one or both positions can be "fixed" by substituting preferred anchor residues in accordance with the respective supertype. The analog peptides can then be tested for crossbinding activity.

Another embodiment for generating effective peptide analogs involves the substitution of residues that have an adverse impact on peptide stability or solubility in, e.g., a liquid environment. This substitution may occur at any position of the peptide epitope. For example, a cysteine (C) can be substituted out in favor of gamma-amino butyric acid. Due to its chemical nature, cysteine has the propensity to form disulfide bridges and sufficiently alter the peptide structurally so as to reduce binding capacity. Substituting gamma-amino butyric acid for C not only alleviates this problem, but actually improves binding and crossbinding capability in certain instances (Sette *et al*, *In: Persistent Viral Infections* (Ahmed & Chen, eds., 1998)). Substitution of cysteine with gamma-amino butyric acid may occur at any residue of a peptide epitope, i.e., at either anchor or non-anchor positions.

## 25 Expression Vectors and Construction of a Minigene

The expression vectors of the invention contain at least one promoter element that is capable of expressing a transcription unit encoding the antigen of interest, for example, a MHC class I epitope or a MHC class II epitope and an MHC targeting sequence in the appropriate cells of an organism so that the antigen is expressed and targeted to the appropriate MHC molecule. For example, if the expression vector is administered to a mammal such as a human, a promoter element that functions in a human cell is incorporated into the expression vector. An example of an expression vector useful for expressing the MHC class II epitopes fused to MHC class II targeting

sequences and the MHC class I epitopes described herein is the pEP2 vector described in Example IV.

- This invention relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include
- 5 Sambrook *et al.*, *Molecular Cloning. A Laboratory Manual* (2nd ed. 1989); Krieger, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds., 1994); *Oligonucleotide Synthesis: A Practical Approach* (Gait, ed., 1984); Kuijpers, *Nucleic Acids Research* 18(17):5197 (1994); Ducholm, *J. Org. Chem.* 59:5767-5773 (1994); *Methods in Molecular Biology*, volume
  - 10 20 (Agrawal, ed.); and Tijssen, *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, e.g., Part I, chapter 2 "Overview of principles of hybridization and the strategy of nucleic acid probe assays" (1993).

- The minigenes are comprised of two or many different epitopes (*see, e.g.*, Tables 1-8). The nucleic acid encoding the epitopes are assembled in a minigene
- 15 according to standard techniques. In general, the nucleic acid sequences encoding minigene epitopes are isolated using amplification techniques with oligonucleotide primers, or are chemically synthesized. Recombinant cloning techniques can also be used when appropriate. Oligonucleotide sequences are selected which either amplify (when using PCR to assemble the minigene) or encode (when using synthetic oligonucleotides to
  - 20 assemble the minigene) the desired epitopes.

- Amplification techniques using primers are typically used to amplify and isolate sequences encoding the epitopes of choice from DNA or RNA (*see* U.S. Patents 4,683,195 and 4,683,202; *PCR Protocols: A Guide to Methods and Applications* (Innis *et al.*, eds, 1990)). Methods such as polymerase chain reaction (PCR) and ligase chain
- 25 reaction (LCR) can be used to amplify epitope nucleic acid sequences directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. Restriction endonuclease sites can be incorporated into the primers. Minigenes amplified by the PCR reaction can be purified from agarose gels and cloned into an appropriate vector.

- Synthetic oligonucleotides can also be used to construct minigenes. This
- 30 method is performed using a series of overlapping oligonucleotides, representing both the sense and non-sense strands of the gene. These DNA fragments are then annealed, ligated and cloned. Oligonucleotides that are not commercially available can be chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage & Caruthers, *Tetrahedron Letts.* 22:1859-1862 (1981), using an

automated synthesizer, as described in Van Devanter *et al.*, *Nucleic Acids Res.* 12:6159-6168 (1984). Purification of oligonucleotides is by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson & Reanier, *J. Chrom.* 255:137-149 (1983).

- 5           The epitopes of the minigene are typically subcloned into an expression vector that contains a strong promoter to direct transcription, as well as other regulatory sequences such as enhancers and polyadenylation sites. Suitable promoters are well known in the art and described, e.g., in Sambrook *et al.* and Ausubel *et al.* Eukaryotic expression systems for mammalian cells are well known in the art and are commercially available. Such promoter elements include, for example, cytomegalovirus (CMV), Rous sarcoma virus LTR and SV40.

- 10           The expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the minigene in host cells. A typical expression cassette thus contains a promoter operably linked to the minigene and signals required for efficient polyadenylation of the transcript. Additional elements of the cassette may include enhancers and introns with functional splice donor and acceptor sites.

- 15           In addition to a promoter sequence, the expression cassette can also contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes.

- 20           The particular expression vector used to transport the genetic information into the cell is not particularly critical. Any of the conventional vectors used for expression in eukaryotic cells may be used. Expression vectors containing regulatory elements from eukaryotic viruses are typically used in eukaryotic expression vectors, e.g., SV40 vectors, papilloma virus vectors, and vectors derived from Epstein Bar virus. Other exemplary eukaryotic vectors include pMSG, pAV009/A+, pMTO10/A+, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the SV40 early promoter, SV40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells. In one embodiment, the vector pEP2 is used in the present invention.

25           Other elements that are typically included in expression vectors also include a replicon that functions in *E. coli*, a gene encoding antibiotic resistance to permit

- selection of bacteria that harbor recombinant plasmids, and unique restriction sites in nonessential regions of the plasmid to allow insertion of eukaryotic sequences. The particular antibiotic resistance gene chosen is not critical, any of the many resistance genes known in the art are suitable. The prokaryotic sequences are preferably chosen
- 5 such that they do not interfere with the replication of the DNA in eukaryotic cells, if necessary.

#### Administration *In Vivo*

- The invention also provides methods for stimulating an immune response
- 10 by administering an expression vector of the invention to an individual. Administration of an expression vector of the invention for stimulating an immune response is advantageous because the expression vectors of the invention target MHC epitopes to MHC molecules, thus increasing the number of CTL and HTL activated by the antigens encoded by the expression vector.

- 15 Initially, the expression vectors of the invention are screened in mouse to determine the expression vectors having optimal activity in stimulating a desired immune response. Initial studies are therefore carried out, where possible, with mouse genes of the MHC targeting sequences. Methods of determining the activity of the expression vectors of the invention are well known in the art and include, for example, the uptake of
- 20  $^3\text{H}$ -thymidine to measure T cell activation and the release of  $^{51}\text{Cr}$  to measure CTL activity as described below in Examples II and III. Experiments similar to those described in Example IV are performed to determine the expression vectors having activity at stimulating an immune response. The expression vectors having activity are further tested in human. To circumvent potential adverse immunological responses to encoded
- 25 mouse sequences, the expression vectors having activity are modified so that the MHC class II targeting sequences are derived from human genes. For example, substitution of the analogous regions of the human homologs of genes containing various MHC class II targeting sequences are substituted into the expression vectors of the invention. Examples of such human homologs of genes containing MHC class II targeting sequences
- 30 are shown in Figures 12 to 17. Expression vectors containing human MHC class II targeting sequences, such as those described in Example I below, are tested for activity at stimulating an immune response in human.

The invention also relates to pharmaceutical compositions comprising a pharmaceutically acceptable carrier and an expression vector of the invention.



Pharmaceutically acceptable carriers are well known in the art and include aqueous or non-aqueous solutions, suspensions and emulsions, including physiologically buffered saline, alcohol/aqueous solutions or other solvents or vehicles such as glycols, glycerol, oils such as olive oil or injectable organic esters.

- 5 A pharmaceutically acceptable carrier can contain physiologically acceptable compounds that act, for example, to stabilize the expression vector or increase the absorption of the expression vector. Such physiologically acceptable compounds include, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants such as ascorbic acid or glutathione, chelating agents, low molecular weight polypeptides, antimicrobial agents, inert gases or other stabilizers or excipients. Expression vectors can  
10 additionally be complexed with other components such as peptides, polypeptides and carbohydrates. Expression vectors can also be complexed to particles or beads that can be administered to an individual, for example, using a vaccine gun. One skilled in the art would know that the choice of a pharmaceutically acceptable carrier, including a  
15 physiologically acceptable compound, depends, for example, on the route of administration of the expression vector.

- The invention further relates to methods of administering a pharmaceutical composition comprising an expression vector of the invention to stimulate an immune response. The expression vectors are administered by methods well known in the art as  
20 described in Donnelly *et al.* (*Ann. Rev. Immunol.* 15:617-648 (1997)); Felgner *et al.* (U.S. Patent No. 5,580,859, issued December 3, 1996); Felgner (U.S. Patent No. 5,703,055, issued December 30, 1997); and Carson *et al.* (U.S. Patent No. 5,679,647, issued October 21, 1997), each of which is incorporated herein by reference. In one embodiment, the minigene is administered as naked nucleic acid.

- 25 A pharmaceutical composition comprising an expression vector of the invention can be administered to stimulate an immune response in a subject by various routes including, for example, orally, intravaginally, rectally, or parenterally, such as intravenously, intramuscularly, subcutaneously, intraorbitally, intracapsularly, intraperitoneally, intracisternally or by passive or facilitated absorption through the skin  
30 using, for example, a skin patch or transdermal iontophoresis, respectively. Furthermore, the composition can be administered by injection, intubation or topically, the latter of which can be passive, for example, by direct application of an ointment or powder, or active, for example, using a nasal spray or inhalant. An expression vector also can be administered as a topical spray, in which case one component of the composition is an

- appropriate propellant. The pharmaceutical composition also can be incorporated, if desired, into liposomes, microspheres or other polymer matrices (Felgner *et al.*, U.S. Patent No. 5,703,055; Gregoriadis, *Liposome Technology*, Vols. I to III (2nd ed. 1993), each of which is incorporated herein by reference). Liposomes, for example, which
- 5 consist of phospholipids or other lipids, are nontoxic, physiologically acceptable and metabolizable carriers that are relatively simple to make and administer.

- The expression vectors of the invention can be delivered to the interstitial spaces of tissues of an animal body (Felgner *et al.*, U.S. Patent Nos. 5,580,859 and 5,703,055). Administration of expression vectors of the invention to muscle is a
- 10 particularly effective method of administration, including intradermal and subcutaneous injections and transdermal administration. Transdermal administration, such as by iontophoresis, is also an effective method to deliver expression vectors of the invention to muscle. Epidermal administration of expression vectors of the invention can also be employed. Epidermal administration involves mechanically or chemically irritating the
- 15 outermost layer of epidermis to stimulate an immune response to the irritant (Carson *et al.*, U.S. Patent No. 5,679,647).

- Other effective methods of administering an expression vector of the invention to stimulate an immune response include mucosal administration (Carson *et al.*, U.S. Patent No. 5,679,647). For mucosal administration, the most effective method of
- 20 administration includes intranasal administration of an appropriate aerosol containing the expression vector and a pharmaceutical composition. Suppositories and topical preparations are also effective for delivery of expression vectors to mucosal tissues of genital, vaginal and ocular sites. Additionally, expression vectors can be complexed to particles and administered by a vaccine gun.

- The dosage to be administered is dependent on the method of administration and will generally be between about 0.1  $\mu\text{g}$  up to about 200  $\mu\text{g}$ . For example, the dosage can be from about 0.05  $\mu\text{g}/\text{kg}$  to about 50  $\text{mg}/\text{kg}$ , in particular about 0.005-5  $\text{mg}/\text{kg}$ . An effective dose can be determined, for example, by measuring the immune response after administration of an expression vector. For example, the
- 30 production of antibodies specific for the MHC class II epitopes or MHC class I epitopes encoded by the expression vector can be measured by methods well known in the art, including ELISA or other immunological assays. In addition, the activation of T helper cells or a CTL response can be measured by methods well known in the art including, for

example, the uptake of  $^3\text{H}$ -thymidine to measure T cell activation and the release of  $^{51}\text{Cr}$  to measure CTL activity (*see* Examples II and III below).

The pharmaceutical compositions comprising an expression vector of the invention can be administered to mammals, particularly humans, for prophylactic or  
5 therapeutic purposes. Examples of diseases that can be treated or prevented using the expression vectors of the invention include infection with HBV, HCV, HIV and CMV as well as prostate cancer, renal carcinoma, cervical carcinoma, lymphoma, condyloma acuminatum and acquired immunodeficiency syndrome (AIDS).

In therapeutic applications, the expression vectors of the invention are  
10 administered to an individual already suffering from cancer, autoimmune disease or infected with a virus. Those in the incubation phase or acute phase of the disease can be treated with expression vectors of the invention, including those expressing all universal MHC class II epitopes, separately or in conjunction with other treatments, as appropriate.

In therapeutic and prophylactic applications, pharmaceutical compositions  
15 comprising expression vectors of the invention are administered to a patient in an amount sufficient to elicit an effective immune response to an antigen and to ameliorate the signs or symptoms of a disease. The amount of expression vector to administer that is sufficient to ameliorate the signs or symptoms of a disease is termed a therapeutically effective dose. The amount of expression vector sufficient to achieve a therapeutically  
20 effective dose will depend on the pharmaceutical composition comprising an expression vector of the invention, the manner of administration, the state and severity of the disease being treated, the weight and general state of health of the patient and the judgment of the prescribing physician.

25 All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily  
30 apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

## EXAMPLES

The following example is provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

5

EXAMPLE I: Construction of Expression Vectors Containing MHC Class II Epitopes

This example shows construction of expression vectors containing MHC class II epitopes that can be used to target antigens to MHC class II molecules.

- Expression vectors comprising DNA constructs were prepared using  
10 overlapping oligonucleotides, polymerase chain reaction (PCR) and standard molecular biology techniques (Dieffenbach & Dveksler, *PCR Primer: A Laboratory Manual* (1995); Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual* (2nd ed., 1989), each of which is incorporated herein by reference).

- To generate full length wild type Ii, the full length invariant chain was  
15 amplified, cloned, and sequenced and used in the construction of the three invariant chain constructs. Except where noted, the source of cDNA for all the constructs listed below was Mouse Spleen Marathon-Ready cDNA made from Balb/c males (Clontech; Palo Alto CA). The primer pairs were the oligonucleotide  
GCTAGCGCCGCCACCATGGATGACCAACGCGACCTC (SEQ ID NO:40), which is  
20 designated murli-F and contains an NheI site followed by the consensus Kozak sequence and the 5' end of the Ii cDNA; and the oligonucleotide  
GGTACCTCACAGGGTGACTTGACCCAG (SEQ ID NO:41), which is designated  
murli-R and contains a KpnI site and the 3' end of the Ii coding sequence.

- For the PCR reaction, 5  $\mu$ l of spleen cDNA and 250 nM of each primer  
25 were combined in a 100  $\mu$ l reaction with 0.25 mM each dNTP and 2.5 units of *Pfu* polymerase in *Pfu* polymerase buffer containing 10 mM KCl, 10 mM  $(\text{NH}_4)_2\text{SO}_4$ , 20 mM Tris-chloride, pH 8.75, 2 mM  $\text{MgSO}_4$ , 0.1% TRITON X-100 and 100  $\mu$ g/ml bovine serum albumin (BSA). A Perkin/Elmer 9600 PCR machine (Perkin Elmer; Foster City CA) was used and the cycling conditions were: 1 cycle of 95°C for 5 minutes, followed by 30  
30 cycles of 95°C for 15 seconds, 52°C for 30 seconds, and 72°C for 1 minute. The PCR reaction was run on a 1% agarose gel, and the 670 base pair product was cut out, purified by spinning through a Millipore Ultrafree-MC filter (Millipore; Bedford MA) and cloned into pCR-Blunt from Invitrogen (San Diego, CA). Individual clones were screened by

sequencing, and a correct clone (named bli#3) was used as a template for the helper constructs.

- DNA constructs containing pan DR epitope sequences and MHC II targeting sequences derived from the Ii protein were prepared. The Ii murine protein has been previously described (Zhu & Jones, *Nucleic Acids Res.* 17:447-448 (1989)), which is incorporated herein by reference. Briefly, the IiPADRE construct contains the full length Ii sequence with PADRE precisely replacing the CLIP region. The DNA construct encodes amino acids 1 through 87 of invariant chain, followed with the 13 amino acid PADRE sequence (SEQ ID NO:38) and the rest of the invariant chain DNA sequence (amino acids 101-215). The construct was amplified in 2 overlapping halves that were joined to produce the final construct. The two primers used to amplify the 5' half were murli-F and the oligonucleotide
- CAGGGTCCAGGCAGCCACGAAGCTTGGCCACAGGTTGGCAGA (SEQ ID NO:42), which is designated IiPADRE-R. The IiPADRE-R primer includes nucleotides 303-262 of IiPADRE. The 3' half was amplified with the primer
- GGCTGCCTGGACCTGAAGGCTGCCGCTATGTCCATGGATAAC (SEQ ID NO:43), which is designated IiPADRE-F and includes nucleotides 288-330 of IiPADRE; and murli-R. The PCR conditions were the same as described above, and the two halves were isolated by agarose gel electrophoresis as described above.
- Ten microliters of each PCR product was combined in a 100 µl PCR reaction with an annealing temperature of 50°C for five cycles to generate a full length template. Primers murli-F and murli-R were added and 25 more cycles carried out. The full length IiPADRE product was isolated, cloned, and sequenced as described above. This construct contains the murine Ii gene with a pan DR epitope sequence substituted for the CLIP sequence of Ii (Figure 1).

- A DNA construct, designated I80T, containing the cytoplasmic domain, the transmembrane domain and part of the luminal domain of Ii fused to a string of multiple MHC class II epitopes was constructed (Figure 2). Briefly, the string of multiple MHC class II epitopes was constructed with three overlapping oligonucleotides (oligos). Each oligo overlapped its neighbor by 15 nucleotides and the final MHC class II epitope string was assembled by extending the overlapping oligonucleotides in three sets of reactions using PCR. The three oligonucleotides were: oligo 1, nucleotides 241-310,
- CTTGCATGAAGCTTATCAGCCAGGCTGTGCACGCCGCTCACGCCGAAATCAA  
CGAAGCTGGAAGAACCC (SEQ ID NO:44);

oligo 2, nucleotides 364-295,

TTCTGGTCAGCAGAAAGAACAGGATAGGAGCGTTTGGAGGGCGATAAGCTGG  
AGGGGTTCTTCCAGCTTC (SEQ ID NO:45); and

oligo 3, nucleotides 350-42,

5 TTCTGCTGACCAGAATCCTGACAATCCCCAGTCCCTGGACGCCAAGTTCGTG  
GCTGCCTGGACCCCTGAAG (SEQ ID NO:46).

For the first PCR reaction, 5 µg of oligos 1 and 2 were combined in a 100  
µl reaction containing *Pfu* polymerase. A Perkin/Elmer 9600 PCR machine was used and  
the annealing temperature used was 45°C. The PCR product was gel-purified, and a

10 second reaction containing the PCR product of oligos 1 and 2 with oligo 3 was annealed  
and extended for 10 cycles before gel purification of the full length product to be used as  
a "mega-primer."

The I80T construct was made by amplifying bli#3 with murli-F and the  
mega-primer. The cycling conditions were: 1 cycle of 95°C for 5 minutes, followed by 5  
cycles of 95°C for 15 seconds, 37°C for 30 seconds, and 72°C for 1 minute. Primer Help-  
15 epR was added and an additional 25 cycles were carried out with the annealing

temperature raised to 47°C. The Help-epR primer

GGTACCTCAAGCGGCAGCCTTCAGGGTCCAGGCA (SEQ ID NO:47) corresponds  
to nucleotides 438-405. The full length I80T product was isolated, cloned, and sequenced  
20 as above.

The I80T construct (Figure 2) encodes amino acid residues 1 through 80 of  
Ii, containing the cytoplasmic domain, the transmembrane domain and part of the luminal  
domain, fused to a string of multiple MHC class II epitopes corresponding to: amino acid  
residues 323-339 of ovalbumin

25 (IleSerGlnAlaValHisAlaAlaHisAlaGluIleAsnGluAlaGlyArg; SEQ ID NO:48); amino  
acid residues 128 to 141 of HBV core antigen (amino acids  
ThrProProAlaTyrArgProProAsnAlaProIleLeu; SEQ ID NO:49); amino acid residues 182  
to 196 of HBV env (amino acids PhePheLeuLeuThrArgIleLeuThrIleProGlnSerLeuAsp;  
SEQ ID NO:50); and the pan DR sequence designated SEQ ID NO:38.

30 A DNA construct containing the cytoplasmic domain, transmembrane  
domain and a portion of the luminal domain of Ii fused to the MHC class II epitope string  
shown in Figure 2 and amino acid residues 101 to 215 of Ii encoding the trimerization  
region of Ii was generated (Figure 3). This construct, designated IiThfull, encodes the  
first 80 amino acids of invariant chain followed by the MHC class II epitope string

(replacing CLIP) and the rest of the invariant chain (amino acids 101-215). Briefly, the construct was generated as two overlapping halves that were annealed and extended by PCR to yield the final product.

The 5' end of liThfull was made by amplifying I80T with murli-F (SEQ ID NO:40) and Th-Pad-R. The Th-Pad-R primer AGCGGCAGCCTTCAGGGTC (SEQ ID NO:51) corresponds to nucleotides 429-411. The 3' half was made by amplifying bli#3 with liPADRE-F and murli-R (SEQ ID NO:41). The liPADRE-F primer GGCTGCCTGGACCCTGAAGGCTGCCGCTATGCCATGGATAAC (SEQ ID NO:52) corresponds to nucleotides 402-444. Each PCR product was gel purified and mixed, then denatured, annealed, and extended by five cycles of PCR. Primers murli-F (SEQ ID NO:40) and murli-R (SEQ ID NO:41) were added and another 25 cycles performed. The full length product was gel purified, cloned, and sequenced.

All of the remaining constructs described below were made essentially according to the scheme shown in Figure 18. Briefly, primer pairs 1F plus 1R, designated below for each specific construct, were used to amplify the specific signal sequence and contained an overlapping 15 base pair tail identical to the 5' end of the MHC class II epitope string. Primer pair Th-ova-F, ATCAGCCAGGCTGTGCACGC (SEQ ID NO:53), plus Th-Pad-R (SEQ ID NO:51) were used to amplify the MHC class II epitope string. A 15 base pair overlap and the specific transmembrane and cytoplasmic tail containing the targeting signals were amplified with primer pairs 2F plus 2R.

All three pieces of each cDNA were amplified using the following conditions: 1 cycle of 95°C for 5 minutes, followed by 30 cycles of 95°C for 15 seconds, 52°C for 30 seconds, and 72°C for 1 minute. Each of the three fragments was agarose-gel purified, and the signal sequence and MHC class II string fragments were combined and joined by five cycles in a second PCR. After five cycles, primers 1F and Th-Pad-R were added for 25 additional cycles and the PCR product was gel purified. This signal sequence plus MHC class II epitope string fragment was combined with the transmembrane plus cytoplasmic tail fragment for the final PCR. After five cycles, primers 1F plus 2R were added for 25 additional cycles and the product was gel purified, cloned and sequenced.

A DNA construct containing the murine immunoglobulin kappa signal sequence fused to the T helper epitope string shown in Figure 2 and the transmembrane and cytoplasmic domains of LAMP-1 was generated (Figure 4) (Granger *et al.*, *J. Biol. Chem.* 265:12036-12043 (1990)), which is incorporated by reference (mouse LAMP-1

GenBank accession No. M32015). This construct, designated kappaLAMP-Th, contains the consensus mouse immunoglobulin kappa signal sequence and was amplified from a plasmid containing full length immunoglobulin kappa as depicted in Figure 18. The primer 1F used was the oligonucleotide designated KappaSig-F,

5 GCTAGCGCCGCCACCATGGGAATGCAG (SEQ ID NO:54).

The primer 1R used was the oligonucleotide designated Kappa-Th-R, CACAGCCTGGCTGATTCTCTGGACCC (SEQ ID NO:55).

The primer 2F used was the oligonucleotide designated PAD/LAMP-F, CTGAAGGCTGCCGCTAACACATGTTGATCCCC (SEQ ID NO:56). The primer 2R used was the oligonucleotide designated LAMP-CYTOR,

10 GGTACCCTAGATGGTCTGATAGCC (SEQ ID NO:57).

A DNA construct containing the signal sequence of H2-M fused to the MHC class II epitope string shown in Figure 2 and the transmembrane and cytoplasmic domains of H2-M was generated (Figure 5). The mouse H2-M gene has been described previously, Peleraux *et al.*, *Immunogenetics* 43:204-214 (1996)), which is incorporated herein by reference. This construct was designated H2M-Th and was constructed as depicted in Figure 18. The primer 1F used was the oligonucleotide designated H2-Mb-1F, GCC GCT AGC GCC ACC ATG GCT GCA CTC TGG (SEQ ID NO:58). The primer 1R used was the oligonucleotide designated H2-Mb-1R, CAC AGC CTG GCT

20 GAT CCC CAT ACA GTG CAG (SEQ ID NO:59). The primer 2F used was the oligonucleotide designated H2-Mb-2F, CTG AAG GCT GCC GCT AAG GTC TCT GTG TCT (SEQ ID NO:60). The primer 2R used was the oligonucleotide designated H2-Mb-2R, GCG GGT ACC CTA ATG CCG TCC TTC (SEQ ID NO:61).

A DNA construct containing the signal sequence of H2-DO fused to the MHC class II epitope string shown in Figure 2 and the transmembrane and cytoplasmic domains of H2-DO was generated (Figure 6). The mouse H2-DO gene has been described previously (Larhammar *et al.*, *J. Biol. Chem.* 260:14111-14119 (1985)), which is incorporated herein by reference (GenBank accession No. M19423). This construct, designated H2O-Th, was constructed as depicted in Figure 18. The primer 1F used was the oligonucleotide designated H2-Ob-1F, GCG GCT AGC GCC ACC ATG GGC

30 GCT GGG AGG (SEQ ID NO:62). The primer 1R used was the oligonucleotide designated H2-Ob-1R, TGC ACA GCC TGG CTG ATG GAA TCC AGC CTC (SEQ ID NO:63). The primer 2F used was the oligonucleotide designated H2-Ob-2F, CTG AAG GCT GCC GCT ATA CTG AGT GGA GCT (SEQ ID NO:64). The primer 2R used was



the oligonucleotide designated H2-Ob-2R, GCC GGT ACC TCA TGT GAC ATG TCC CG (SEQ ID NO:65).

A DNA construct containing a pan DR epitope sequence (SEQ ID NO:38) fused to the amino-terminus of influenza matrix protein is generated (Figure 7). This construct, designated PADRE-Influenza matrix, contains the universal MHC class II epitope PADRE attached to the amino terminus of the influenza matrix coding sequence. The construct is made using a long primer on the 5' end primer. The 5' primer is the oligonucleotide  
 GCTAGCGCCGCCACCATGGCCAAGTTCGTGGCTGCCTGGACCCTGAAGGCTGC  
 CGCTATGAGTCTTCTAACCGAGGTCTGA (SEQ ID NO:66). The 3' primer is the oligonucleotide TCACTGAATCGCTGCATCTGCACCCCAT (SEQ ID NO:67). Influenza virus from the America Type Tissue Collection (ATCC) is used as a source for the matrix coding region (Perdue *et al. Science* 279:393-396 (1998)), which is incorporated herein by reference (GenBank accession No. AF036358).

A DNA construct containing a pan DR epitope sequence (SEQ ID NO:38) fused to the amino-terminus of HBV-S antigen was generated (Figure 8). This construct is designated PADRE-HBV-s and was generated by annealing two overlapping oligonucleotides to add PADRE onto the amino terminus of hepatitis B surface antigen (Michel *et al., Proc. Natl. Acad. Sci. USA* 81:7708-7712 (1984); Michel *et al., Proc. Natl. Acad. Sci. USA* 92:5307-5311 (1995)), each of which is incorporated herein by reference. One oligonucleotide was  
 GCTAGCGCCGCCACCATGGCCAAGTTCGTGGCTGCCTGGACCCTGAAGGCTGC  
 CGCTC (SEQ ID NO:68). The second oligonucleotide was  
 CTCGAGAGCGGCAGCCTTCAGGGTCCAGGCAGCCACGAAGTTCGGCCATGGTGC  
 GCGGCG (SEQ ID NO:69). When annealed, the oligos have NheI and XhoI cohesive ends. The oligos were heated to 100°C and slowly cooled to room temperature to anneal. A three part ligation joined PADRE with an XhoI-KpnI fragment containing HBV-s antigen into the NheI plus KpnI sites of the expression vector.

A DNA construct containing the signal sequence of Ig- $\alpha$  fused to the MHC class II epitope string shown in Figure 2 and the transmembrane and cytoplasmic domains of Ig- $\alpha$  was generated (Figure 9). The mouse Ig- $\alpha$  gene has been described previously (Kashiwamura *et al., J. Immunol.* 145:337-343 (1990)), which is incorporated herein by reference (GenBank accession No. M31773). This construct, designated Ig-alphaTh, was constructed as depicted in Figure 18. The primer 1F used was the oligonucleotide

designated Ig alpha-1F, GCG GCT AGC GCC GCC ACC ATG CCA GGG GGT CTA (SEQ ID NO:70). The primer 1R used was the oligonucleotide designated Igalpha-1R, GCA CAG CCT GGC TGA TGG CCT GGC ATC CGG (SEQ ID NO:71). The primer 2F used was the oligonucleotide designated Igalpha-2F, CTG AAG GCT GCC GCT GGG  
 5 ATC ATC TTG CTG (SEQ ID NO:72). The primer 2R used was the oligonucleotide designated Igalpha-2R, GCG GGT ACC TCA TGG CTT TTC CAG CTG (SEQ ID NO:73).

A DNA construct containing the signal sequence of Ig- $\beta$  fused to the MHC class II string shown in Figure 2 and the transmembrane and cytoplasmic domains of Ig $\beta$  was generated (Figure 10). The Ig- $\beta$  sequence is the B29 gene of mouse and has been  
 10 described previously (Hermanson *et al.*, *Proc. Natl. Acad. Sci. USA* 85:6890-6894 (1988)), which is incorporated herein by reference (GenBank accession No. J03857). This construct, designated Ig-betaTh, was constructed as depicted in Figure 18. The primer 1F used was the oligonucleotide designated B29-1F (33mer) GCG GCT AGC  
 15 GCC GCC ACC ATG GCC ACA CTG GTG (SEQ ID NO:74). The primer 1R used was the oligonucleotide designated B29-1R (30mer) CAC AGC CTG GCT GAT CGG CTC ACC TGA GAA (SEQ ID NO:75). The primer 2F used was the oligonucleotide designated B292F (30mer) CTG AAG GCT GCC GCT ATT ATC TTG ATC CAG (SEQ ID NO: 76). The primer 2R used was the oligonucleotide designated B29-2R (27mer),  
 20 GCC GGT ACC TCA TTC CTG GCC TGG ATG (SEQ ID NO:77).

A DNA construct containing the signal sequence of the kappa immunoglobulin signal sequence fused to the MHC class II epitope string shown in Figure 2 was constructed (Figure 11). This construct is designated SigTh and was generated by using the kappaLAMP-Th construct (shown in Figure 4) and amplifying  
 25 with the primer pair KappaSig-F (SEQ ID NO:54) plus Help-epR (SEQ ID NO:47) to create SigTh. SigTh contains the kappa immunoglobulin signal sequence fused to the T helper epitope string and terminated with a translational stop codon.

Constructs encoding human sequences corresponding to the above described constructs having mouse sequences are prepared by substituting human  
 30 sequences for the mouse sequences. Briefly, for the II PADRE construct, corresponding to Figure 1, amino acid residues 1-80 from the human Ii gene HLA-DR sequence (Figure 12) (GenBank accession No. X00497 M14765) is substituted for the mouse Ii sequences, which is fused to PADRE, followed by human invariant chain HLA-DR amino acid residues 114-223. For the I80T construct, corresponding to Figure 2, amino acid residues

1-80 from the human sequence of Ii is followed by a MHC class II epitope string. For the IiThfull construct, corresponding to Figure 3, amino acid residues 1-80 from the human sequence of Ii, which is fused to a MHC class II epitope string, is followed by human invariant chain amino acid residues 114-223.

- 5 For the LAMP-Th construct, similar to Figure 4, the signal sequence encoded by amino acid residues 1-19 (nucleotides 11-67) of human LAMP-1 (Figure 13) (GenBank accession No. J04182), which is fused to the MHC class II epitope string, is followed by the transmembrane (nucleotides 1163-1213) and cytoplasmic tail (nucleotides 1214-1258) region encoded by amino acid residues 380-416 of human
- 10 LAMP-1.

- For the HLA-DM-Th construct, corresponding to Figure 5, the signal sequence encoded by amino acid residues 1-17 (nucleotides 1-51) of human HLA-DMB (Figure 14) (GenBank accession No. U15085), which is fused to the MHC class II epitope string, is followed by the transmembrane (nucleotides 646-720) and cytoplasmic tail
- 15 (nucleotides 721-792) region encoded by amino acid residues 216-263 of human HLA-DMB.

- For the HLA-DO-Th construct, corresponding to Figure 6, the signal sequence encoded by amino acid residues 1-21 (nucleotides 1-63) of human HLA-DO (Figure 15) (GenBank accession No. L29472 J02736 N00052), which is fused to the
- 20 MHC class II epitope string, is followed by the transmembrane (nucleotides 685-735) and cytoplasmic tail (nucleotides 736-819) region encoded by amino acid residues 223-273 of human HLA-DO.

- For the Ig-alphaTh construct, corresponding to Figure 9, the signal sequence encoded by amino acid residues 1-29 (nucleotides 1-87) of human Ig-alpha MB-1 (Figure 16) (GenBank accession No. U05259), which is fused to the MHC class II epitope
- 25 string, is followed by the transmembrane (nucleotides 424-498) and cytoplasmic tail (nucleotides 499-678) region encoded by amino acid residues 142-226 of human Ig-alpha MB-1.

- For the Ig-betaTh construct, corresponding to Figure 10, the signal
- 30 sequence encoded by amino acid residues 1-28 (nucleotides 17-100) of human Ig-beta B29 (Figure 17) (GenBank accession No. M80461), which is fused to the MHC class II epitope string, is followed by the transmembrane (nucleotides 500-547) and cytoplasmic tail (nucleotides 548-703) region encoded by amino acid residues 156-229 of human Ig-beta.

The SigTh construct shown in Figure 11 can be used in mouse and human. Alternatively, a signal sequence derived from an appropriate human gene containing a signal sequence can be substituted for the mouse kappa immunoglobulin sequence in the Sig Th construct.

- 5           The PADRE-Influenza matrix construct shown in Figure 7 and the PADRE-HBV's construct shown in Figure 8 can be used in mouse and human.

Some of the DNA constructs described above were cloned into the vector pEP2 (Figure 19; SEQ ID NO:35). The pEP2 vector was constructed to contain dual CMV promoters. The pEP2 vector used the backbone of pcDNA3.1(-)Myc-His A from  
10   Invitrogen and pIRES1hyg from Clontech. Changes were made to both vectors before the CMV transcription unit from pIRES1hyg was moved into the modified pcDNA vector.

- The pcDNA3.1(-)Myc-His A vector (<http://www.invitrogen.com>) was modified. Briefly, the PvuII fragment (nucleotides 1342-3508) was deleted. A BspHI fragment that contains the Ampicillin resistance gene (nucleotides 4404-5412) was cut  
15   out. The Ampicillin resistance gene was replaced with the kanamycin resistance gene from pUC4K (GenBank Accession #X06404). pUC4K was amplified with the primer set: TCTGATGTTACATTGCACAAG (SEQ ID NO:78) (nucleotides 1621-1601) and GCGCACTCATGATGCTCTGCCAGTGTACAACC (SEQ ID NO:79) (nucleotides 682-702 plus the addition of a BspHI restriction site on the 5' end). The PCR product  
20   was digested with BspHI and ligated into the vector digested with BspHI. The region between the PmeI site at nucleotide 905 and the EcoRV site at nucleotide 947 was deleted. The vector was then digested with PmeI (cuts at nucleotide 1076) and ApaI (cuts at nucleotide 1004), Klenow filled in at the cohesive ends and ligated. The KpnI site at nucleotide 994 was deleted by digesting with KpnI and filling in the ends with Klenow  
25   DNA polymerase, and ligating. The intron A sequence from CMV (GenBank accession M21295, nucleotides 635-1461) was added by amplifying CMV DNA with the primer set: GCGTCTAGAGTAAGTACCGCCTATAGACTC (SEQ ID NO:80) (nucleotides 635-655 plus an XbaI site on the 5' end) and CCGCTAGCCTGCAGAAAAGACCCATGGAA (SEQ ID NO:81) (nucleotides 1461-1441 plus an NheI site on the 3' end). The PCR  
30   product was digested with XbaI and NheI and ligated into the NheI site of the vector (nucleotide 895 of the original pcDNA vector) so that the NheI site was on the 3' end of the intron.

          To modify the pIRES1hyg vector (GenBank Accession U89672, Clontech), the KpnI site (nucleotide 911) was deleted by cutting and filling in with

Klenow. The plasmid was cut with NotI (nucleotide 1254) and XbaI (nucleotide 3196) and a polylinker oligo was inserted into the site. The polylinker was formed by annealing the following two oligos:

GGCCGCAAGGAAAAAATCTAGAGTCGGCCATAGACTAATGCCGTACCG (SEQ

5 ID NO:82) and

CTAGCGGTACCGGCATTAGTCTATGGCCGACTCTAGATTTTTCCTTGC (SEQ  
ID NO:83). The resulting plasmid was cut with HincII and the fragment between HincII  
sites 234 and 3538 was isolated and ligated into the modified pcDNA vector. This  
fragment contains a CMV promoter, intron, polylinker, and polyadenylation signal.

10 The pRES1hyg piece and the pcDNA piece were combined to form pEP2.  
The modified pcDNA3.1(-)Myc-His A vector was partially digested with PvuII to isolate  
a linear fragment with the cut downstream of the pcDNA polyadenylation signal (the  
other PvuII site is the CMV intron). The HincII fragment from the modified pRES1hyg  
vector was ligated into the PvuII cut vector. The polyadenylation signal from the pcDNA  
15 derived transcription unit was deleted by digesting with EcoRI (pcDNA nucleotide 955)  
and XhoI (pRES1hyg nucleotide 3472) and replaced with a synthetic polyadenylation  
sequence. The synthetic polyadenylation signal was described in Levitt *et al.*, *Genes and  
Development* 3:1019-1025 (1989)).

Two oligos were annealed to produce a fragment that contained a  
20 polylinker and polyadenylation signal with EcoRI and XhoI cohesive ends. The oligos  
were:  
AATTCGGATATCCAAGCTTGATGAATAAAAGATCAGAGCTCTAGTGATCTGTGT  
GTTGGTTTTTTGTGTGC (SEQ ID NO:84) and  
TCGAGCACACAAAAACCAACACACAGATCACTAGAGCTCTGATCTTTTATT  
25 CATCAAGCTTGGATATCCG (SEQ ID NO:85).

The resulting vector is named pEP2 and contains two separate  
transcription units. Both transcription units use the same CMV promoter but each  
contains different intron, polylinker, and polyadenylation sequences.

The pEP2 vector contains two transcription units. The first transcription  
30 unit contains the CMV promoter initially from pcDNA (nucleotides 210-862 in Figure  
19), CMV intron A sequence (nucleotides 900-1728 in Figure 19), polylinker cloning site  
(nucleotides 1740-1760 in Figure 19) and synthetic polyadenylation signal (nucleotides  
1764-1769 in Figure 19). The second transcription unit, which was initially derived from  
pRES1hyg, contains the CMV promoter (nucleotides 3165-2493 in Figure 19), intron

sequence (nucleotides 2464-2173 in Figure 19), polylinker clone site (nucleotides 2126-2095 in Figure 19) and bovine growth hormone polyadenylation signal (nucleotides 1979-1974 in Figure 19). The kanamycin resistance gene is encoded in nucleotides 4965-4061 (Figure 19).

- 5 The DNA constructs described above were digested with NheI and KpnI and cloned into the XbaI and KpnI sites of pEP2 (the second transcription unit).

Additional vectors were also constructed. To test for the effect of co-expression of MHC class I epitopes with MHC class II epitopes, an insert was generated, designated AOS, that contains nine MHC class I epitopes. The AOS insert was initially  
 10 constructed in the vector pMIN.0 (Figure 20; SEQ ID NO:36). Briefly, the AOS insert contains nine MHC class I epitopes, six restricted by HLA-A2 and three restricted by HLA-A11, and the universal MHC class II epitope PADRE. The vector pMIN.0 contains epitopes from HBV, HIV and a mouse ovalbumin epitope. The MHC class I epitopes appear in pMIN.0 in the following order:

- 15 consensus mouse Ig Kappa signal sequence (pMIN.0 amino acid residues 1-20, nucleotides 16-81) MQVQIQSLFLLLLWVPGSRG (SEQ ID NO:86) encoded by nucleotides ATG CAG GTG CAG ATC CAG AGC CTG TTT CTG CTC CTC CTG TGG GTG CCC GGG TCC AGA GGA (SEQ ID NO:87);

HBV pol 149-159 (A11 restricted)

- 20 (pMIN.0 amino acid residues 21-31, nucleotides 82-114)  
 HTLWKAGILYK (SEQ ID NO:88) encoded by nucleotides CAC ACC CTG TGG AAG GCC GGA ATC CTG TAT AAG (SEQ ID NO:89);

- PADRE-universal MHC class II epitope (pMIN.0 amino acid residues 32-45, nucleotides 115-153) AKFVAAWTLKAAA (SEQ ID NO:38) encoded by nucleotides  
 25 GCC AAG TTC GTG GCT GCC TGG ACC CTG AAG GCT GCC GCT (SEQ ID NO:90);

HBV core 18-27 (A2 restricted) (pMIN.0 amino acid residues 46-55, nucleotides 154-183) FLPSDFFPVS (SEQ ID NO:91) encoded by nucleotides TTC CTG CCT AGC GAT TTC TTT CCT AGC GTG (SEQ ID NO:92);

- 30 HIV env 120-128 (A2 restricted) (pMIN.0 amino acid residues 56-64, nucleotides 184-210) KLTPLCVTL (SEQ ID NO:93) encoded by nucleotides AAG CTG ACC CCA CTG TGC GTG ACC CTG (SEQ ID NO:94);

HBV pol 551-559 (A2 restricted) (pMIN.0 amino acid residues 65-73, nucleotides 211-237) YMDDVVLGA (SEQ ID NO:95) encoded by nucleotides TAT ATG GAT GAC GTG GTG CTG GGA GCC (SEQ ID NO:96);

mouse ovalbumin 257-264 (K<sup>b</sup> restricted) (pMIN.0 amino acid residues 74-81, nucleotides 238-261) SIINFELK (SEQ ID NO:97) encoded by nucleotides AGC ATC ATC AAC TTC GAG AAG CTG (SEQ ID NO:98);

HBV pol 455-463 (A2 restricted) (pMIN.0 amino acid residues 82-90, nucleotides 262-288) GLSRYVARL (SEQ ID NO:99) encoded by nucleotides GGA CTG TCC AGA TAC GTG GCT AGG CTG (SEQ ID NO:100);

10 HIV pol 476-84 (A2 restricted) (pMIN.0 amino acid residues 91-99, nucleotides 289-315) LKPEPVHGV (SEQ ID NO:101) encoded by nucleotides ATC CTG AAG GAG CCT GTG CAC GGC GTG (SEQ ID NO:102);

HBV core 141-151 (A11 restricted)

(pMIN.0 amino acid residues 100-110, nucleotides 316-348)

15 STLPETTVVRR (SEQ ID NO:103) encoded by nucleotides TCC ACC CTG CCA GAG ACC ACC GTG GTG AGG AGA (SEQ ID NO:104);

HIV env 49-58 (A11 restricted) (pMIN.0 amino acid residues 111-120, nucleotides 349-378) TVYYGVPVWK (SEQ ID NO:105) encoded by nucleotides ACC GTG TAC TAT GGA GTG CCT GTG TGG AAG (SEQ ID NO:106); and

20 HBV env 335-343 (A2 restricted) (pMIN.0 amino acid residues 121-129, nucleotides 378-405) WLSLLVPFV (SEQ ID NO:107) encoded by nucleotides TGG CTG AGC CTG CTG GTG CCC TTT GTG (SEQ ID NO:108).

The pMIN.0 vector contains a KpnI restriction site (pMIN.0 nucleotides 406-411) and a NheI restriction site (pMIN.0 nucleotides 1-6). The pMIN.0 vector  
25 contains a consensus Kozak sequence (nucleotides 7-18) (GCCGCCACCATG; SEQ ID NO:109) and murine Kappa Ig-light chain signal sequence followed by a string of 10 MHC class I epitopes and one universal MHC class II epitope. The pMIN.0 sequence encodes an open reading frame fused to the Myc and His antibody epitope tag coded for by the pcDNA 3.1 Myc-His vector. The pMIN.0 vector was constructed with eight  
30 oligonucleotides:

MinI oligo

GAGGAGCAGAAACAGGCTCTGGATCTGCACCTGCAITCCCATGGTGGCGGCGC  
TAGCAAGCTTCTTGCGC (SEQ ID NO:110);

## Min2 oligo

CCTGTTTCTGCTCCTCTGTGGGTGCCCGGTCCAGAGGACACACCTGTGGA  
AGGCCGGAATCCTGTATA (SEQ ID NO:111);

## Min3 oligo

- 5 TCGTAGGCAGGAAAGCGGCAGCCTTCAGGGTCCAGGCAGCCACGAACTGG  
CCTTATACAGGATTCCGG (SEQ ID NO:112);

## Min4 oligo

CTTTCCTGCCTAGCGATTCTTTCTAGCGTGAAGCTGACCCCACTGTGCGTGA  
CCCTGTATATGGATGAC (SEQ ID NO:113);

- 10 Min5 oligo

CGTACCTGGACAGTCCAGCTTCTCGAAGTTGATGATGCTGGCT  
CCCAGCACCACGTCATCCATATACAG (SEQ ID NO:114);

## Min6 oligo

- GGACTGTCCAGATACGTGGCTAGGCTGATCCTGAAGGAGCCTGTGCACGGCGT  
15 GTCCACCCTGCCAGAGAC (SEQ ID NO:115);

## Min7 oligo

GCTCAGCCACTCCACACAGGCACTCCATAGTACACGGTCTCCTCACCACGG  
TGGTCTCTGGCAGGGTG (SEQ ID NO:116);

## Min8 oligo

- 20 GTGGAAGTGGCTGAGCCTGCTGGTGCCCTTTGTGGGTACCTGATCTAGAGC  
(SEQ ID NO:117).

Additional primers were flanking primer 5', GCG CAA GAA GCT TGC  
TAG CG (SEQ ID NO:118) and flanking primer 3', GCT CTA GAT CAG GTA CCC  
CAC (SEQ ID NO:119).

- 25 The original pMIN.0 minigene construction was carried out using eight  
overlapping oligos averaging approximately 70 nucleotides in length, which were  
synthesized and HPLC purified by Operon Technologies Inc. Each oligo overlapped its  
neighbor by 15 nucleotides, and the final multi-epitope minigene was assembled by  
extending the overlapping oligos in three sets of reactions using PCR (Ho *et al.*, *Gene*  
30 77:51-59 (1989).

For the first PCR reaction, 5 µg of each of two oligos were annealed and  
extended: 1+2, 3+4, 5+6, and 7+8 were combined in 100 µl reactions containing 0.25 mM  
each dNTP and 2.5 units of Pfu polymerase in Pfu polymerase buffer containing 10 mM  
KCl, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 20 mM Tris-chloride, pH 8.75, 2 mM MgSO<sub>4</sub>, 0.1% TRITON



X-100 and 100 mg/ml BSA. A Perkin/Elmer 9600 PCR machine was used and the annealing temperature used was 5°C below the lowest calculated  $T_m$  of each primer pair. The full length dimer products were gel-purified, and two reactions containing the product of 1-2 and 3-4, and the product of 5-6 and 7-8 were mixed, annealed and extended for 10 cycles. Half of the two reactions were then mixed, and 5 cycles of annealing and extension carried out before flanking primers were added to amplify the full length product for 25 additional cycles. The full length product was gel purified and cloned into pCR-blunt (Invitrogen) and individual clones were screened by sequencing. The Min insert was isolated as an NheI-KpnI fragment and cloned into the same sites of pcDNA3.1(-)/Myc-His A (Invitrogen) for expression. The Min protein contains the Myc and His antibody epitope tags at its carboxyl-terminal end.

For all the PCR reactions described, a total of 30 cycles were performed using Pfu polymerase and the following conditions: 95°C for 15 seconds, annealing temperature for 30 seconds, 72°C for one minute. The annealing temperature used was 5°C below the lowest calculated  $T_m$  of each primer pair.

Three changes to pMIN.0 were made to produce pMIN.1 (Figure 21; SEQ ID NO:37, also referred to as pMIN-AOS). The mouse ova epitope was removed, the position 9 alanine anchor residue (#547) of HBV pol 551-560 was converted to a valine which increased the *in vitro* binding affinity 40-fold, and a translational stop codon was introduced at the end of the multi-epitope coding sequence. The changes were made by amplifying two overlapping fragments and combining them to yield the full length product.

The first reaction used the 5' pcDNA vector primer T7 and the primer Min-ovaR (nucleotides 247-218) TGGACAGTCCCACTCCAGCACCACGTCAT (SEQ ID NO:120). The 3' half was amplified with the primers: Min-ovaF (nucleotides 228-257) GCTGGGAGTGGGACTGTCCAGTGACGTGGC (SEQ ID NO:121) and Min-StopR (nucleotides 390-361) GGTACCTCACACAAAGGGCACCAGCAGGC (SEQ ID NO:122)

The two fragments were gel purified, mixed, denatured, annealed, and filled in with five cycles of PCR. The full length fragment was amplified with the flanking primers T7 and Min-Stop for 25 more cycles. The product was gel purified, digested with NheI and KpnI and cloned into pcDNA3.1 for sequencing and expression. The insert from pMin.1 was isolated as an NheI-KpnI fragment and cloned into pEP2 to make pEP2-AOS.

EXAMPLE II: Assay for T Helper Cell Activation

This example shows methods for assaying T helper cell activity. One method for assaying T helper cell activity uses spleen cells of an immunized organism.

- 5 Briefly, a spleen cell pellet is suspended with 2-3 ml of red blood cell lysis buffer containing 8.3 g/liter ammonium chloride in 0.001 M Tris-HCl, pH 7.5. The cells are incubated in lysis buffer for 3-5 min at room temperature with occasional vortexing. An excess volume of 50 ml of R10 medium is added to the cells, and the cells are pelleted. The cells are resuspended and pelleted one or two more times in R2 medium or R10  
10 medium.

The cell pellet is suspended in R10 medium and counted. If the cell suspension is aggregated, the aggregates are removed by filtration or by allowing the aggregates to settle by gravity. The cell concentration is brought to  $10^7$ /ml, and 100  $\mu$ l of spleen cells are added to 96 well flat bottom plates.

- 15 Dilutions of the appropriate peptide, such as pan DR epitope (SEQ ID NO:145), are prepared in R10 medium at 100, 10, 1, 0.1 and 0.01  $\mu$ g/ml, and 100  $\mu$ l of peptide are added to duplicate or triplicate wells of spleen cells. The final peptide concentration is 50, 5, 0.5, 0.05 and 0.005  $\mu$ g/ml. Control wells receive 100  $\mu$ l R10 medium.

- 20 The plates are incubated for 3 days at 37°C. After 3 days, 20  $\mu$ l of 50  $\mu$ Ci/ml  $^3$ H-thymidine is added per well. Cells are incubated for 18-24 hours and then harvested onto glass fiber filters. The incorporation of  $^3$ H-thymidine into DNA of proliferating cells is measured in a beta counter.

- A second assay for T helper cell activity uses peripheral blood  
25 mononuclear cells (PBMC) that are stimulated *in vitro* as described in Alexander *et al.*, *supra* and Sette (WO 95/07,707), as adapted from Manca *et al.*, *J. Immunol.* 146:1964-1971 (1991), which is incorporated herein by reference. Briefly, PBMC are collected from healthy donors and purified over Ficoll-Plaque (Pharmacia Biotech; Piscataway, NJ). PBMC are plated in a 24 well tissue culture plate at  $4 \times 10^6$  cells/ml. Peptides are  
30 added at a final concentration of 10  $\mu$ g/ml. Cultures are incubated at 37°C in 5% CO<sub>2</sub>.

On day 4, recombinant interleukin-2 (IL-2) is added at a final concentration of 10 ng/ml. Cultures are fed every 3 days by aspirating 1 ml of medium and replacing with fresh medium containing IL-2. Two additional stimulations of the T cells with antigen are performed on approximately days 14 and 28. The T cells (3 x

10<sup>5</sup>/well) are stimulated with peptide (10 µg/ml) using autologous PBMC cells (2 x 10<sup>6</sup> irradiated cells/well) (irradiated with 7500 rads) as antigen-presenting cells in a total of three wells of a 24 well tissue culture plate. In addition, on day 14 and 28, T cell proliferative responses are determined under the following conditions: 2 x 10<sup>4</sup> T cells/well; 1 x 10<sup>5</sup> irradiated PBMC/well as antigen-presenting cells; peptide concentration varying between 0.01 and 10 µg/ml final concentration. The proliferation of the T cells is measured 3 days later by the addition of <sup>3</sup>H-thymidine (1 µCi/well) 18 hr prior to harvesting the cells. Cells are harvested onto glass filters and <sup>3</sup>H-thymidine incorporation is measured in a beta plate counter. These results demonstrate methods for assaying T helper cell activity by measuring <sup>3</sup>H-thymidine incorporation.

#### EXAMPLE III: Assay for Cytotoxic T Lymphocyte Response

This example shows a method for assaying cytotoxic T lymphocyte (CTL) activity. A CTL response is measured essentially as described previously (Vitiello *et al.*, *Eur. J. Immunol.* 27:671-678 (1997), which is incorporated herein by reference). Briefly, after approximately 10-35 days following DNA immunization, splenocytes from an animal are isolated and co-cultured at 37°C with syngeneic, irradiated (3000 rad) peptide-coated LPS blasts (1 x 10<sup>6</sup> to 1.5 x 10<sup>6</sup> cells/ml) in 10 ml R10 in T25 flasks. LPS blasts are obtained by activating splenocytes (1 x 10<sup>6</sup> to 1.5 x 10<sup>6</sup> cells/ml) with 25 µg/ml lipopolysaccharides (LPS) (Sigma cat. no. L-2387; St. Louis, MO) and 7 µg/ml dextran sulfate (Pharmacia Biotech) in 30 ml R10 medium in T75 flasks for 3 days at 37°C. The lymphoblasts are then resuspended at a concentration of 2.5 x 10<sup>7</sup> to 3.0 x 10<sup>7</sup>/ml, irradiated (3000 rad), and coated with the appropriate peptides (100µg/ml) for 1 h at 37°C. Cells are washed once, resuspended in R10 medium at the desired concentration and added to the responder cell preparation. Cultures are assayed for cytolytic activity on day 7 in a <sup>51</sup>Cr-release assay.

For the <sup>51</sup>Cr-release assay, target cells are labeled for 90 min at 37°C with 150 µl sodium <sup>51</sup>chromate (<sup>51</sup>Cr) (New England Nuclear; Wilmington DE), washed three times and resuspended at the appropriate concentration in R10 medium. For the assay, 10<sup>4</sup> target cells are incubated in the presence of different concentrations of effector cells in a final volume of 200 µl in U-bottom 96 well plates in the presence or absence of 10 µg/ml peptide. Supernatants are removed after 6 h at 37°C, and the percent specific lysis is determined by the formula: percent specific lysis = 100 x (experimental release - spontaneous release) / (maximum release - spontaneous release). To facilitate comparison

of responses from different experiments, the percent release data is transformed to lytic units  $30 \text{ per } 10^6 \text{ cells (LU30/10}^6\text{)}$ , with 1 LU30 defined as the number of effector cells required to induce 30% lysis of  $10^6$  target cells in a 6 h assay. LU values represent the LU30/ $10^6$  obtained in the presence of peptide minus LU30/ $10^6$  in the absence of peptide.

- 5 These results demonstrate methods for assaying CTL activity by measuring  $^{51}\text{Cr}$  release from cells.

EXAMPLE IV: T Cell Proliferation in Mice Immunized with Expression Vectors  
Encoding MHC Class II Epitopes and MHC Class II Targeting Sequences

- 10 This example demonstrates that expression vectors encoding MHC class II epitopes and MHC class II targeting sequences are effective at activating T cells.

The constructs used in the T cell proliferation assay are described in Example I and were cloned into the vector pEP2, a CMV driven expression vector. The peptides used for T cell *in vitro* stimulation are: Ova 323-339, ISQAVHAAHAEINEAGR (SEQ ID NO:123); HBVcore128, TPPAYRPPNAPILF (SEQ ID NO:124); HBVenv182, FLLLRILRLTPQSLD (SEQ ID NO:125); and PADRE, AKFVAAWTLKAAA (SEQ ID NO:38).

- 20 T cell proliferation was assayed essentially as described in Example II. Briefly, 12 to 16 week old B6D2 F1 mice (2 mice per construct) were injected with 100  $\mu\text{g}$  of the indicated expression vector (50  $\mu\text{g}$  per leg) in the anterior tibialis muscle. After eleven days, spleens were collected from the mice and separated into a single cell suspension by Dounce homogenization. The splenocytes were counted and one million splenocytes were plated per well in a 96-well plate. Each sample was done in triplicate. Ten  $\mu\text{g/ml}$  of the corresponding peptide encoded by the respective expression vectors was added to each well. One well contained splenocytes without peptide added for a negative control. Cells were cultured at  $37^\circ\text{C}$ , 5%  $\text{CO}_2$  for three days.

- 30 After three days, one  $\mu\text{Ci}$  of  $^3\text{H}$ -thymidine was added to each well. After 18 hours at  $37^\circ\text{C}$ , the cells were harvested onto glass filters and  $^3\text{H}$  incorporation was measured on an LKB  $\beta$  plate counter. The results of the T cell proliferation assay are shown in Table 9. Antigen specific T cell proliferation is presented as the stimulation index (SI); this is defined as the ratio of the average  $^3\text{H}$ -thymidine incorporation in the presence of antigen divided by the  $^3\text{H}$ -thymidine incorporation in the absence of antigen.

The immunogen "PADRE + IFA" is a positive control where the PADRE peptide in incomplete Freund's adjuvant was injected into the mice and compared to the

response seen by injecting the MHC class II epitope constructs containing a PADRE sequence. As shown in Table 9, most of the expression vectors tested were effective at activating T cell proliferation in response to the addition of PADRE peptide. The activity of several of the expression vectors was comparable to that seen with immunization with the PADRE peptide in incomplete Freund's adjuvant. The expression vectors containing both MHC class I and MHC class II epitopes, pEP2-AOS and pcDNA-AOS, were also effective at activating T cell proliferation in response to the addition of PADRE peptide.

These results show that expression vectors encoding MHC class II epitopes fused to a MHC class II targeting sequence is effective at activating T cell proliferation and are useful for stimulating an immune response.

#### EXAMPLE V: *In vivo* assay Using Transgenic Mice

##### *A. Materials and methods*

Peptides were synthesized according to standard F-moc solid phase

- synthesis methods which have been previously described (Ruppert *et al.*, *Cell* 74:929 (1993); Sette *et al.*, *Mol. Immunol.* 31:813 (1994)). Peptide purity was determined by analytical reverse-phase HPLC and purity was routinely >95%. Synthesis and purification of the Theradigm-HBV lipopeptide vaccine is described in (Vitiello *et al.*, *J. Clin. Invest.* 95:341 (1995)).

##### Mice

HLA-A2.1 transgenic mice used in this study were the F1 generation derived by crossing transgenic mice expressing a chimeric gene consisting of the  $\alpha 1$ ,  $\alpha 2$  domains of HLA-A2.1 and  $\alpha 3$  domain of H-2K<sup>b</sup> with SJL/J mice (Jackson Laboratory, Bar Harbor, ME). This strain will be referred to hereafter as HLA-A2.1/K<sup>b</sup>-H-2<sup>bxs</sup>. The parental HLA-A2.1/K<sup>b</sup> transgenic strain was generated on a C57BL/6 background using the transgene and methods described in (Vitiello *et al.*, *J. Exp. Med.* 173:1007 (1991)). HLA-A11/K<sup>b</sup> transgenic mice used in the current study were identical to those described in (Alexander *et al.*, *J. Immunol.* 159:4753 (1997)).

##### Cell lines, MHC purification, and peptide binding assay

Target cells for peptide-specific cytotoxicity assays were Jurkat cells transfected with the HLA-A2.1/K<sup>b</sup> chimeric gene (Vitiello *et al.*, *J. Exp. Med.* 173:1007

(1991)) and .221 tumor cells transfected with HLA-A11/K<sup>b</sup> (Alexander *et al.*, *J. Immunol.* 159:4753 (1997)).

To measure presentation of endogenously processed epitopes, Jurkat-A2.1/K<sup>b</sup> cells were transfected with the pMin.1 or pMin.2-GFP minigenes then tested in a cytotoxicity assay against epitope-specific CTL lines. For transfection, Jurkat-A2.1/K<sup>b</sup> cells were resuspended at 10<sup>7</sup> cells/ml and 30 µg of DNA was added to 600 µl of cell suspension. After electroporating cells in a 0.4 cm cuvette at 0.25 kV, 960 µF, cells were incubated on ice for 10 min then cultured for 2 d in RPMI culture medium. Cells were then cultured in medium containing 200 U/ml hygromycin B (Calbiochem, San Diego CA) to select for stable transfectants. FACS was used to enrich the fraction of green fluorescent protein (GFP)-expressing cells from 15% to 60% (data not shown).

Methods for measuring the quantitative binding of peptides to purified HLA-A2.1 and -A11 molecules is described in Ruppert *et al.*, *Cell* 74:929 (1993); Sette *et al.*, *Mol. Immunol.* 31:813 (1994); Alexander *et al.*, *J. Immunol.* 159:4753 (1997).

All tumor cell lines and splenic CTLs from primed mice were grown in culture medium (CM) that consisted of RPMI 1640 medium with HEPES (Life Technologies, Grand Island, NY) supplemented with 10% FBS, 4 mM L-glutamine, 5 X 10<sup>-5</sup> M 2-ME, 0.5 mM sodium pyruvate, 100 µg/ml streptomycin, and 100 U/ml penicillin.

#### Construction of minigene multi-epitope DNA plasmids

pMIN.0 and pMIN.1 (i.e., pMIN-AOS) were constructed as described above and in USSN 60/085,751.

pMin.1-No PADRE and pMin.1-Anchor. pMin.1 was amplified using two overlapping fragments which was then combined to yield the full length product. The first reaction used the 5' pcDNA vector primer T7 and either primer ATCGCTAGGCAGGAACCTTATACAGGATTCC (SEQ ID NO:126) for pMin.1-No PADRE or TGGACAGTCCGGCTCCAGCACCACGT (SEQ ID NO:127) for pMin.1-Anchor. The 3' half was amplified with the primers TTCTGCGCTAGCGATTTC (SEQ ID NO:128) (No PADRE) or GCTGGGAGCCGGACTGTCCAGGTACGT (SEQ ID NO:129) (Anchor) and Min-StopR. The two fragments generated from amplifying the 5' and 3' ends were gel purified, mixed, denatured, annealed, and filled in with five cycles

of PCR. The full length fragment was further amplified with the flanking primers T7 and Min-StopR for 25 more cycles.

- 5 pMin.1-No Sig. The Ig signal sequence was deleted from pMin.1 by PCR amplification with primer GCTAGCGCCGCCACCATGCACACCCTGTGGAAGGC CGGAATC (SEQ ID NO:130) and pcDNA rev (Invitrogen) primers. The product was cloned into PCR-blunt and sequenced.

- 10 pMin.1-Switch. Three overlapping fragments were amplified from pMin.1, combined, and extended. The 5' fragment was amplified with the vector primer T7 and primer GGGCACCAGCAGGCTCAGCCACACTCCCAGCACCACGTC (SEQ ID NO:131). The second overlapping fragment was amplified with primers AGCCTGCTGGTGCCCTTTGTGATCCTGAAGGAGCCTGTGC (SEQ ID NO:132) and AGCCACGTACCTGGACAGTCCCTTCCACACAGGCACTCCAT (SEQ ID NO:133). Primer TGTCCAGGTACGTGGCTAGGCTGTGAGGTACC (SEQ ID NO:134) and the vector primer pcDNA rev (Invitrogen) were used to amplify the third (3') fragment. Fragments 1, 2, and 3 were amplified and gel purified. Fragments 2 and 3 were mixed, annealed, amplified, and gel purified. Fragment 1 was combined with the product of 2 and 3, and extended, gel purified and cloned into pcDNA3.1 for expression.

20

- pMin.2-GFP. The signal sequence was deleted from pMin.0 by PCR amplification with Min.0-No Sig-5' plus pcDNA rev (Invitrogen) primers GCTAGCGCCGCCACCATGCACACCCTGTGGAAGGCCGGAATC (SEQ ID NO:135). The product was cloned into PCR-blunt and sequenced. The insert containing the open reading frame of the signal sequence-deleted multi-epitope construct was cut out with *NheI* plus *HindIII* and ligated into the same sites of pEGFPN1 (Clontech). This construct fuses the coding region of the signal-deleted pMin.0 construct to the N-terminus of green fluorescent protein (GFP).

30

#### Immunization of mice

For DNA immunization, mice were pretreated by injecting 50  $\mu$ l of 10  $\mu$ M cardiotoxin (Sigma Chem. Co., #C9759) bilaterally into the tibialis anterior muscle. Four or five days later, 100  $\mu$ g of DNA diluted in PBS were injected in the same muscle.

Theradigm-HBV lipopeptide (10 mg/ml in DMSO) that was stored at -20°C, was thawed for 10 min at 45°C before being diluted 1:10 (v/v) with room temperature PBS. Immediately upon addition of PBS, the lipopeptide suspension was vortexed vigorously and 100 µl was injected s.c. at the tail base (100 µg/mouse).

- 5 Immunogenicity of individual CTL epitopes was tested by mixing each CTL epitope (50 µg/mouse) with the HBV core 128-140 peptide (TPPAYRPPNAPIL (SEQ ID NO:124), 140 µg/mouse) which served to induce I-A<sup>b</sup>-restricted Th cells. The peptide cocktail was then emulsified in incomplete Freund's adjuvant (Sigma Chem. Co.) and 100 µl of peptide emulsion was injected s.c. at the tail base.

10

*In vitro* CTL cultures and cytotoxicity assays

- Eleven to 14 days after immunization, animals were sacrificed and a single cell suspension of splenocytes prepared. Splenocytes from cDNA-primed animals were stimulated *in vitro* with each of the peptide epitopes represented in the minigene.
- 15 Splenocytes (2.5-3.0 X 10<sup>7</sup>/flask) were cultured in upright 25 cm<sup>2</sup> flasks in the presence of 10 µg/ml peptide and 10<sup>7</sup> irradiated spleen cells that had been activated for 3 days with LPS (25 µg/ml) and dextran sulfate (7 µg/ml). Triplicate cultures were stimulated with each epitope. Five days later, cultures were fed with fresh CM. After 10 d of *in vitro* culture, 2-4 X 10<sup>6</sup> CTLs from each flask were restimulated with 10<sup>7</sup> LPS/dextran sulfate-
- 20 activated splenocytes treated with 100 µg/ml peptide for 60-75 min at 37°C, then irradiated 3500 rads. CTLs were restimulated in 6-well plates in 8 ml of cytokine-free CM. Eighteen hr later, cultures received cytokines contained in con A-activated splenocyte supernatant (10-15% final concentration, v/v) and were fed or expanded on the third day with CM containing 10-15% cytokine supernate. Five days after restimulation,
- 25 CTL activity of each culture was measured by incubating varying numbers of CTLs with 10<sup>4</sup> <sup>51</sup>Cr-labelled target cells in the presence or absence of peptide. To decrease nonspecific cytotoxicity from NK cells, YAC-1 cells (ATCC) were also added at a YAC-1:<sup>51</sup>Cr-labeled target cell ratio of 20:1. CTL activity against the HBV Pol 551 epitope was measured by stimulating DNA-primed splenocytes *in vitro* with the native A-
- 30 containing peptide and testing for cytotoxic activity against the same peptide.

To more readily compare responses, the standard E:T ratio vs % cytotoxicity data curves were converted into LU per 10<sup>6</sup> effector cells with one LU defined as the lytic activity required to achieve 30% lysis of target cells at a 100:1 E:T



- ratio. Specific CTL activity ( $\Delta$ LU) was calculated by subtracting the LU value obtained in the absence of peptide from the LU value obtained with peptide. A given culture was scored positive for CTL induction if all of the following criteria were met: 1)  $\Delta$ LU  $> 2$ ; 2) LU(+ peptide) + LU(- peptide)  $> 3$ ; and 3) a  $> 10\%$  difference in % cytotoxicity tested with and without peptide at the two highest E:T ratios (starting E:T ratios were routinely between 25-50:1).

- CTL lines were generated from pMin.1-primed splenocytes through repeated weekly stimulations of CTLs with peptide-treated LPS/DxS-activated splenocytes using the 6-well culture conditions described above with the exception that CTLs were expanded in cytokine-containing CM as necessary during the seven day stimulation period.

#### Cytokine assay

- To measure IFN- $\gamma$  production in response to minigene-transfected target cells,  $4 \times 10^4$  CTLs were cultured with an equivalent number of minigene-transfected Jurkat-A2.1/K<sup>b</sup> cells in 96-well flat bottom plates. After overnight incubation at 37°C, culture supernatant from each well was collected and assayed for IFN- $\gamma$  concentration using a sandwich ELISA. Immulon II microtiter wells (Dynatech, Boston, MA) were coated overnight at 4°C with 0.2  $\mu$ g of anti-mouse IFN- $\gamma$  capture Ab, R4-6A2 (Pharmingen). After washing wells with PBS/0.1% Tween-20 and blocking with 1% BSA, Ab-coated wells were incubated with culture supernate samples for 2 hr at room temperature. A secondary anti-IFN- $\gamma$  Ab, XMG1.2 (Pharmingen), was added to wells and allowed to incubate for 2 hr at room temperature. Wells were then developed by incubations with Avidin-DH and finally with biotinylated horseradish peroxidase H (Vectastain ABC kit, Vector Labs, Burlingame, CA) and TMB peroxidase substrate (Kirkegaard and Perry Labs, Gaithersburg, MD). The amount of cytokine present in each sample was calculated using a rIFN- $\gamma$  standard (Pharmingen).

#### *b. Results*

- Selection of epitopes and minigene construct design

In the first series of experiments, the issue was whether a balanced multispecific CTL response could be induced by simple minigene cDNA constructs that encode several dominant HLA class I-restricted epitopes. Accordingly, nine CTL

epitopes were chosen on the basis of their relevance in CTL immunity during HBV and HIV infection in humans, their sequence conservancy among viral subtypes, and their class I MHC binding affinity (Table 10). Of these nine epitopes, six are restricted by HLA-A2.1 and three showed HLA-A11-restriction. One epitope, HBV Pol 551, was studied in two alternative forms: either the wild type sequence or an analog (HBV Pol 551-V) engineered for higher binding affinity.

As referenced in Table 10, several independent laboratories have reported that these epitopes are part of the dominant CTL response during HBV or HIV infection. All of the epitopes considered showed greater than 75% conservancy in primary amino acid sequence among the different HBV subtypes and HIV clades. The MHC binding affinity of the peptides was also considered in selection of the epitopes. These experiment addressed the feasibility of immunizing with epitopes possessing a wide range of affinities and, as shown in Table 10, the six HBV and three HIV HLA-restricted epitopes covered a spectrum of MHC binding affinities spanning over two orders of magnitude, with IC<sub>50</sub> concentrations ranging from 3 nM to 200 nM.

The immunogenicity of the six A2.1- and three A11-restricted CTL epitopes in transgenic mice was verified by co-immunization with a helper T cell peptide in an IFA formulation. All of the epitopes induced significant CTL responses in the 5 to 73  $\Delta$ LU range (Table 10). As mentioned above, to improve the MHC binding and immunogenicity of HBV Pol 551, the C-terminal A residue of this epitope was substituted with V resulting in a dramatic 40-fold increase in binding affinity to HLA-A2.1 (Table 10). While the parental sequence was weakly or nonimmunogenic in HLA transgenic mice, the HBV Pol 551-V analog induced significant levels of CTL activity when administered in IFA (Table 10). On the basis of these results, the V analog of the HBV Pol 551 epitope was selected for the initial minigene construct. In all of the experiments reported herein, CTL responses were measured with target cells coated with the native HBV Pol 551 epitope, irrespective of whether the V analog or native epitope was utilized for immunization.

Finally, since previous studies indicated that induction of T cell help significantly improved the magnitude and duration of CTL responses (Vitiello *et al.*, *J. Clin. Invest.* 95:341 (1995); Livingston *et al.*, *J. Immunol.* 159:1383 (1997)), the universal Th cell epitope PADRE was also incorporated into the minigene. PADRE has been shown previously to have high MHC binding affinity to a wide range of mouse and

human MHC class II haplotypes (Alexander *et al.*, *Immunity* 1:751 (1994)). In particular, it has been previously shown that PADRE is highly immunogenic in H-2<sup>b</sup> mice that are used in the current study (Alexander *et al.*, *Immunity* 1:751 (1994)).

- pMin.1, the prototype cDNA minigene construct encoding nine CTL epitopes and PADRE, was synthesized and subcloned into the pcDNA3.1 vector. The position of each of the nine epitopes in the minigene was optimized to avoid junctional mouse H-2<sup>b</sup> and HLA-A2.1 class I MHC epitopes. The mouse Ig  $\kappa$  signal sequence was also included at the 5' end of the construct to facilitate processing of the CTL epitopes in the endoplasmic reticulum (ER) as reported by others (Anderson *et al.*, *J. Exp. Med.* 174:489 (1991)). To avoid further conformational structure in the translated polypeptide gene product that may affect processing of the CTL epitopes, an ATG stop codon was introduced at the 3' end of the minigene construct upstream of the coding region for c-myc and poly-his epitopes in the pcDNA3.1 vector.

#### 15 Immunogenicity of pMin.1 in transgenic mice

- To assess the capacity of the pMin.1 minigene construct to induce CTLs *in vivo*, HLA-A2.1/K<sup>b</sup>-H-2<sup>bxs</sup> transgenic mice were immunized intramuscularly with 100  $\mu$ g of naked cDNA. As a means of comparing the level of CTLs induced by cDNA immunization, a control group of animals was also immunized with Theradigm-HBV, a
- 20 palmitoylated lipopeptide consisting of the HBV Core 18 CTL epitope linked to the tetanus toxin 830-843 Th cell epitope.

- Splenocytes from immunized animals were stimulated twice with each of the peptide epitopes encoded in the minigene, then assayed for peptide-specific cytotoxic activity in a <sup>51</sup>Cr release assay. A representative panel of CTL responses of pMin.1-primed splenocytes, shown in Figure 22, clearly indicates that significant levels of CTL
- 25 induction were generated by minigene immunization. The majority of the cultures stimulated with the different epitopes exceeded 50% specific lysis of target cells at an E:T ratio of 1:1. The results of four independent experiments, compiled in Table 11, indicate that the pMin.1 construct is indeed highly immunogenic in HLA-A2.1/K<sup>b</sup>-H-2<sup>bxs</sup>
- 30 transgenic mice, inducing a broad CTL response directed against each of its six A2.1-restricted epitopes.

To more conveniently compare levels of CTL induction among the different epitopes, the % cytotoxicity values for each splenocyte culture was converted to

$\Delta$ LU and the mean  $\Delta$ LU of CTL activity in positive cultures for each epitope was determined (see Example V, materials and methods, for positive criteria). The data, expressed in this manner in Table 11, confirms the breadth of CTL induction elicited by pMin.1 immunization since extremely high CTL responses, ranging between 50 to 700  $\Delta$ LU, were observed against the six A2.1-restricted epitopes. More significantly, the responses of several hundred  $\Delta$ LU observed for five of the six epitopes approached or exceeded that of the Theradigm-HBV lipopeptide, a vaccine formulation known for its high CTL-inducing potency (Vitiello *et al.*, *J. Clin. Invest.* 95:341 (1995); Livingston *et al.*, *J. Immunol.* 159:1383 (1997)). The HBV Env 335 epitope was the only epitope showing a lower mean  $\Delta$ LU response compared to lipopeptide (Table 11, 44 vs 349  $\Delta$ LU).

#### Processing of minigene epitopes by transfected cells

The decreased CTL response observed against HBV Env 335 was somewhat unexpected since this epitope had good A2.1 binding affinity (IC50%, 5 nM) and was also immunogenic when administered in IFA. The lower response may be due, at least in part, to the inefficient processing of this epitope from the minigene polypeptide by antigen presenting cells following *in vivo* cDNA immunization. To address this possibility, Jurkat-A2.1 K<sup>b</sup> tumor cells were transfected with pMin.1 cDNA and the presentation of the HBV Env 335 epitope by transfected cells was compared to more immunogenic A2.1-restricted epitopes using specific CTL lines. Epitope presentation was also studied using tumor cells transfected with a control cDNA construct, pMin.2-GFP, that encoded a similar multi-epitope minigene fused with GFP which allows detection of minigene expression in transfected cells by FACS.

Epitope presentation of the transfected Jurkat cells was analyzed using specific CTL lines, with cytotoxicity or IFN- $\gamma$  production serving as a read-out. It was found that the levels of CTL response correlated directly with the *in vivo* immunogenicity of the epitopes. Highly immunogenic epitopes *in vivo*, such as HBV Core 18, HIV Pol 476, and HBV Pol 455, were efficiently presented to CTL lines by pMin.1- or pMin.2-GFP-transfected cells as measured by IFN- $\gamma$  production (Figure 23A, >100 pg/ml for each epitope) or cytotoxic activity (Figure 23C, >30% specific lysis). In contrast to these high levels of *in vitro* activity, the stimulation of the HBV Env 335-specific CTL line against both populations of transfected cells resulted in less than 12 pg/ml IFN- $\gamma$  and 3% specific

lysis. Although the HBV Env 335-specific CTL line did not recognize the naturally processed epitope efficiently, this line did show an equivalent response to peptide-loaded target cells, as compared to CTL lines specific for the other epitopes (Figure 23B, D).

- Collectively, these results suggest that a processing and/or presentation defect associated with the HBV Env 335 epitope that may contribute to its diminished immunogenicity *in vivo*.

#### Effect of the helper T cell epitope PADRE on minigene immunogenicity

- Having obtained a broad and balanced CTL response in transgenic mice immunized with a minigene cDNA encoding multiple HLA-A2.1-restricted epitopes, next possible variables were examined that could influence the immunogenicity of the prototype construct. This type of analysis could lead to rational and rapid optimization of future constructs. More specifically, a cDNA construct based on the pMin.1 prototype was synthesized in which the PADRE epitope was deleted to examine the contribution of T cell help in minigene immunogenicity (Figure 24A).

- The results of the immunogenicity analysis indicated that deletion of the PADRE Th cell epitope resulted in significant decreases in the frequency of specific CTL precursors against four of the minigene epitopes (HBV Core 18, HIV Env 120, HBV Pol 455, and HBV Env 335) as indicated by the 17 to 50% CTL-positive cultures observed against these epitopes compared to the 90-100% frequency in animals immunized with the prototype pMin.1 construct (Figure 25). Moreover, for two of the epitopes, HBV Core 18 and HIV Env 120, the magnitude of response in positive cultures induced by pMin.1-No PADRE was 20- to 30-fold less than that of the pMin.1 construct (Figure 25A).

#### Effect of modulation of MHC binding affinity on epitope immunogenicity

Next a construct was synthesized in which the V anchor residue in HBV Pol 551 was replaced with alanine, the native residue, to address the effect of decreasing MHC binding on epitope immunogenicity (Figure 24B).

- Unlike deletion of the Th cell epitope, decreasing the MHC binding capacity of the HBV Pol 551 epitope by 40-fold through modification of the anchor residue did not appear to affect epitope immunogenicity (Figure 25B). The CTL response against the HBV Pol 551 epitope, as well as to the other epitopes, measured either by LU or frequency of CTL-positive cultures, was very similar between the constructs

- containing the native A or improved V residue at the MHC binding anchor site. This finding reinforces the notion that minimal epitope minigenes can efficiently deliver epitopes of vastly different MHC binding affinities. Furthermore, this finding is particularly relevant to enhancing epitope immunogenicity via different delivery methods, especially in light of the fact that the wild type HBV Pol 551 epitope was essentially nonimmunogenic when delivered in a less potent IFA emulsion.

#### Effect of the signal sequence on minigene construct immunogenicity

- The signal sequence was deleted from the pMin.1 construct, thereby preventing processing of the minigene polypeptide in the ER (Figure 24C). When the immunogenicity of the pMin.1-No Sig construct was examined, an overall decrease in response was found against four CTL epitopes. Two of these epitopes, HIV Env 120 and HBV Env 335, showed a decrease in frequency of CTL-positive cultures compared to pMin.1 while the remaining epitopes, HBV Pol 455 and HIV Pol 476, showed a 16-fold (from 424 to 27  $\Delta$ LU) and 3-fold decrease (709 to 236  $\Delta$ LU) in magnitude of the mean CTL response, respectively (Figure 25C). These findings suggest that allowing ER-processing of some of the epitopes encoded in the pMin.1 prototype construct may improve immunogenicity, as compared with constructs that allow only cytoplasmic processing of the same panel of epitopes.

#### Effect of epitope rearrangement and creation of new junctional epitopes

- In the final construct tested, the immunogenicity of the HBV Env 335 epitope was analyzed to determine whether it may be influenced by its position at the 3' terminus of the minigene construct (Figure 24D). Thus, the position of the Env epitope in the cDNA construct was switched with a more immunogenic epitope, HBV Pol 455, located in the center of the minigene. It should be noted that this modification also created two potentially new epitopes. As shown in Figure 25D, the transposition of the two epitopes appeared to affect the immunogenicity of not only the transposed epitopes but also more globally of other epitopes. Switching epitopes resulted in obliteration of CTL induction against HBV Env 335 (no positive cultures detected out of six). The CTL response induced by the terminal HBV Pol 455 epitope was also decreased but only slightly (424 vs 78 mean  $\Delta$ LU). In addition to the switched epitopes, CTL induction against other epitopes in the pMin.1-Switch construct was also markedly reduced

compared to the prototype construct. For example, a CTL response was not observed against the HIV Env 120 epitope and it was significantly diminished against the HBV Core 18 (4 of 6 positive cultures, decrease in mean  $\Delta$ LU from 306 to 52) and HBV Pol 476 (decrease in mean  $\Delta$ LU from 709 to 20) epitopes (Figure 25D).

5 As previously mentioned, it should be noted that switching the two epitopes had created new junctional epitopes. Indeed, in the pMin.1-Switch construct, two new potential CTL epitopes were created from sequences of HBV Env 335-HIV Pol 476 (LLVPFVIL (SEQ ID NO:135), H-2K<sup>b</sup>-restricted) and HBV Env 335-HBV Pol 551 (VLGVWLSLLV (SEQ ID NO:136), HLA-A2.1-restricted) epitopes. Although these  
10 junctional epitopes have not been examined to determine whether or not they are indeed immunogenic, this may account for the low immunogenicity of the HBV Env 335 and HIV Pol 476 epitopes. These findings suggest that avoiding junctional epitopes may be important in designing multi-epitope minigenes as is the ability to confirm their immunogenicity *in vivo* in a biological assay system such as HLA transgenic mice.

15

#### Induction of CTLs against A11 epitopes encoded in pMin.1

To further examine the flexibility of the minigene vaccine approach for inducing a broad CTL response against not only multiple epitopes but also against epitopes restricted by different HLA alleles, HLA-A11/K<sup>b</sup> transgenic mice were  
20 immunized to determine whether the three A11 epitopes in the pMin.1 construct were immunogenic for CTLs, as was the case for the A2.1-restricted epitopes in the same construct. As summarized in Table 12, significant CTL induction was observed in a majority of cultures against all three of the HLA-A11-restricted epitopes and the level of CTL immunity induced for the three epitopes, in the range of 40 to 260  $\Delta$ LU, exceeded  
25 that of peptides delivered in IFA (Table 10). Thus, nine CTL epitopes of varying HLA restrictions incorporated into a prototype minigene construct all demonstrated significant CTL induction *in vivo*, confirming that minigene DNA plasmids can serve as means of delivering multiple epitopes, of varying HLA restrictions and MHC binding affinities, to the immune system in an immunogenic fashion and that appropriate transgenic mouse  
30 strains can be used to measure DNA construct immunogenicity *in vivo*.

CTLs were also induced against three A11 epitopes in A11/K<sup>b</sup> transgenic mice. These responses suggest that minigene delivery of multiple CTL epitopes that confers broad population coverage may be possible in humans and that transgenic animals

- of appropriate haplotypes may be a useful tools in optimizing the *in vivo* immunogenicity of minigene DNA. In addition, animals such as monkeys having conserved HLA molecules with cross reactivity to CTL and HTL epitopes recognized by human MHC molecules can be used to determine human immunogenicity of HTL and CTL epitopes
- 5 (Bertoni *et al.*, *J. Immunol.* 161:4447-4455 (1998)).

- This study represents the first description of the use of HLA transgenic mice to quantitate the *in vivo* immunogenicity of DNA vaccines, by examining response to epitopes restricted by human HLA antigens. *In vivo* studies are required to address the variables crucial for vaccine development, that are not easily evaluated by *in vitro* assays,
- 10 such as route of administration, vaccine formulation, tissue biodistribution, and involvement of primary and secondary lymphoid organs. Because of its simplicity and flexibility, HLA transgenic mice represent an attractive alternative, at least for initial vaccine development studies, compared to more cumbersome and expensive studies in higher animal species, such as nonhuman primates. The *in vitro* presentation studies
- 15 described above further supports the use of HLA transgenic mice for screening DNA constructs containing human epitopes inasmuch as a direct correlation between *in vivo* immunogenicity and *in vitro* presentation was observed. Finally, strong CTL responses were observed against all six A 2.1 restricted viral epitopes and in three A11 restricted epitopes encoded in the prototype pMin.1 construct. For five of the A 2.1 restricted
- 20 epitopes, the magnitude of CTL response approximated that observed with the lipopeptide, Theradigm-HBV, that previously was shown to induce strong CTL responses in humans (Vitiello *et al.*, *J. Clin. Invest.* 95:341 (1995); Livingston *et al.*, *J. Immunol.* 159:1383 (1997)).



Table 1  
HBV derived HTL epitopes

Pepide	Sequence	Source	SEQ ID NO.
1298.06	KQAFITSPITYKALFC	HBV POL 661	
F107.03	LQSLTNLLSSNLSWL	HBV POL 412	
1280.06	AGFELLTRLTIPOS	HBV ENV 180	
1280.09	GTSFYVVISALNPAD	HBV POL 774	
CF-08	VSEGYWIRTPAYRPINAPH	HBV NLIC 120	
27.0280	GVWIRTPAYRPINA	HBV NLIC 121	
1186.25	SHGVWIRTPAYRPIN	HBV NLIC 121	
27.0281	RHYLHLLWKAGILYK	HBV POL 145	
F107.04	PELLAQFTSACSVV	HBV POL 523	
1186.15	LVPFQVWFVGLSFTV	HBV ENV 339	
1280.15	LHLVSHPIILGFRKI	HBV POL 501	
1298.04	KQCFRKL PVNRPIDW	HBV POL 615	
1298.07	AAHWILRGTSFYVP	HBV POL 764	
857.02	PHHTALRQAILCWGELMTLA	HBV CORE 50	
35.0100	LQCVFADATPTWGGL	HBV POL 683	
35.0096	ESRLVVDIFSQRGN	HBV POL 387	
35.0093	VGPLTYNEKRRLKLI	HBV POL 96	
1186.18	NLSWLSLDVSAAFYH	HBV POL 422	

Table 2  
HIV derived CTL epitopes

Supertype	Pepitide	Sequence	Source	SEQ ID NO.
A2	924.07	HLSDHFRSV	HIV env 1827	
	1013.002	WLSLIYPRV	HIV-1 <sub>CRF01_AG</sub> (p335.343)	
	777.01	HLIIRLIIT	HIV ENV 183	
	927.15	ALMPLVACI	HIV syw pol 642	
	1168.02	GLSRVVARL	HIV POL 455	
A3	927.11	FLSLGIL	HIV pol 562	
	1147.16	HTLWKAGILYK	HIV POL 149	
	1083.01	STLPETTVRR	HIV env 141	
	1090.11	SACSVVRR	HIV pol 531	
	1090.10	QAFTSTPYK	HIV pol 665	
	1069.16	INVSIPWTK	HIV pol 47	
	1069.20	LVDVFSQSR	HIV pol 388	
	1142.05	KVGNFTGLY	HIV adr POL 629	
	1083.02	TLWKAGILYK	HIV pol 150	
	1145.04	ILPSRISV	HIV ENV 313	
B7	988.05	LPDSRVSF	HIV env 1527	
A2	1147.04	TPARVTGGVF	HIV POL 354	
	1069.06	LLVPFQWVF	HIV env 338-341	
	1147.13	FLAQFTSAI	HIV POL 513	
	1147.14	VLLDYQGMPLV	HIV ENV 259	
	1132.01	LVPPVQWVF	HIV ENV 339	
	1069.05	LLAQFTSAI	HIV pol 504-512	
	927.42	NI.SWLSI.DV	HIV pol 411	
	927.41	LISSNI.SWL	HIV pol 992	
	927.46	KLIILYSIPI	HIV pol 489	
	1069.071	FLAQFTSA	HIV pol 503	
	1142.07	GLIGWSIQA	HIV ENV 62	
	927.47	HLYSIPHL	HIV syw pol 1076	
	1142.08	VLLPFCL	HIV env 377-385	
	1013.1402	VLLPFCL	HIV-1 <sub>CRF01_AG</sub> (p335.343)	
	1090.14	YMDDYVIGIA	HIV POL 340	
A3	26.0339	PLVDFQFSR	HIV pol 376	
	26.0335	GVWIRTPPYAR	HIV X <sub>1</sub> mac lat 299	

Table 2 (Cont'd)  
HIV derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO.
A3	26.0153	SSAGFGALR	HIV X 64	
	1.0993	KVPLGGCR	HIV ad "X" 1548	
	26.0149	CALRTSAR	HIV X 69	
	26.0023	VSRGVWIR	HIV x nuc fus 296	
	26.0545	TLPEITVVRKR	HIV x nuc fus 318	
	20.0131	SVRRRAFTII	HIV POL 524	
	1.0219	FVIGGGRHK	HIV ad "X" 1550	
	26.0008	FTSPITYK	HIV pol 656	
	20.0130	AFSPITYK	HIV POL 655	
	1147.05	PHCLAFSYM	HIV POL 530	
B7	1147.08	YPALMPLYA	HIV POL 640	
	1147.06	LPVCAISSA	HIV X 58	
	1147.02	HPAAMPHLL	HIV POL 429	
	26.0570	YALMAYACI	HIV POL 640	
	1145.08	YHCLMPLY	HIV POL 541	
	1145.08	PHCLALSY	HIV POL 541	
	1090.02	AYRPNAAH	HIV NUC 131	
	1.0519	DLIDTASALY	HIV ad CORE 419	
	13.0129	EVLVSRGVWI	HIV NUC 117	
	20.0254	FAAPTQCCY	HIV POL 631	
Other	2.0060	GYPALMPLY	HIV ALL 1224	
	1069.04	ITLWKAGILY	HIV pol 149	
	1.0166	ILLCLIFLL	HIV env 249-258	
	1.0166	KYGNFTGLY	HEV ad POL 629	
	1069.23	KYTSPPWLL	HIV POL 745	
	1069.01	LLDTASALY	HIV core 59	
	2.0239	LSIDVASALY	HIV ALL 1000	
	2.0181	LYSHIHGR	HIV POL 492	
	1039.01	MAWYWGFSLY	HIV 360	
	2.0126	MSLLENALY	HIV ad 1521	
Other	1069.05	ITLWKAGILY	HIV pol 124	
	1069.09	ITLWKAGILY	HIV pol 808	
	20.0135	PTWIKVGNF	HIV POL 51	
	20.0135	RWMLIRFI	HIV ENV 216	
	20.0269	RWMLIRFI	HIV ENV 216	
	20.0269	SFCGSPYSW	HIV POL 167	
	20.0139	SFCGSPYSW	HIV POL 167	

Table 2 (Cont'd)  
HBV derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO.
Other	1069.02	SLDVSAAY	HBV pol 427	
	20.0136	SWLSLVPH	HBV ENV 134	
	20.0276	SWWTSNRL	HBV POL 392	
	20.0137	SWWTSNRL	HBV ENV 197	
	2.0173	SYQHRKLLI	HBV POL	
	13.0073	WFHSLCLTF	HBV NUC 102	
	1.0774	WLWGMDDPY	HBV analog CORE 416	
	1039.06	WMWVYWGPSLY	HBV env 350	
	924.14	FLPSDIFPSI	HBV 18-27 1a var.	
	1090.77	YMDDVVLGV	HBV pol 538-546 sub	
	941.01	FLJSDYFISV	11b18-27 analog	
	1083.02	STLPETVYVRR	HBV core1 41-151 analog	
	1145.05	FPFSSWAF	HBV ENV 313 analog	
	1145.11	FPICLAFSL	HBV POL 541 analog	
	1145.24	FPICLAFAL	HBV POL 541 analog	
	1145.06	IPITSSWAF	HBV ENV 313 analog	
	1145.25	IPFSSWAF	HBV ENV 313 analog	
	1145.07	FPFCLASSY	HBV POL 441 analog	
	1145.09	FPICLAFSY	HBV ENV 313 analog	
	1145.10	FPICLAFAY	HBV POL 541 analog	

Table 3  
HCV derived HTL epitopes

Peptide	Sequence	Source	SEQ ID NO:
P98.03	AAVAAQGYKVLVINSVAATLGRGAY	HCV NS3 1242-1267	
P98.04	AAVAAQGYKVLVINSVAAT	HCV NS3 1242	
P98.05	GKVLVINSVAATLGRGAY	HCV NS3 1248	
1283.21	GKVLVINSVAAT	HCV NS3 1248	
1283.20	AGQYKVLVINSVAA	HCV NS3 1253	
F134.08	GEGAVQWMNRLIAFASRGNIVS	HCV NS4 1914-1935	
1283.44	GEGAVQWMNRLIAFASRGNIV	HCV NS4 1914	
1283.46	MNRLIAFASRGNIVS	HCV NS4 1921	
1283.55	SKGWILLAPITAYAQ	HCV NS3 1025	
F134.05	GSSTGRQYSPGRQVE	HCV NS5 2641	
1283.61	NPAQGLGGLGGLPLGNPA	HCV NS4 1772	
1283.25	ASCLRLGVRLPLGNPA	HCV NS5 2939	
35.0107	GHLIFCHSKKKCFE	HCV NS3 1393	
	TVDFSLDPTFTFTT	HCV NS4 1436	
35.0106	VVVVATDALMTGTG	HCV NS3 1437	

Table 4  
HCV derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO.
A2	1090.18	ELLADARY	HCV NS3/NS5 728	
	1073.05	LLPILGGWV		
	1013.02	YLVAVQATV		
	1013.1002	DLMGVHLV		
	1090.22	RLVFPDLGV		
	24.0075	VLVGGVLA		
	24.0073	WMNRLLAFA		
	1174.08	HMWNFISGI		
	1073.06	ILAGYGAGV		
	24.0071	ELLFLLADA		
	1001.07	YLPRGGPRL		
	1010.19	ELYTRHADV		
A3	1093.2	YSERSQPR	HCV NS3 1136	
	1073.10	GVVGGVFAK		
	1012.3	LIQHSFAK		
	1095.5	QLFTSPRR		
	1073.11	RLGVRAIRK		
	1073.13	RMVVGVEHRR		
	24.0090	VAGALVAFK		
	F104.01	VGITLLPNR		
	1145.12	LPGCSFISF		
	29.0035	IPPTGKAI		
Other	1009.62	CTASNDLY	HCV Core 168	
	24.0092	FWAKIMWNF	HCV NS3 1128	
	13.0019	LSAIFSLISY	HCV NS4 1765	
			HCV NS5 2922	

Table 4 (Cont'd)  
HCV derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO:
A3	24.0086	LFQGYMSK	HCV NS3 1267	
	1174.21	RVGERMALY	HCV NS5 2621	
	1174.6	WRNSGTFR	HCV NS1/2 357	
	1073.04	TLIGITILLI	HCV NS3 1622	
	16.0013	PLNSVYQA	HCV NS3 1388	
B7	15.0047	NYGVNNTY	HCV NS3 1388	
		NYGVNNTY	HCV NS1/2 623	
Other	24.0093	EVGGVRLIIV	HCV NS3 2129	
	3.0417	LTCGFADLMY	HCV 126	
	1073.01	NVDVQVLY	HCV EL 80	
	1.0509	GLSASLSHY	HCV NS5 2931	
	1073.17	MYVGBLCGSVF	HCV EL 275	
	1073.18	MYVGGVGHRL	HCV NS1/2 633	
	13.075	QYLAGLSTL	HCV NS4 1778	
	1145.13	IFGCSFSIF	HCV Core 1 68	
	1145.25	LPGCMFSIF	HCV Core 1 68	
	1292.24	LPGCSFSI	HCV Core 1 69	
	1145.14	LPGCSFSIF	HCV Core 1 68	
	1145.15	LPGCSFSYF	HCV Core 1 68	

Table 5  
HIV derived HTL epitopes

Peptide	Sequence	Source	SEQ ID NO:
	GEIYKRWILLGLNKIVRMYSPTSLD	IIIV1 GAG 294-319	
	KRWILLGLNKIVRMYSPTSLD	IIIV gag 298-319	
27.0313	KRWILLGLNKIVRMYSPTSLD	IIIV1 GAG 298	
27.0311	GEIYKRWILLGLNKI	IIIV1 GAG 294	
27.0354	WEPYNTPLVLKLYQ	IIIV1 POL 596	
27.0377	QKQTKQNERVYVR	IIIV1 POL 956	
	EKYLLAWYPAHKGIGG	IIIV1 POL 711-726	
1280.03	KYLLAWYPAHKGIGG	IIIV POL 712	
27.0361	EKYLLAWYPAHKGIG	IIIV1 POL 711	
	PIVQCCQMVHQASPTSLNA	IIIV1 gag 165-186	
27.0304	QQQWYHQASPTSLN	IIIV1 GAG 171	
27.0297	QILLQWYHQASPTSLN	IIIV1 ENV 729	
27.0305	QILLQWYHQASPTSLN	IIIV1 ENV 735	
F09.15	IKQPNNAWQBYGKAMY	IIIV1 ENV 566	
27.0341	FRKYTAFTISINNE	IIIV1 POL 303	
27.0364	HSNWRAASDINLPP	IIIV1 POL 758	
27.0373	KTAVQMAVEHNFR	IIIV1 POL 915	
	DRYHPVHAGPIAQCMREPRGS	IIIV GAG 245	
	AFSPFVIMFALSSEGAITQDLTML	IIIV gag 195-216	
	AFSPFVIMFALSSEGAITQDL	IIIV gag 205	
200.06	SALSSEGAITQDLTML	IIIV gag 205	
27.0307	SPEVIMFALSSEGA	IIIV gag 197	
	LAQIQGWMTNNPIPVGHYKR	IIIV gag 275	
27.0310	QQQGWMTNNPIPVGHYKR	IIIV gag 276	
35.0135	WKKILQRKIDRLID	IIIV VIF 31	
35.0131	WAKIQGFITFQ	IIIV POL 874	
35.0132	RYNTPDQALN	IIIV POL 674	
35.0133	ARTFYDQALNRTK	IIIV POL 674	
35.0133	GAVVIQDNDIEVTP	IIIV POL 989	



Table 6  
HIV derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO:
A2	25.0148	MASDNLHPV	HIV1 POL 70	
	106932	VLAEMSQV	HIV gag 397	
	1211.04	KLTPLCYTL	HIV ENV 134	
	25.0062	KLVGKLNWA	HIV1 POL 87	
	25.0039	LTFGWCFKL	HIV1 NEF 62	
	941.031	ILKBPVHG	HIV1 pol 476-484	
	25.0035	MTNPPHPV	HIV1 GAG 34	
	25.0057	RILQQLLFI	HIV1 VPR 72	
A3	1.0944	AVFHNEKR	HIV POL 1434	
	1.1056	KQNFVYYR	HIV POL 1474	
	106949	QMAVFLNFK	HIV pol 1432	
	946.012	AIQSSMTK	HIV pol 337	
	1150.14	AVFTHNEKR	HIV pol 909	
	940.03	QVFLRMTYK	HIV nef 75-82	
	25.0175	TTLEASIAK	HIV ENV 81	
	1069.43	TVYGVFVWK	HIV env 49	
	25.0209	VTIKGGOLK	HIV1 POL 65	
	1146.01	FVRFQVPL	HIV nef 84-92	
	29.0060	IPHYCAVA	HIV env 293	
B7	15.0073	IPHSLETV	HIV POL 171	
	29.0056	CPKVSFEPI	HIV env 285	
	29.0107	IPYNIQSQGVV	HIV pol 883	
	25.0151	CTLNIPHSI	HIV1 POL 96	
	25.0143	LTPGWCHKL	HIV1 NEF 62	
	25.0043	YTAFTHSI	HIV1 POL 83	
	25.0055	AIIRLQQL	HIV1 VPR 76	
	25.0049	ALVEICTEM	HIV1 POL 52	
	25.0032	LLQLTYWGI	HIV1 ENV 61	
	25.0050	LYQPTVNI	HIV1 POL 100	
	25.0047	YVSGVWGI	HIV1 POL 65	
	25.0162	KAVGGQGI	HIV1 POL 96	
A2	25.0032	KAMASDNL	HIV1 POL 98	
	1211.09	SLLNATDIIV	HIV ENV 814	

Table 6 (Cont'd)  
HIV derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO:
A2	25.0041	TLNFRSRI	HIV POL 96	
	1.0046	IVNGGTRK	HIV POL 1075	
	25.0064	MVHQASRK	HIV GAG 45	
	1.0062	YLAWPAIHK	HIV POL 127	
	1.0942	MTKILFHRK	HIV POL 185	
	25.0184	QMVHQASRK	HIV GAG 45	
	1069.48	AVFIINERKK	HIV pol 1614	
	1069.44	KLGRWPK	HIV pol 1358	
	1069.42	KVYLAWFALIK	HIV pol 1225	
	1.0024	NTPYFAIKK	HIV pol 752	
	25.0062	RIVELLGR	IIIV1 ENV 53	
	25.0095	TKIGGQQLK	IIIV1 POL 65	
	25.0078	TLFCSDAK	IIIV1 ENV 82	
	25.0104	VMVWQYDR	IIIV1 YIF 83	
B7	1069.57	VTYYGVVWK	HIV env 48	
	1069.58	YPLASRLF	HIV GAG 307	
	1202.1	YPLASRLA	HIV GAG 248	
	19.0044	YPLQRLPA	HIV env REV 71	
Other	1.0431	ENVNTDSQV	HIV GAG 147	
	1.0014	FRDYVDREY	HIV GAG 298	
	25.0113	IVGCSGKLI	IIIV1 ENV 68	
	25.0127	IVETYGDTW	IIIV1 VPR 92	
	1069.60	IVQEFKNL	HIV pol 1036	
	2.0129	IVQYMDOLY	HIV pol 359	
	25.0128	PNYNTVFAI	IIIV1 VPR 56	
	25.0123	RYLKDQQLL	IIIV1 POL 74	
	1069.57	RYLRDQQLL	IIIV env 2778	
	1069.58	TYQIQEPEF	HIV pol 1033	
	1069.59	VTYQYMDOLY	HIV pol 358	
	1069.27	VTYLDVGDAY	HIV pol 265	
	1069.26	YKREATTLL	IIIV1 ENV 47	
	25.0118	YKREATTLL	IIIV1 ENV 47	
	25.0218	YKREATTLL	IIIV1 ENV 47	
	25.0219	YKQATWFEW	IIIV1 POL 96	

Table 6 (Cont'd)  
HIV derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO:
A2 A3	1211.4	SLINATAIV	HIV MN gp160 E14(a)	
	F105.21	AIQSSMTK	HIV pol 337(a)	
	F105.17	AIQSSMTK	HIV pol 337(a)	
	F105.02	AIQSSMTK	HIV pol 337(a)	
	F105.03	AIQSSMTK	HIV pol 337(a)	
	F105.04	AIQSSMTK	HIV pol 337(a)	
	F105.05	AIQSSMTK	HIV pol 337(a)	
	F105.06	AIQSSMTK	HIV pol 337(a)	
	F105.07	AIQSSMTK	HIV pol 337(a)	
	F105.08	AIQSSMTK	HIV pol 337(a)	
	F105.09	AIQSSMTK	HIV pol 337(a)	
	F105.11	AIQSSMTK	HIV pol 337(a)	
	F105.12	AIQSSMTK	HIV pol 337(a)	
	F105.13	AIQSSMTK	HIV pol 337(a)	
B7	1181.03	FPVRLQVPL	HIV nef 84-92 analog	
	1181.04	FPVRLQVPL	HIV nef 84-92(a)	
	1292.14	FPVRLQVPL	HIV nef 84-92(a)	
	1292.09	FPVRLQVPL	HIV nef 84-92(a)	
	1145.02	FPVRLQVPL	HIV nef 84-92 analog	
	1145.22	FPVRLQVPL	HIV nef 84-92 analog	
	1181.04	FPVRLQVPL	HIV nef 84-92(a)	
	1181.01	FPVRLQVPL	HIV nef 84-92(a)	
	1181.02	FPVRLQVPL	HIV nef 84-92(a)	
	1181.05	FPVRLQVPL	HIV nef 84-92(a)	
	1181.06	FPVRLQVPL	HIV nef 84-92(a)	
	1181.07	FPVRLQVPL	HIV nef 84-92(a)	
	1181.08	FPVRLQVPL	HIV nef 84-92(a)	
	1181.09	FPVRLQVPL	HIV nef 84-92(a)	

Table 7  
*P. falciparum* derived HTL epitopes

Peptide	Sequence	Source	SEQ ID NO:
F125.04	RINWVNHAVPLAMKLI	Pf SSP2 61	
1188.34	INWYNHIAVPLAMKLI	Pf SSP2 62	
1188.16	KSKYKATSLVLAGLI	Pf EXP1 71	
	LVNLLIFHNGKIKNSE	Pf LSA1 13	
F125.02	LVNLLIFHNGKIKNS	Pf LSA1 16	
27.0402	LLIFHNGKIKNSE	Pf SSP2 512	
1188.32	GLAYKEVVGGAATPY	Pf CSP 410	
27.0392	SSVTVVYNSIGLIM	Pf SSP2 223	
27.0393	VSYGPHRMKCYE	Pf CSP 2	
27.0387	MRKAGSSLLP	Pf CSP 33	
27.0388	MRNYGQBNWVSLKK	Pf SSP2 84	
1188.38	KYKIAGGIAGELALI	Pf EXP1 82	
1188.13	AGLLGNVSTVLLGV	Pf LSA1 94	
27.0408	QTNFKSLRLNLOVSE	Pf SSP2 165	
35.0171	PDSIQSLKESKRLN	Pf SSP2 211	
35.0172	KCNLYADSAVENVKN		

Table 8  
*P. falciparum* derived CTL epitopes

Supertype	Peptide	Sequence	Source	SEQ ID NO:
A2	1167.21	FLIFDLFLV	Pf SSP2 14	
	1167.08	GLIMVSEL	Pf CSP 425	
	1167.12	VLAGLLGNV	Pf EXP1 80	
	1167.13	KILSVFLA	Pf EXP1 2	
	1167.10	GLLGNVSTV	Pf EXP1 83	
	1167.18	ILSVSSFLV	Pf CSP 7	
	1167.19	VLLGGVGLV	Pf EXP1 91	
A3	1167.36	LACAGLAYK	Pf SSP2 511	
	1167.22	QNFKSLIR	Pf LSA1 94	
	1167.43	VTGNGQVR	Pf CSP 375	
	1167.24	ASLNNK	Pf EXP1 10	
	1167.28	GVSRHEIK	Pf LSA1 165	
	1167.47	HVLSHNVK	Pf LSA1 59	
	1167.51	LLACAGLAYK	Pf SSP2 510	
	1167.46	FILNLIIRI	Pf LSA1 11	
	1101.03	MPLTQLAI	Pf SHED 77	
	1167.61	TYVAGEPAF	Pf SSP2 539	
A2	1167.14	FLIFDLFL	Pf SSP2 14	
	1167.16	PMKAVCEV	Pf SSP2 230	
	1167.15	LIFFDLFLV	Pf SSP2 15	
	1167.17	LLMDCGSI	Pf SSP2 51	
	1167.09	VILGGVGLV	Pf EXP1 91	
B7	19.0051	TPVGHITNI	Pf SSP2 126	
Other	16.0245	PDDEENIGV	Pf LSA1 1794	
	16.0040	FVEALLQEV	Pf CSP 15	
	1167.54	KYFELVRL	Pf LSA1 9	
	1167.55	KYFELVRL	Pf EXP1 73	
	1167.56	KYLVVRL	Pf SSP2 163	
	15.0184	LRSENERGY	Pf LSA1 663	
	16.0130	PSDCKONLY	Pf SSP2 207	
	16.0077	PSENERGYV	Pf LSA1 1664	
	1167.57	TYVAGEPAF	Pf SSP2 528	
	1167.55	YYPHQSSL	Pf LSA1 1671	

Table 9. Activation of T Cell Proliferation by Expression  
 Vectors Encoding MHC Class II Epitopes Fused to MHC  
 Class II Targeting Sequences

	Immunogen	Stimulating Peptide <sup>1</sup>		
		PADRE	OVA 323	CORE 128
	peptide + CFA <sup>2</sup>	3.0 (1.1)	2.7 (1.2)	3.2 (1.4)
10	pEP2.(PAOS).(-)	-	-	-
	pEP2.(AOS).(-)	5.6 (1.8)	-	-
	pEP2.(PAOS).(sigTh)	5.0 (2.9)	-	2.6 (1.5)
	pEP2.(PAOS).(I $\alpha$ Th)	5.6 (2.1)	-	3.0 (1.6)
	pEP2.(PAOS).(LampTh)	3.8 (1.7)	-	3
15	pEP2.(PAOS).(IiTh)	5.2 (2.0)	3.2 (1.5)	3.7 (1.5)
	pEP2.(PAOS).(H2M)	3.3 (1.3)	-	2.8

<sup>1</sup>Geometric mean of cultures with SI  $\geq 2$ .

<sup>2</sup>Proliferative response measured in the lymph node.

Table 10  
CTL Epitopes in cDNA Minigene

Epitope	Sequence	MHC Restrict.	MHC Binding Affinity [IC30% (nM)]	Immunogenicity In Vivo (IFA)	
				No. CTL- Positive Cultures	CTL Response (Geo. Mean x/±SD) <sup>b</sup> ΔLU
HBV Core 18	FLPSDFFPSV	A2.1	3	6 / 6	73.0 (1.1)
HBV Env 335	WLSLLVPFV	A2.1	5	4 / 6	5.3 (1.6)
HBV Pol 455	GLSRVVARL	A2.1	76	ND <sup>c</sup>	ND
HIV Env 120	KLTPLCVTL	A2.1	102	2 / 5	6.4 (1.3)
HIV Pol 476	ILKEPVHGV	A2.1	192	2 / 5	15.2 (2.9)
HBV Pol 551-A	YMDDVVLGA	A2.1	200	0 / 6	-
HBV Pol 551-V	YMDDVVLGV	A2.1	5	6 / 6	8.2 (2.3)
HIV Env 49	TVYYGVPVWK	A11	4	28 / 33	13.4 (3.1)
HBV Core 141	STLPETTVVRR	A11	4	6 / 6	12.1 (2.6)
HBV Pol 149	HTLWKAGILYK	A11	14	6 / 6	13.1 (1.2)

a Peptide tested in HLA-A2.1/K<sup>b</sup> H-2<sup>baw</sup> transgenic mice by co-immunizing with a T helper cell peptide in IFA.

b Geometric mean CTL response of positive cultures.

c ND, not done.

Table 11  
Summary of Immunogenicity of pMin.1 DNA  
construct in HLA A2.1/K<sup>b</sup> transgenic mice

Epitope	CTL Response <sup>a</sup>	
	No. Positive Cultures/Total <sup>b</sup>	Geo. Mean Response Positive Cultures [ $\bar{x}$ /±SD]
		ΔLU
HBV Core 18	9 / 9	455.5 [2.2]
HIV Env 120	12 / 12	211.9 [3.7]
HBV Pol 551-V	9 / 9	126.1 [2.8]
HBV Pol 455	12 / 12	738.6 [1.3]
HIV Pol 476	11 / 11	716.7 [1.5]
HBV Env 335	12 / 12	43.7 [1.8]
HBV Core 18 (Theradigm) <sup>c</sup>	10 / 10	349.3 [1.8]

- <sup>a</sup> Mice were immunized with pMin.1 DNA or Theradigm-HBV lipopeptide and CTL activity in splenocyte cultures was determined after in vitro stimulation with individual peptide epitopes. Results from four independent experiments are shown.
- <sup>b</sup> See Example V, Materials and Methods for definition of a CTL-positive culture.
- <sup>c</sup> Response of mice immunized with Theradigm-HBV lipopeptide containing the HBV Core 18 epitope.



Table 12  
Summary of immunogenicity  
in HLA A11/K<sup>b</sup> transgenic mice

Epitope	CTL Response <sup>a</sup>	
	No. Positive Cultures/Total <sup>b</sup>	Geo. Mean Response Positive Cultures [ $\bar{x}$ $\pm$ SD]
HBV Core 141	5/9	$\Delta$ LU 128.1 [1.6]
HBV Pol 149	6/9	267.1 [2.2]
HIV Env 43	9/9	40.1 [2.9]

5     <sup>a</sup> Mice were immunized with pMin.1 DNA and CTL activity in splenocyte cultures was determined after in vitro stimulation with individual A11-restricted epitopes. The geometric mean CTL response from three independent experiments are shown.

<sup>b</sup> Definition of a CTL-positive culture is described in Example V, Materials and Methods.

## WHAT IS CLAIMED IS:

- 1                   1.     An expression vector comprising a promoter operably linked to a  
2     first nucleotide sequence encoding a major histocompatibility (MHC) targeting sequence  
3     fused to a second nucleotide sequence encoding two or more heterologous peptide  
4     epitopes, wherein the heterologous peptide epitopes comprise two HTL peptide epitopes  
5     or a CTL peptide epitope and a universal HTL peptide epitope.
- 1                   2.     The expression vector of claim 1, wherein the heterologous peptide  
2     epitopes comprise two or more heterologous HTL peptide epitopes.
- 1                   3.     The expression vector of claim 1, wherein the heterologous peptide  
2     epitopes comprise a CTL peptide epitope and a universal HTL peptide epitope.
- 1                   4.     The expression vector of claim 2, wherein the heterologous peptide  
2     epitopes further comprise one or more CTL peptide epitopes.
- 1                   5.     The expression vector of claim 3, wherein the heterologous peptide  
2     epitopes further comprise two or more CTL peptide epitopes.
- 1                   6.     The expression vector of claim 3, wherein the heterologous peptide  
2     epitopes further comprise two or more HTL peptide epitopes.
- 1                   7.     The expression vector of claim 2, wherein one of the HTL peptide  
2     epitopes is a universal HTL epitope.
- 1                   8.     The expression vector of claim 3 or 7, wherein the universal HTL  
2     epitope is a pan DR epitope.
- 1                   9.     The expression vector of claim 8, wherein the pan DR epitope has  
2     the sequence AlaLysPheValAlaAlaTrpThrLeuLysAlaAlaAla (SEQ ID NO:38).
- 1                   10.    The expression vector of claim 1, wherein the peptide epitopes are  
2     hepatitis B virus epitopes, hepatitis C virus epitopes, human immunodeficiency virus  
3     epitopes, human papilloma virus epitopes, MAGE epitopes, PSA epitopes, PSM epitopes,  
4     PAP epitopes, p53 epitopes, CEA epitopes, Hcr2/neu epitopes, or *Plasmodium* epitopes.

- 1                    11.    The expression vector of claim 10, wherein the peptide epitopes  
2 each have a sequence selected from the group consisting of the peptides depicted in  
3 Tables 1-8.
- 1                    12.    The expression vector of claim 11, wherein at least one of the  
2 peptide epitopes is an analog of a peptide depicted in Tables 1-8.
- 1                    13.    The expression vector of claim 1, wherein the MHC targeting  
2 sequence comprises a region of a polypeptide selected from the group consisting of the li  
3 protein, LAMP-I, HLS-DM, HLA-DO, H2-DO, influenza matrix protein, hepatitis B  
4 surface antigen, hepatitis B virus core antigen, Ty particle, Ig- $\alpha$  protein, Ig- $\beta$  protein, and  
5 Ig kappa chain signal sequence.
- 1                    14.    The expression vector of claim 1, wherein the expression vector  
2 further comprises a second promoter sequence operably linked to a third nucleotide  
3 sequence encoding one or more heterologous HTL or CTL peptide epitopes.
- 1                    15.    The expression vector of claim 1, wherein the vector comprises  
2 pMin1 or pEP2.
- 1                    16.    The expression vector of claim 3 or 4, wherein the CTL peptide  
2 epitope comprises a structural motif for an HLA supertype, whereby the peptide CTL  
3 epitope binds to two or more members of the supertype with an affinity of greater than  
4 500 nM.
- 1                    17.    The expression vector of claim 4 or 5, wherein the CTL peptide  
2 epitopes have structural motifs that provide binding affinity for more than one HLA allele  
3 supertype.
- 1                    18.    A method of inducing an immune response *in vivo* comprising  
2 administering to a mammalian subject an expression vector comprising a promoter  
3 operably linked to a first nucleotide sequence encoding a major histocompatibility (MHC)  
4 targeting sequence fused to a second nucleotide sequence encoding two or more  
5 heterologous peptide epitopes, wherein the heterologous peptide epitopes comprise two  
6 HTL peptide epitopes or a CTL peptide epitope and a universal HTL peptide epitope.

- 1                   19.    The method of claim 18, wherein the heterologous peptide epitopes  
2   comprise two or more heterologous HTL peptide epitopes.
- 1                   20.    The method of claim 18, wherein the heterologous peptide epitopes  
2   comprise a CTL peptide epitope and a universal HTL peptide epitope.
- 1                   21.    The method of claim 19, wherein the heterologous peptide epitopes  
2   further comprise one or more CTL peptide epitopes.
- 1                   22.    The method of claim 20, wherein the heterologous peptide epitopes  
2   further comprise two or more CTL peptide epitopes.
- 1                   23.    The method of claim 20, wherein the heterologous peptide epitopes  
2   further comprise two or more HTL peptide epitopes.
- 1                   24.    The method of claim 19, wherein the HTL peptide epitope is a  
2   universal HTL epitope.
- 1                   25.    The method of claim 20 or 24, wherein the universal HTL epitope  
2   is a pan DR epitope.
- 1                   26.    The method of claim 25, wherein the pan DR epitope has the  
2   sequence AlaLysPheValAlaAlaTrpThrLeuLysAlaAlaAla (SEQ ID NO:38).
- 1                   27.    The method of claim 18, wherein the peptide epitopes are hepatitis  
2   B virus epitopes, hepatitis C virus epitopes, human immunodeficiency virus epitopes,  
3   human papilloma virus epitopes, MAGE epitopes, PSA epitopes, PAP epitopes, PSM  
4   epitopes, p53 epitopes, CEA epitopes, Her2/neu epitopes, or *Plasmodium* epitopes.
- 1                   28.    The method of claim 27, wherein the peptide epitopes each have a  
2   sequence selected from the group consisting of the peptides depicted in Tables 1-8.
- 1                   29.    The method of claim 28, wherein least one of the peptide epitopes  
2   is an analog of a peptide depicted in Tables 1-8.
- 1                   30.    The method of claim 18, wherein the MHC targeting sequence  
2   comprises a region of a polypeptide selected from the group consisting of the Ii protein,  
3   LAMP-I, HLS-DM, HLA-DO, H2-DO, influenza matrix protein, hepatitis B surface

4 antigen, hepatitis B virus core antigen, Ty particle, Ig- $\alpha$  protein, Ig- $\beta$  protein, and Ig  
5 kappa chain signal sequence.

1 31. The method of claim 18, wherein the expression vector further  
2 comprises a second promoter sequence operably linked to a third nucleotide sequence  
3 encoding one or more heterologous HTL or CTL peptide epitopes.

1 32. The method of claim 18, wherein the vector comprises pMin.1 or  
2 pEP2.

1 33. The method of claim 20 or 21, wherein the CTL peptide epitope  
2 comprises a structural motif for an HLA supertype, whereby the peptide epitope binds to  
3 two or more members of the supertype with an affinity of greater than 500 nM.

1 34. The method of claim 21 or 22, wherein the CTL peptide epitopes  
2 have structural motifs that provide binding affinity for more than one HLA allele  
3 supertype.

1 35. A method of inducing an immune response *in vivo* comprising  
2 administering to a mammalian subject an expression vector comprising a promoter  
3 operably linked to a first nucleotide sequence encoding a major histocompatibility (MHC)  
4 targeting sequence fused to a second nucleotide sequence encoding a heterologous human  
5 HTL peptide epitope.

1 36. The method of claim 35, wherein the second nucleotide sequence  
2 further comprises two or more heterologous HTL peptide epitopes.

1 37. The method of claim 35, wherein the second nucleotide sequence  
2 further comprises one or more heterologous CTL peptide epitopes.

1 38. The method of claim 35, wherein the HTL peptide epitope is a  
2 universal HTL peptide epitope

1 39. The method of claim 38, wherein the universal HTL epitope is a  
2 pan DR epitope.

1 40. The method of claim 39, wherein the pan DR epitope has the  
2 sequence AlaLysPheValAlaAlaTrpThrLeuLysAlaAlaAla (SEQ ID NO:38).

- 1                   41.    The method of claim 37, wherein the HTL and CTL peptide  
2    epitopes are hepatitis B virus epitopes, hepatitis C virus epitopes, human  
3    immunodeficiency virus epitopes, human papilloma virus epitopes, MAGE epitopes, PSA  
4    epitopes, PAP epitopes, PSM epitopes, p53 epitopes, CEA epitopes, Her2/neu epitopes,  
5    or *Plasmodium* epitopes.
- 1                   42.    The method of claim 41, wherein the peptide epitopes each have a  
2    sequence selected from the group consisting of the peptides depicted in Tables 1-8.
- 1                   43.    The method of claim 42, wherein at least one of the peptide  
2    epitopes is an analog of a peptide depicted in Tables 1-8.
- 1                   44.    The method of claim 35, wherein the MHC targeting sequence  
2    comprises a region of a polypeptide selected from the group consisting of the  $\Pi$  protein,  
3    LAMP-I, HLS-DM, HLA-DO, H2-DO, influenza matrix protein, hepatitis B surface  
4    antigen, hepatitis B virus core antigen, Ty particle, Ig- $\alpha$  protein, Ig- $\beta$  protein, and Ig  
5    kappa chain signal sequence.
- 1                   45.    The method of claim 35, wherein the expression vector further  
2    comprises a second promoter sequence operably linked to a third nucleotide sequence  
3    encoding one or more heterologous HTL or CTL peptide epitopes.
- 1                   46.    The method of claim 37, wherein the CTL peptide epitope  
2    comprises a structural motif for an HLA supertype, whereby the peptide epitope binds to  
3    two or more members of the supertype with an affinity of greater than 500 nM.
- 1                   47.    The method of claim 37, wherein the CTL peptide epitopes have  
2    structural motifs that provide binding affinity for more than one HLA allele supertype.
- 1                   48.    A method of assaying the human immunogenicity of a human T  
2    cell peptide epitope *in vivo* in a non-human mammal, comprising the step of  
3    administering to the non-human mammal an expression vector comprising a promoter  
4    operably linked to a first nucleotide sequence encoding a heterologous human CTL or  
5    HTL peptide epitope.

- 1                   49.   The method of claim 48, wherein the first nucleotide sequence  
2   encodes two or more heterologous CTL or HTL peptide epitopes.
- 1                   50.   The method of claim 48, wherein the non-human mammal is a  
2   transgenic mouse that expresses a human HLA allele.
- 1                   51.   The method of claim 50, wherein the human HLA allele is selected  
2   from the group consisting of A11 and A2.1.
- 1                   52.   The method of claim 48, wherein the expression vector further  
2   comprise a second nucleotide sequence encoding a major histocompatibility (MHC)  
3   targeting sequence.
- 1                   53.   The method of claim 48, wherein the HTL peptide epitope is a  
2   universal HTL epitope.
- 1                   54.   The method of claim 53, wherein the universal HTL epitope is a  
2   pan DR epitope.
- 1                   55.   The method of claim 54, wherein the pan DR epitope has the  
2   sequence AlaLysPheValAlaAlaTrpThrLeuLysAlaAlaAla (SEQ ID NO:38).
- 1                   56.   The method of claim 48, wherein the CTL or HTL peptide epitopes  
2   are hepatitis B virus epitopes, hepatitis C virus epitopes, human immunodeficiency virus  
3   epitopes, human papilloma virus epitopes, MAGE epitopes, PSA epitopes, PSM epitopes,  
4   PAP epitopes, p53 epitopes, CEA epitopes, Her2/neu epitopes, or *Plasmodium* epitopes.
- 1                   57.   The method of claim 56, wherein the CTL or HTL peptide epitopes  
2   each have a sequence selected from the group consisting of the peptides depicted in  
3   Tables 1-8.
- 1                   58.   The method of claim 57, wherein at least one of the peptide  
2   epitopes is an analog of a peptide depicted in Tables 1-8.
- 1                   59.   The method of claim 52, wherein the MHC targeting sequence  
2   comprises a region of a polypeptide selected from the group consisting of the Ii protein,

3 LAMP-I, HLS-DM, HLA-DO, H2-DO, influenza, hepatitis B virus core antigen, Ty  
4 particle, Ig- $\alpha$  protein, Ig- $\beta$  protein, and Ig kappa chain signal sequence.

1           60.    The method of claim 48, wherein the expression vector further  
2 comprises a second promoter sequence operably linked to a third nucleotide sequence  
3 encoding one or more heterologous human CTL or HTL peptide epitopes.

1           61.    The method of claim 48, wherein the vector comprises pMin.1 or  
2 pEP2.

1           62.    The method of claim 48, wherein the CTL peptide epitope has a  
2 structural motif that provides binding affinity for an HLA allele supertype.

1           63.    The method of claim 49, wherein the CTL peptide epitopes have  
2 structural motifs that provide binding affinity for more than one HLA allele supertype.

1           64.    The method of claim 48, wherein the expression vector comprises  
2 both HTL and CTL peptide epitopes.



```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
GCTAGCCGCGCCACCATGGATGACCAACGCGACCTCATCTTAACCATGAGCAATGCCCATACTGGGCA
CGATCCGCGCGGTGGTACTACTCTGGTGGCGCTGGAGTAGAGATTGGTACTCGTTAACGGGTATGACCCCGT
      M D D Q R D L I S N H E Q L P I L G >

      80      90      100     110     120     130     140
      *      *      *      *      *      *      *
ACCGCCCTAGAGAGCCAGAAAGGTGCAGCGGTGGAGCTCTGTACACCGGTGTTCTGTCTGGTGGCTCT
TGGCGGGATCTCTGGTCTTTCCACGTCCGCACTCGAGACATGTGGCCACAAAGACAGGACCCACGAGA
N R P R E P E R C S R G A L Y T G V S V L V A L>

      150     160     170     180     190     200     210
      *      *      *      *      *      *      *
GCTCTGGCTGGGCGAGGCCACCACTGCTTACTTCCGTGTAACAGCAACAGGGCCGCTAGACAAGCTGACC
CGAGAACCGACCCGTCGGTGGTACGAATGAAGGACATGGTCGTTGTCGGCGGGATCTGTTGACTGG
      L L A G Q A T T A Y F L Y Q Q Q G R L D K L T>

      220     230     240     250     260     270     280
      *      *      *      *      *      *      *
ATCACCTCCAGAACCTGCAACTGGAGAGCCTTCGCATGAAGCTTCGAAATCTGCCAACCTGTGGCCA
TAGTAGAGGTCTTGGACCTTGACCTCTCGGAAGCGTACTCGAAGCGTTAGACGGTTTGGACACCGGT
      I T S Q N L Q L E S L R M K L P K S A K P V A>

      290     300     310     320     330     340     350
      *      *      *      *      *      *      *
AGTCTGTGGCTGCTCGGACCTGAAGGCTGCCGCTATGTCCATGGAACATGCTCCTTGGGCTGTGAA
TCAAGCACCGACGACCTGGGACTTCGACGCGGATACAGGTACTATTGTACGAGGAACCCGAGCACTCT
      K F V A A W T L K A A A M S M D N M L L G P V K>

      360     370     380     390     400     410     420
      *      *      *      *      *      *      *
GAACGTTACCAAGTACGSCAACATGACCCAGGACCATGTGATGCATCTGCTCACGAGGTCTGGACCCCTG
CTTGCAATGGTTTCATGCGGTGTACTGGGTCTGTGACACTAGTAGACGAGTGTCCAGACCTGGGGAC
      N V T K Y G N M T Q D H V M H L L T R S G F L>

      430     440     450     460     470     480     490
      *      *      *      *      *      *      *
GAGTACCGCGAGCTGAAGGGGACCTTCCAGAGAATCTGAGCATCTTAAAGAACTCCATGATGGCGGTA
CTCATGGGCGTGCACCTCCCTGGAAAGGCTCTTTAGACTTCGTAGAATTCTTGAGGTACCTACCGACT
      E Y P Q L K G T F P E N L K H L K N S M D G V>

      500     510     520     530     540     550     560
      *      *      *      *      *      *      *
ACTGGAAGATCTCGAGAGCTGGATGAAGCAGTGCTCTGTTGAGATGAGCAAGAACTCCCTGGAGGA
TGAATCTTAGAAGCTCTCGACCTACTCTGTCAACCGAACAACACTACTGCTTCTTGAGGGACCTCCT
      N W X I F E S W M M K Q W L L F E M S K N S L E E>

      570     580     590     600     610     620     630
      *      *      *      *      *      *      *
GAAGAGCCACCGAGGCTCCACTAAAGAGCCACTGACATGGAAGACTTCTTCTGGCTGGGAGTG
CTTCTGGGTGGCTCCGAGGTGGATTCTCGTGACCTGTACTCTCGATAGAGACCGGACCTCTCAC
      K K P T E A P P K E P L D M E D L S S G L G V>

      640     650     660
      *      *      *
ACCAGGAGGAACCTGGGTCAAGTCAACCTGTGAGGTACC
TGTGCTGCTTGAACCACTGAGTGGGACACTCCATGG
      T R Q E L G Q V T L >>

```

FIGURE 1

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
GCTAGCGCCGCCACCATGGATGACCAACGCGACCTCATCTCTAACCATGAGCAATTGCCCATCTGGGCA
CGATCGCGCGCGGTGTACTCTACTGGTTGCGCTGGAGTAGAGATTGGTACTCGTTAACGGGTATGACCGGT
      M D D Q R D L I S N H E Q L P I L G>

      80      90      100     110     120     130     140
      *      *      *      *      *      *      *
ACCGCCCTAGAGAGCCAGAAAGGTGCAGCGGTGGAGCTCTGTACACCGGTGTTTCTGTCTGGTGGCTCT
TGCGCGGATCTCTCGTCTTTCCAGCTCGGCACCTCGAGACATGTGGCCACAAAGACAGGACCACCGAGA
N R P R E P E R C S R G A L Y T G V S V L V A L>

      150     160     170     180     190     200     210
      *      *      *      *      *      *      *
GCTCTGGGCTGGGCGAGGCCACCACTGCTTACTTCTCTGTACAGCAACAGGGCCGCTAGACAAGCTGACC
CGAGAACCGACCCGTCGCGTGGTGACGAATGAAGGACATGGTCGTTGTCCCGCGCATCTGTCTCGACTGG
      L L A G Q A T T A Y F L Y Q Q G R L D K L T>

      220     230     240     250     260     270     280
      *      *      *      *      *      *      *
ATCACCTCCCAAGACCTGCAACTGGAGAGCCTTCGCATGAAGCTTATCAGCCAGGCTGTGCAGCGCGCTC
TAGTGAGGGGTCTTGGACCTTGACCTCTCGGAAGCGTACTTCGAATAGTCGGTCCGACAGCTGGCGGAG
      I T S Q N L Q L E S L R M K L I S Q A V H A A>

      290     300     310     320     330     340     350
      *      *      *      *      *      *      *
ACGCGGAATCAACGAAGCTGGAAGAACCCCTCCAGCTTATCGCCCTCCAAACGCTCCTATCTGTTCTT
TGCGCGCTTAGTTGCTTCGACCTTCTTGGGAGGTGCAATAGCGGGAGGTTTGGAGGATAGGACAAGAA
H A E I N E A G R T P P A Y R P P N A P I L F F>

      360     370     380     390     400     410     420
      *      *      *      *      *      *      *
TCTGTGACCAGAATCCTGACAATCCCCAGTCCCTGGACGCCAAGTTCTGGTGTGCTGGACCTGAAG
AGAAGACTGGTCTTAGGACTGTTAGGGGGTCAGGGACCTCGGTTCAAGCACCGACCGGACCTGGGACTTC
      L L T R I L T I P Q S L D A K F V A A W T L K>

      430
      *      *      *
GCTGCGCTTGAGGTACC
CGACGCGGAACCTCATGG
      A A A *>

```

FIGURE 2

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
      GCTAGCGCGGCCACCATGGATGACCAACGCGACCTCATCTCTAACCATGAGCAATTGCCATCTGGGCA
      CGATCGCGCGGTGGTCTACTCTGTTGCGCTGGAGTAGAGATTGGTACTCGTTAACGGGTATGACCGGT
      M D D Q R D L I S N H E Q L P I L G>

      80      90      100     110     120     130     140
      *      *      *      *      *      *      *
      ACCGCGCTAGAGAGCCCGAAAGGTGCAGCGGTGGAGCTCTGTACACGGTGTCTTCTGTCTGGTGGCTCT
      TGGCGGGATCTCTCGGTCTTTCCACGTCCGCACTCGAGACATGTGGCCACAAAGACAGGACCACCGAGA
      N R P R E P E R C S R G A L Y T G V S V L V A L>

      150     160     170     180     190     200     210
      *      *      *      *      *      *      *
      GCTCTTGGCTGGGCGAGCCACCACTGCTTACTTCTGTACCAAGCAAGGGCGCTAGACAAGCTGACC
      CGAGAAGCAAGCGCTCGGTGGTGCAGAAAGGACATGGTGTGTCGGCGGGATCTGTTCGACTGG
      L L A G Q A T T A Y F L Y Q Q G R L D K L T>

      220     230     240     250     260     270     280
      *      *      *      *      *      *      *
      ATCACCTCCAGAACCTGCAACTGGAGAGCTTGGCATGAAGCTTATCAGCCAGGCTGTGCACGCGGCTC
      TAGTGGAGGCTCTTGGAGCGTTGACCTCTCGGAAGCGTACTCGAATAGTCGGTCCGACACCTGCGGCGAG
      I T S Q N L Q L E S L R M K L I S Q A V H A A>

      290     300     310     320     330     340     350
      *      *      *      *      *      *      *
      ACGCCGAATCAACGAAGCTGGAAGAAGCCCTCCAGCTTATCGCCCTCCAAACGCTCCTATCTGTCTTT
      TGGCGCTTAGTTGCTTCGACCTTTCTGGGAGGTGCAATAGCGGGAGGTTTGCAGGATAGGACAAGAA
      H A E I N E A G R T P P A Y R P P N A P I L F F>

      360     370     380     390     400     410     420
      *      *      *      *      *      *      *
      TCTGCTGACCAAGATCCTGACAACTCCCGAGTCCCTGGAGCCCAAGTTCGTGGCTGCTGGACCTCGAAG
      AGACGACTGGTCTTAGGACTGTTAGGGGCTCAGGAGCTCGGGTTCAGACACCGACGGACCTGGGACTTC
      L L T R I L T T P Q S L D A K F V A A W T L K>

      430     440     450     460     470     480     490
      *      *      *      *      *      *      *
      GCTGCGCTATGTCCATGGATAACATGCTCCTTGGGCGCTGTGAAGAAGCTTACCAAGTACGGCAACATGA
      CAGCGCGATACAGGTGACTTATGTACGAGGAAGCCCGACACTCTTCGAATGGTTCATGCGGTGTGTACT
      A A A M S M D N M L L G P V K N V T K Y G N M>

      500     510     520     530     540     550     560
      *      *      *      *      *      *      *
      CCCAGGACATGTGATGCATGCTCACGAGGTCTGGACCCCTGGAGTACCCGACGCTGAAGGGGACCTT
      GGTCTCTGTACACTACGTAGACAGGTGCTCCAGACCTCGGGACCTCATGGGCTGCGACTTCCCTCGAA
      T Q D H V M H L L T R S G P L E Y P Q L K G T F>

      570     580     590     600     610     620     630
      *      *      *      *      *      *      *
      CCCAGAGATCTGAAGCATCTTAAGAACTCCATGGATGGCGTGAAGTGAAGATCTTCGAGAGCTGGATG
      GGTCTCTTAGACTTCGTAGAAATCTTGAAGGTACCTACCGCACTTGACCTTCTAGAAGCTCTCGACCTAC
      P E N L K H L K N S M D G V N W K I F E S W M>

```

FIGURE 3

```

      640      650      660      670      680      690      700
      *      *      *      *      *      *      *
AAGCAGTGGCTCTTGTTTGAGATGAGCAAGAACTCCCTGGAGGAGAAGAACCCACCGAGGCTCCACCTA
TTCGTCAACCGAGAACAACTCTACTCGTTCTTGAGGGACCTCCTCTTCTTCGGGTGGCTCCGAGGTGGAT
K Q W L L F E M S R N S L E E K K P T E A P P>

      710      720      730      740      750      760      770
      *      *      *      *      *      *      *
AAGAGCCACTGGACATGGAAGACCTATCTTCTGGCTGGGAGTGACCAGGCAGGAAGTGGGTCAAGTCAC
TTCCTCGTGACCTGTACCTTCTGGATAGAAGACCGGACCTCACTGGTCCGTCTTGACCCAGTTTCAGTG
K E P L D M E D L S S G L G V T R Q E L G Q V T>

      780
      *
CCTGTGAGGTACC
GGACACTCCATGG
L *>

```

FIGURE 3 CONTINUED

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
GCTAGCGCCGCCACCATGGGAATGCAGGTGCAGATCCAGAGCCTGTTTCTGCTCCTCTGTTGGGTGCCCG
CGATCGCGCGGTGGTACCTTTACGTCCAGTCTAGGTCTCGGACAAAGACGAGGAGGACACCCACGSGGC
      M G M Q V Q I Q S L F L L L L W V P>

      80      90      100     110     120     130     140
      *      *      *      *      *      *      *
GGTCCAGAGGAATCAGCCAGGCTGTGCACGCCGCTCACGCCGAATCAACGAAGCTGGAAGAACCCTCC
CCAGGTCTCCTTAGTTCGGTCCGACACGTGCGCGGAGTGCGGCTTTAGTTGCTTCGACCTTCTTGGGAGG
G S R G I S Q A V H A A H A E I N E A G R T P>

      150     160     170     180     190     200     210
      *      *      *      *      *      *      *
AGCTTATCGCCTCCAAACGCTCCTATCCTGTTCTTTCTGCTGACCAGAATCCTGACAATCCCCAGTCC
TCGAATAGCGGGAGGTTTGGAGGATAGGACAAGAAAGACACTGGTCTTAGGACTGTTAGGGGTCAGG
  A Y R P P N A P I L F F L L T R I L T I P O S>

      220     230     240     250     260     270     280
      *      *      *      *      *      *      *
CTGGACGCCAAGTTCGTGGCTGCCTGGACCTGAAGGCTGCCGCTAACACATGTTGATCCCCATTGCTG
GACTTCGGGTTCAAGCACCGACGGACCTGGGACTTCGAGGGGATTTGTACAACCTAGGGGTAACGAC
  L D A K F V A A W T L K A A A N N M L I P I A>

      290     300     310     320     330     340     350
      *      *      *      *      *      *      *
TGGGCGGTGCCCTGGCAGGGCTGGTCTCATCGTCTCATTCGCTACCTCATTGGCAGGAAGAGGAGTCA
ACCCGCCAGGGACCGTCCCGACGAGGAGTAGCAGGAGTAACGGATGGAGTAACCGTCTCTTCTCCTCAGT
V G G A L A G L V L I V L I A Y L I G R K R S H>

      360     370
      *      *
CGCGGCTATCAGACCATCTAGGGTACC
GCGGCGGATAGTCTGAGTAGATCCCATGG
  A G Y Q T I >>

```

FIGURE 6

```

      10      20      30      40      50      60      70
*      *      *      *      *      *      *
GCTAGCGCGCCACCATGGCTGCACTCTGGCTGCTGCTGGTCTCTCAGTCTGCACTGTATGGGGATCA
CGATCGCGCGGTGGTACCGACGTGAGACCGACGACGACACGAGGTGAGACGTGACATACCCCTAGT
      M A A L W L L L L V L S L H C M G I>

      80      90      100      110      120      130      140
*      *      *      *      *      *      *
GCCAGGCTGTGCACGCGCTCAGCCGAAATCAACGAAGCTGGAAGAACCCTCCAGCTTATCGCCCTCC
CGGTCCGACACGTGCGGCGAGTGCGGCTTTAGTTGCTTCGACCTTCTTGGGGAGGTGGAATAGCGGGAGG
S Q A V H A A H A E I N E A G R T P P A Y R P P>

      150      160      170      180      190      200      210
*      *      *      *      *      *      *
AAACGCTCCTATCCTGTTCTTTCTGCTGACCAATCTGACAAATCCCCAGTCCCTGGACGCCAAGTTC
TTTGCGAGGATAGGACAAGAAAGACGACTGGTCTTAGGACTGTTAGGGGGTCAAGGACCTGCGGTTCAAG
N A P I L F F L L T R I L T I P Q S L D A K P>

      220      230      240      250      260      270      280
*      *      *      *      *      *      *
GTGGCTGCCTGGACCTGAAGGCTGCCGCTAAGTCTCTGTGCTGCAACCTGGGCTGGGCTTCA
CACCGACGGACCTGGGACTTCCGACGGCGATTCCAGAGACACAGACGTGGTGGGACCCGACCCGAAAT
V A A W T L K A A A K V S V S A A T L G L G F>

      290      300      310      320      330      340      350
*      *      *      *      *      *      *
TCATCTTCTGTGTGGCTTCTTCAGATGGCGCAAGTCTCATTCTCTCAGCTACACTCCTCTCCCTGGATC
AGTGAAGACACACACCAAGAAAGTCTACCGCGTTACAGAGTAAGGAGGTGATGTGAGGAGAGGGACCTAG
I I F C V G F F R W R K S H S S S Y T P L P G S>

      360      370      380
*      *      *
CACCTACCCAGAAGGACGGCATTAGGGTACC
GTGGATGGGCTCTTCTGCCGTAATCCCATGG
      T Y P E G R H >

```

FIGURE 5

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
GCTAGCGCCGCCACCATGGGCGCTGGGAGGGGCCCTCGGTGGTGGCTCTGTTGGTGAACCTCATGAGGC
CGATCGCGCGGTGGTACCCGCGACCTCCCGGGGACCCACCACCGAGACAACCACTTGGAGTACTCCG
      M G A G R A P W V V A L L V N L M R>

      80      90      100     110     120     130     140
      *      *      *      *      *      *      *
TGGATTCCATCAGCCAGGCTGTGCACGCCGCTCACGCCGAATCAACGAAGCTGGAAGAACCCCTCCAGC
ACCTAAGGTAGTTCGGTCCGACACGTGCGGCGAGTGGCGCTTAGTTGCTTCGACCTTCTTGGGAGGTGCG
L D S I S Q A V H A A H A E I N E A G R T P P A>

      150     160     170     180     190     200     210
      *      *      *      *      *      *      *
TTATCGCCCTCCAAACGCTCCTATCCTGTTCTTCTGCTGACCAGAATCCTGACAAATCCCCAGTCCCTG
AATAGCGGGAGGTTTGGAGGATAGGACAGAAGACGACTGGTCTTAGGACTGTTAGGGGGTCAGGGAC
Y R F P N A P I L F F L L T R I L T I P Q S L>

      220     230     240     250     260     270     280
      *      *      *      *      *      *      *
GACGCCAAGTTCGTGGCTSCCTGGACCCCTGAAGGCTGCCGCTATACTGAGTGGAGCTGCAGTGTTCCTGC
CTGCGGTTCAAGCAGCAGCAGGACCTGGGACTTCGACGCGGATATGACTCACCTCGACGTCACAAGGACG
D A K F V A A W T L K A A A I L S G A A V F L>

      290     300     310     320     330     340     350
      *      *      *      *      *      *      *
TTGGGCTGATTGTCTTCCTGGTGGGGTTGTATCCATCTCAAGGCTCAGAAAGCATCTGTGGAGACTCA
AACCCTGACTAACAGAGGAGGACCCCCCAACATAGGTAGAGTTCCGAGTCTTTCGTAGACACCTCTGAGT
L G L I V F L V G V I H L K A Q K A S V E T Q>

      360     370     380     390     400     410     420
      *      *      *      *      *      *      *
GCTTGGCAATGAGAGTAGTCCCGGATGATGGAGCGGCTAACCAAGTTCAAGGCTGGACCGGGACATGTC
CGGACCGTTACTCTCATCCAGGGCTACTACCTCGCGGATGGTTCAAGTTCGACCTGGCCCTGTACAG
P G N E S R S R M M E R L T K F K A G P G H V>

      430
      *      *
ACATGAGGTACC
TGTAATCCATGG
T *>

```

FIGURE 6

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
GCTAGGCGCGCCACCATGCGCCAASTTCGTGGCTGCCTGGACCTCGAAGGCTGCGCTATGAGTCTTTCTAA
CGATCGCGCGGTGGTACCGGTTCAAGCACCGAGGACTCGGACTTCCGACGGCGATCTCAGAAAGATT
      M A K F V A A W T L K A A A M S L L >

      80      90      100     110     120     130     140
      *      *      *      *      *      *      *
CGAGGTGCGAAACGTACGTTCTCTTATCATCCATCAGGCCCTCAAAGCCGAGATCGCGCAGAGACT
GGCTCCAGCTTTCATGCAAGAGAGATAGTAGGGTAGTCGCGGGGAGTTTCGGCTCTAGCGGGTCTCTGA
T E V E T Y V L S I I P S G P L K A E I A Q R L >

      150     160     170     180     190     200     210
      *      *      *      *      *      *      *
TGAGGATGTTTTGCGAGGGAAGAACACAGNCTTGAGGCTCTCATGGAAATGGCTAAAGACAAGACCAATC
ACTCTACAAAAAGTCCCTCTCTGTCTAGAACTCGAGAGTACCTTACCGAATTTCTGTTCTGGTTAG
E D V F A G K N T D L E A L M E W L K T R P I >

      220     230     240     250     260     270     280
      *      *      *      *      *      *      *
CTGTCACTCTGACTAAGGGAATTTAGGGTTTGTTCACGCTCACCGTGCCCGTGAAGGAGTCTGC
GACAGTGGAGACTGATTCCTTAAATCCAAACCAAGTGCAGTGGCGCGGCTCACTCGTCTCTGACG
L S P L T K G I L G F V F T L T V P S E R G L >

      290     300     310     320     330     340     350
      *      *      *      *      *      *      *
AGCGTAGACGATTGTCCAAATGCGCTAAATGGGAATGGAGACCCAAACCAATGGACAGGGCAGTTAA
TCGACTCTGCTAAACAGGTTTACGGGAATTAACCTTACCTCTGGGTTTGTGTACCTGTCCCCTCAATT
Q R R R F V Q N A L N G N G D P N N M D R A V K >

      360     370     380     390     400     410     420
      *      *      *      *      *      *      *
ACTATACAAGAGCTGAGAGGGGAATGACATTCATGGAGCAAGGAAGTTGCACTCAGTTACTCAACT
TGATATGTTCTTGACTTCTCCCTTACTGTAAGTACTGTTCTCTCAACGTGAGTCAATGAGTTGA
L Y K K L K R E M T F H G A K E V A L S Y S T >

      430     440     450     460     470     480     490
      *      *      *      *      *      *      *
GGTGGCTTGCCAGTTGCATGGTCTCATATACACCGGATGGGAACAGTGACCACAGAAAGTGCTTTG
CCAAGCGAAGCGGTCAAGTACCCAGAGTATATTTGGGCTACCTTGTCACTGGTGCTTCCACGAGAAC
G A L A . S C M G L I Y N R M G T V T T E V A L >

      500     510     520     530     540     550     560
      *      *      *      *      *      *      *
GCCTAGTATGTGCCACTTGTGACAGATTGCTGATGCCCAACATCGGTCCTCAAGGAGAGTGGGACTAC
CGATCATACAGGTGAACTGTCTTAAGACTACGGGTTGTAGCCAGGGGTCTCGCTACCGGTGATG
G L V C A T C E Q I A D A Q H R S H R Q M A T T >

      570     580     590     600     610     620     630
      *      *      *      *      *      *      *
CACCAACCCATTAATCAGGCATGAGAACAGAAATGGTACTAGCCAGCACTACGGCTAAGGCCATGGAGCAA
TGTGTGGGTGATTAGTCCGTACTCTGTCTTACCATGATCGTGCTGTATGCGAATCCGGTACTCGGTT
T N P L I R E E N R M V L A S T T A K A M E Q >

      640     650     660     670     680     690     700
      *      *      *      *      *      *      *
ATGGCTGGATCAAGTGAGCGGAGCAGAGGCGCATGGAAGTGCAGGCTAGACAAATGGTGAGG
TACGACCTAGTTACTCTCTCGCTCTCTCGGCTACTCTCAGGTTCACTCCGATCTGTTTACCACTGTC
M A G S S E Q A A E A M E V A S Q A R Q M V Q >

```

FIGURE 7



```
      710      720      730      740      750      760      770
*   *   *   *   *   *   *   *   *   *   *   *
CAATGAGGACAAITGGGACTCACCTAGCTCCAGTGCAGGTCTAAAAGATGATCTTATTGAAAATTGCA
GTTACTCTCTTTAACCTCTGAGTGGGATCGAGGTCACTCCAGATTTTCTACTAGAATAACTTTTAAACGT
A M R T I G T H P S S S A G L K D D L I E N L Q>

      780      790      800      810
*   *   *   *   *   *   *   *
GGCTTACCAGAAAACGGATGGGGGTGCAGATGCAGCGATTCAAGTGA
CCGAATGGTCTTTGCCTACCCCAAGTCTAAGTCGCTAAGTTCACT
A Y Q K R M G V Q M Q R F K *>
```

FIGURE 7 CONTINUED

10 20 30 40 50 60 70

GCTAGCGCCGACCAATGGCCAAAGTTCTGTGGCTGCTGGACCTGAAGGCTCGCGCTCTGGAGATTGGG  
CGATCGCGCGGTGTGATCCGGTTCAAGCACCGAGCGACCTGGGACTTCCGCGCGAGAGACTTAACCG  
M A K F V A A N T L K A A A L E I G >

80 90 100 110 120 130 140

GACCTGCTCTGAAGCGCGAGAACATCACATCGAGATTCTTAGGACCCCTTCTCTGTGTACAGGGCGGGTT  
CTGGGACGACTTGGCGCTCTGTGATGTAGTCTTAAGATTCTGGGGAAGAGCACAATTGCGGCAAT  
G P C L N A E N I T S G F L G P L L V L Q G G C C A F >

150 160 170 180 190 200 210

TTTCTGTGTGACAAGAATTCTCCAAATACGCGAGAGTCTAGACTGGTGGTGAAGCTCTCTCAATTCTCTA  
AAGAAGCAACTGTTCTTAGAGTGTATTAGCGTCTCAGATCTGAGCACCAGCTGAAGAGTGTAAAGAT  
F L L T R I A L T I P Q S L D S W W T S L N F L >

220 230 240 250 260 270 280

GGGGGAATACCTGTGTCTTGGCCAAATTCGAGCTGCCCAACTCCAATCACTCACCACCACTCTTGTG  
CCCCCTTAGTCGACGACAGAACCGGTTTTAAGCTCAGGGGTGGAGGTAGTGAAGTGTGGGAGACAG  
G G T T V C L G Q N S Q S P T S N H S P T S C >

290 300 310 320 330 340 350

CTCCAACTGTCTGGTATATGCTCGGATGTGTCTCGCGGCTTTTATCACTTCTCTCTATCTCTCTGCT  
GAGGTGTGAACAGGACCAATGACAGCACTACACAGACGCGCAAAATAGTAGAAGSAGAGATAGGAGACGCA  
P P T C T G G Y R W M C L R R F I I F L F I L L >

360 370 380 390 400 410 420

ATGCTCACTCTCTTGTGGTCTCTCTGAGTACTACAGGATATGTTCGCGTGTGTCTCTTAATTCAGGA  
TAGGAGTAGAAGAACCAACAGGAAGACCTTAGTGTCCATACACGGGCAACAGGAGATATAGGTCT  
C L I F L L V L L D Y Q G N L P V C F L I P >

430 440 450 460 470 480 490

TTCTCAACACCGACGCGGACCATCGCGGACTGCATGACTACTGCTCAAGGAACCTCTATGTATCCCT  
AGGAGTTGTGGTCTGCCTTGATACGGCTGAGCTGACTTACTGATCAGAGTTCTCTGGAGATCATAGGGA  
S S T T S T G P C R T C M T T A Q G T S M Y P >

500 510 520 530 540 550 560

CTGTGTGCTTACCAAACTCTCGGACGGAAATGCACTGTATTCCCATCCCATCATCTGGGCTTTGG  
GGACACGACATGGTTTGGAGGCTGCTTTAACTGGGATCATTAGGTTAGGTTAGTAGACCGCAAGGCG  
S C C C T K P S D N G N C T C I P I P T S S W A F G >

570 580 590 600 610 620 630

AAAATTCTATGGGAGTGGGCTCAGCGCGTTTCTCTGGCTCAGTTTACTAGTGCATTTGTTCAATGG  
TTTTAAGGATACCTACCCGGAGTCGGGCAAGGAGCCGAGTCAAAATGATCAGGTTAAACAGATCAAC  
K F L W E W A S R A F S W L S L V P F V Q W >

FIGURE 8

**PCT/US99/10646**

FIGURE 8 CONTINUED

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
GCTAGCCGCCCCACCATGCGAGGGGCTCTAGAAGCCCTCAGAGCCCTGCTCTCTCTCTCTAT
CGATCGCGGGCGGTGCTACGGTCCCGCAGATCTTGGCAGTCTCGGACGGAGAGGAGGAGAAGACGTA
      *      *      *      *      *      *      *
      H P G C L E A L R A L P L L L F L S>

      80      90      100     110     120     130     140
      *      *      *      *      *      *      *
ACGCGCTGTTTGGGTCCCGATGCCAGGCCATCAGCCAGGCTGTGCAAGCCGCTCAGCCGGAATCAAGGA
TGCGGACAAACCCAGGGCTACGGTCCGGTAGTTCGGTCCGACACGCTCGCGGAGTCGGCTTTAGTTCGT
Y A C L G P G C Q A I S Q A V H A H A E I N E>

      150     160     170     180     190     200     210
      *      *      *      *      *      *      *
AGCTGGAGAACCCCTCCAGCTTATCGCCCTCCAAACGCTCCTATCCTGTTCTTCTGCTGACCAGAACT
TCGACCTCTCTGGGAGGTGGAATAGCGGGAGGTTTGGGAGGATAGGACAAGAAAGAGGACTGCTCTAG
A G R T P P A Y R P P N A P I L F F L L T R L>

      220     230     240     250     260     270     280
      *      *      *      *      *      *      *
CTGCAATCCCGGAGTCCCTGGAGCCAAAGTTCTGCTGCTGCGTGGACCTGAAGGCTCGCGCTGGATCA
GACTGTTAGGGGGTCAGGGACCTTCGGCTTCAAGCACCGGACGCTGGGACTTCGGACGGGACGCTAGT
L T I P Q S L D A K F V A A W T L K A A A G L>

      290     300     310     320     330     340     350
      *      *      *      *      *      *      *
TCTGCTGTTCTGTCAGTGTGCCAGGGACGCTGCTGCTATTCAGCAAAACGCTGCCAAATGAGAAGTT
AGAACGACAAGACAGCTCAGCAGGCTCCCTGGCAGGAGGATAGTCCCTTGGCACCGCTTTACTCTTCAA
I L L F C A V V P G T L L L F R K R W Q N E K F>

      360     370     380     390     400     410     420
      *      *      *      *      *      *      *
TGGGGTGGACATGCCAGATGACTATGAGATGAAAATCTCTATGAGGSCCTGAACCTTGATGACTGTTCT
ACCCCACTGTACGGTCTACTGATCTTCTACTTTTAGAGATACTCCCGGACTTGGAACTACTGCAAGA
G V D M P D D Y E D E N L Y E G L N L D D C S>

      430     440     450     460     470     480     490
      *      *      *      *      *      *      *
ATGTATGAGGACATCTCCAGGGGACTCCAGGGCACCTACAGGATGTGGGCAACCTCCACATTGGAGATG
TACATACTCTCTGAGAGGTCCTGAGGTCCCTGGATGGTCTACACCCGTTGGAGGTGTAACCTCTAC
H Y E D I S R G L Q G T Y Q D V G N L H I G C D>

      500     510
      *      *
CCCAGCTGGAAAGCCATGAGGTACC
GGGTGACCTTTTCGTACTCCATGG
A Q L E K P ">

```

FIGURE 9

```

      10      20      30      40      50      60      70
GCTAGCGCGCCGCAOCATGCGCCACACTGGTGTGTCTTCCATGCCCTGCCACTGGCTGTGTCTCTCTCC
CGATCGCGCGCGGTGGTACCGGTGTGACACGACAGAGGTACGGGACGGTGACCGACACAAAGGACGAG
      K A T L V L S S M P C H W L L F L L
      80      90      100      110      120      130      140
TGCTCTTCTCAGGTGAGCGGATCAGCCAGGCTGTGCAAGCGCGCTCAGCCGAAATCAGCAAGCTGGAG
ACGAGAAGAGTCCACTGGCTAGTGGTCCGACAGGTGCGGAGTGGCGCTTAGTGTCTCGAGCTTC
L L F S G E P I S Q A V H A A H A E I N E A G R
      150      160      170      180      190      200      210
AACCCCTCCAGCTTATCGCCCTCCAAAGCTCCTATCTGTCTTCTCTCTGACCAAAATCCTGACAATC
TTGGGAGAGTCCGAATAGCGGAGGTTTTCGAGGATAGGACAAAGAGACGACTGGTCTTAGGACTGTAG
T P P A Y R P P N A P I L F F L L T R I L T I
      220      230      240      250      260      270      280
CCCCAGTCCCTGACGCCAAGTTCGTGGCTGCGTGGACCCCTGAAGGCTGCGCTATTATCTTGATCCAGA
GGGCTCAGGACCTCGCGTTCAAGCACCGACGACCTGGGACTTCGACGCGCGATAATAGAACTGGTCT
P Q S L D A K F V A A W T L K A A A I I L I Q
      290      300      310      320      330      340      350
CCCTCTCATCATCTCTTCATCATGTGCCCCATCTTCTGTCTACTTGACAAGGATGACGCCAAGCTGG
GGGAGGAGTAGTAGGAGAAGTAGTAACACGGGTAGAAGGACGATGAAGTGTCTCTACTTCCCGTTCCGACC
T L L I I L F I I V P I F L L L D K D D G K A G
      360      370      380      390      400      410      420
GATCGAGGAAGATCACACTATGAGGGCTTGAACATTGACCAGACAGCCACCTATGAAGACATAGTGAAT
CTACCTCTTCTAGTGTGGATCTCCGAACTGTGAAGTGTCTGTGCGTGGATCTCTGTATCACTGA
M E E D H T Y E G L N I D Q T A T Y E D I V T
      430      440      450      460      470      480
CTTCGACACGGGAGGTAAAGTGGTGGTAGGAGGACATCCAGGCCAGGAATGAGGTACC
GAAGCCTCTCCCTCCATTTCACAGCCATCTCTCGTAGGTCCGCTCTACTCCATGG
L R T G E V K W S V G E H P G Q E

```

FIGURE 10

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
GCTAGCGCGCCACCATGGGAATCCAGGTGCAGATCCAGAGCCTGTTTCTGCTCCTCCTGTGGGTGCCCG
CGATCGCGCGCGGTGGTACCCCTTAGCTCCAGCTTAGGTCTCCGACAAAGACGAGGAGGACACCCACGGGC
      H G H Q V Q I Q S L F L L L W V P>

      80      90      100      110      120      130      140
      *      *      *      *      *      *      *
GGTCCCGAGGAATCAGCCAGGCTGTGCACGCCGCTCAGCCGGAATCAACGAAGCTGGAAGAACCCTCC
CCAGGCTCCTTAGTCCGTCCGACACGTGCCCGCAGTCCGGCTTTAGTTGCTTCGACCTTCTTGGGAGG
G S R G I S Q A V R A A H A E I N E A G R T P P>

      150      160      170      180      190      200      210
      *      *      *      *      *      *      *
AGCTTATCGCCCTCCAAACGCTCCTATCCTGTTCTTTCTGCTGACGAGAATCCTGACAAATCCCCAGTCC
TCGAATAGCGGGAGGTTTGGCAGGATAGGACAAGAAAGACGACTGGTCTTAGGACTGTTAGGGGCTCAGG
A Y R P P N A P I L F F L L T R I L T I P Q S>

      220      230      240      250      260
      *      *      *      *      *
CTGGACGCCAAGTTGCTGCGCTGCCCTCGACCCCTGAAGGCTGCCGCTTGAGGTACC
GACCTGCGGTTCAAGCACCGACGGACCTGGGACTTCGGACGGGGAATCCATGG
L D A K F V A A W T L K A A A ">

```

FIGURE 11

TTCCCCAG	ATG	CAC	AGG	AGG	AGA	AGC	AGG	AGC	TGT	CGG	GAA	GAT	CAG	AAG	49	
	Met	His	Arg	Arg	Arg	Ser	Arg	Ser	Cys	Arg	Glu	Asp	Gln	Lys		
	1				5				10							
CCA	GTC	ATG	GAT	GAC	CAG	CGC	GAC	CTT	ATC	TCC	AAC	AAT	GAG	CAA	CTG	97
Pro	Val	Met	Asp	Asp	Gln	Arg	Asp	Leu	Ile	Ser	Asn	Asn	Glu	Gln	Leu	
15					20				25						30	
CCC	ATG	CTG	GGC	CGG	CGC	CCT	GGG	GCC	CCG	GAG	AGC	AAG	TGC	AGC	CGC	145
Pro	Met	Leu	Gly	Arg	Arg	Pro	Gly	Ala	Pro	Glu	Ser	Lys	Cys	Ser	Arg	
				35				40					45			
GGA	GCC	CTG	TAC	ACA	GGC	TTT	TCC	ATC	CTG	GTG	ACT	CTG	CTC	CTC	GCT	193
Gly	Ala	Leu	Tyr	Thr	Gly	Phe	Ser	Ile	Leu	Val	Thr	Leu	Leu	Leu	Ala	
			50					55					60			
GGC	CAG	GCC	ACC	ACC	GCC	TAC	TTT	CTG	TAC	CAG	CAG	CAG	GGC	CGG	CTG	241
Gly	Gln	Ala	Thr	Thr	Ala	Tyr	Phe	Leu	Tyr	Gln	Gln	Gln	Gly	Arg	Leu	
			65				70					75				
GAC	AAA	CTG	ACA	GTC	ACC	TCC	CAG	AAC	CTG	CAG	CTG	GAG	AAC	CTG	CGC	289
Asp	Lys	Leu	Thr	Val	Thr	Ser	Gln	Asn	Leu	Gln	Leu	Glu	Asn	Leu	Arg	
	80					85						90				
ATG	AAG	CTT	CCC	AAG	CCT	CCC	AAG	CCT	GTG	AGC	AAG	ATG	CGC	ATG	GCC	337
Met	Lys	Leu	Pro	Lys	Pro	Pro	Lys	Pro	Val	Ser	Lys	Met	Arg	Met	Ala	
	95				100				105						110	
ACC	CCG	CTG	CTG	ATG	CAG	GCG	CTG	CCC	ATG	GGA	GCC	CTG	CCC	CAG	GGG	385
Thr	Pro	Leu	Leu	Met	Gln	Ala	Leu	Pro	Met	Gly	Ala	Leu	Pro	Gln	Gly	
				115				120						125		
CCC	ATG	CAG	AAT	GCC	ACC	AAG	TAT	GGC	AAC	ATG	ACA	GAG	GAC	CAT	GTG	433
Pro	Met	Gln	Asn	Ala	Thr	Lys	Tyr	Gly	Asn	Met	Thr	Glu	Asp	His	Val	
			130					135					140			
ATG	CAC	CTG	CTC	CAG	AAT	GCT	GAC	CCC	CTG	AAG	GTG	TAC	CCG	CCA	CTG	481
Met	His	Leu	Leu	Gln	Asn	Ala	Asp	Pro	Leu	Lys	Val	Tyr	Pro	Pro	Leu	
			145				150					155				
AAG	GGG	AGC	TTC	CCG	GAG	AAC	CTG	AGA	CAC	CTT	AAG	AAC	ACC	ATG	GAG	529
Lys	Gly	Ser	Phe	Pro	Glu	Asn	Leu	Arg	His	Leu	Lys	Asn	Thr	Met	Glu	
	160					165					170					
ACC	ATA	GAC	TGG	AAG	GTC	TTT	GAG	AGC	TGG	ATG	CAC	CAT	TGG	CTC	CTG	577
Thr	Ile	Asp	Trp	Lys	Val	Phe	Glu	Ser	Trp	Met	His	His	Trp	Leu	Leu	
	175				180					185					190	

FIGURE 12

TTT GAA ATG AGC AGG CAC TCC TTG GAG CAA AAG CCC ACT GAC GCT CCA	625
Phe Glu Met Ser Arg His Ser Leu Glu Gln Lys Pro Thr Asp Ala Pro	
195 200 205	
CCG AAA GAG TCA CTG GAA CTG GAG GAC CCG TCT TCT GGG CTG GGT GTG	673
Pro Lys Glu Ser Leu Glu Leu Glu Asp Pro Ser Ser Gly Leu Gly Val	
210 215 220	
ACC AAG CAG GAT CTG GGC CCA GTC CCC ATG TGAGAGCAGC AGAGGCGGTC	723
Thr Lys Gln Asp Leu Gly Pro Val Pro Met	
225 230	

FIGURE 12 Continued



CCGCCTGGCGC	ATG GCG CCC CGC AGC GCC CGG CGA CCC CTG CTG CTG CTA	229
	Met Ala Pro Arg Ser Ala Arg Arg Pro Leu Leu Leu Leu	
	1 5 10	
CTG CCT GTT GCT GCT GCT CGG CCT CAT GCA TTG TCG TCA GCA GCC ATG	277	
Leu Pro Val Ala Ala Ala Arg Pro His Ala Leu Ser Ser Ala Ala Met		
	15 20 25	
TTT ATG GTG AAA AAT GGC AAC GGG ACC GCG TGC ATA ATG GCC AAC TTC	325	
Phe Met Val Lys Asn Gly Asn Gly Thr Ala Cys Ile Met Ala Asn Phe		
	30 35 40 45	
TCT GCT GCC TTC TCA GTG AAC TAC GAC ACC AAG AGT GGC CCC AAG AAC	373	
Ser Ala Ala Phe Ser Val Asn Tyr Asp Thr Lys Ser Gly Pro Lys Asn		
	50 55 60	
ATG ACC TTT GAC CTG CCA TCA GAT GCC ACA GTG GTG CTC AAC CGC AGC	421	
Met Thr Phe Asp Leu Pro Ser Asp Ala Thr Val Val Leu Asn Arg Ser		
	65 70 75	
TCC TGT GGA AAA GAG AAC ACT TCT GAC CCC AGT CTC GTG ATT GCT TTT	469	
Ser Cys Gly Lys Glu Asn Thr Ser Asp Pro Ser Leu Val Ile Ala Phe		
	80 85 90	
GGA AGA GGA CAT ACA CTC ACT CTC AAT TTC ACG AGA AAT GCA ACA CGT	517	
Gly Arg Gly His Thr Leu Thr Leu Asn Phe Thr Arg Asn Ala Thr Arg		
	95 100 105	
TAC AGC GTT CAG CTC ATG AGT TTT GTT TAT AAC TTG TCA GAC ACA CAC	565	
Tyr Ser Val Gln Leu Met Ser Phe Val Tyr Asn Leu Ser Asp Thr His		
	110 115 120 125	
CTT TTC CCC AAT GCG AGC TCC AAA GAA ATC AAG ACT GTG GAA TCT ATA	613	
Leu Phe Pro Asn Ala Ser Ser Lys Glu Ile Lys Thr Val Glu Ser Ile		
	130 135 140	
ACT GAC ATC AGG GCA GAT ATA GAT AAA AAA TAC AGA TGT GTT AGT GGC	661	
Thr Asp Ile Arg Ala Asp Ile Asp Lys Lys Tyr Arg Cys Val Ser Gly		
	145 150 155	
ACC CAG GTC CAC ATG AAC AAC GTG ACC GGA ACG CTC CAT GAT GCC ACC	709	
Thr Gln Val His Met Asn Asn Val Thr Val Thr Leu His Asp Ala Thr		
	160 165 170	
ATC CAG GCG TAC CTT TCC AAC AGC AGC TTC ACG AGG GGA GAG ACA CGC	757	
Ile Gln Ala Tyr Leu Ser Asn Ser Ser Phe Ser Arg Gly Glu Thr Arg		
	175 180 185	

FIGURE 13

TGT GAA CAA GAC AGG CCT TCC CCA ACC ACA GCG CCC CCT GCG CCA CCC Cys Glu Gln Asp Arg Pro Ser Pro Thr Thr Ala Pro Pro Ala Pro Pro 190 195 200 205	805
AGC CCC TCG CCC TCA CCC GTG CCC AAG AGC CCC TCT GTG GAC AAG TAC Ser Pro Ser Pro Ser Pro Val Pro Lys Ser Pro Ser Val Asp Lys Tyr 210 215 220	853
AAC GTG AGC GGC ACC AAC GGG ACC TGC CTG GCG AGC ATG GGG CTG Asn Val Ser Gly Thr Asn Gly Thr Cys Leu Leu Ala Ser Met Gly Leu 225 230 235	901
CAG CTG AAC CTC ACC TAT GAG AGG AAG GAC AAC ACG ACG GTG ACA AGG Gln Leu Asn Leu Thr Tyr Glu Arg Lys Asp Asn Thr Thr Val Thr Arg 240 245 250	949
CTT CTC AAC ATC AAC CCC AAC AAG ACC TCG GCC AGC GGG AGC TGC GGC Leu Leu Asn Ile Asn Pro Asn Lys Thr Ser Ala Ser Gly Ser Cys Gly 255 260 265	997
GCC CAC CTG GTG ACT CTG GAG CTG CAC AGC GAG GGC ACC ACC GTC CTG Ala His Leu Val Thr Leu Glu Leu His Ser Ser Ser Arg Thr Thr Val Leu 270 275 280 285	1045
CTC TTC CAG TTC GGG ATG AAT GCA AGT TCT AGC CGG TTT TTC CTA CAA Leu Phe Gln Phe Gly Met Asn Ala Ser Ser Ser Arg Phe Phe Leu Gln 290 295 300	1093
GGA ATC CAG TTG AAT ACA ATT CTT CCT GAC GCC AGA GAC CCT GCC TTT Gly Ile Gln Leu Asn Thr Ile Leu Pro Asp Ala Arg Asp Pro Ala Phe 305 310 315	1141
AAA GCT GCC AAC GGC TCC CTG CGA GCG CTG GAC GCC ACA GTC GGC AAT Lys Ala Ala Asn Gly Ser Leu Arg Ala Leu Gln Ala Thr Val Gly Asn 320 325 330	1189
TCC TAC AAG TGC AAC GCG GAG GAG CAC GTC GCT GTC ACG GCG TTT Ser Tyr Lys Cys Asn Ala Glu Glu His Val Arg Val Thr Lys Ala Phe 335 340 345	1237
TCA GTC AAT ATA TTC AAA GTG TGG GTC CAG GCT TTC AAG GTG GAA GGT Ser Val Asn Ile Phe Lys Val Trp Val Gln Ala Phe Lys Val Glu Gly 350 355 360 365	1285
GCC CAG TTT GGC TCT GTG GAG GAG TGT CTG CTG GAC GAG AAC AGC ACG Gly Gln Phe Gly Ser Val Glu Glu Cys Leu Leu Asp Glu Asn Ser Thr 370 375 380	1333

FIGURE 13. CONTINUED

CTG ATC CCC ATC GCT GTG GGT GGT GCC CTG GCG GGG CTG GTC CTC ATC	1381
Leu Ile Pro Ile Ala Val Gly Gly Ala Leu Ala Gly Leu Val Leu Ile	
385 390 395	
GTG CTC ATC GCC TAC CTC GTC GGC AGG AAG AGG AGT CAC GCA GGC TAC	1429
Val Leu Ile Ala Tyr Leu Val Gly Arg Lys Arg Ser His Ala Gly Tyr	
400 405 410	
CAG ACT ATC TAGCCTGGTG CACGCAGGCA CAGCAGCTGC AGGGGCCTCT	1478
Gln Thr Ile	
415	

FIGURE 13 CONTINUED

```

      10      20      30      40      50      60      70
* * * * *
ATGATCACCATTCTGCGCGTCTGCTGCTGGGGCTCAGCCTGGGCTGCACAGGAGCAGGTGGCTTCTGTGGCCC
TACTAGTGTGAAGGACGGCAGCAGCAGCCCGAGCTGGGACCGGACGTGTCCTGTCACCGAAGCAGCCGG
M I T F L P L L L G L S L G C T G A G G F V A>

      80      90      100      110      120      130      140
* * * * *
ATGTGGAAAGCACCTGCTGTGTTGGATGATGCTGGGACTCCAAAGGATTTCACACTGCATCTCCCTTCAA
TACACCTTCTGTGGACAGACAACTACTACGACCTGAGGTTTCTTAAAGTGTTATGACGTAGAGGAAGTT
H V E S T C L L D D A G T P K D F T Y C I S P N>

      150      160      170      180      190      200      210
* * * * *
CAAGGATCTGCTGACCTGCTGGGATCCAGAGGAGATAAGATGGCCCTTGGCAATTTGGGGTGTCTGAAT
GTTCCTAGACGACTGGACGACCTAGGTCTCTCTTATCTACCGGGGAACGCTTAAACCCCAAGCACTTA
K D L L T C W D P E E N K M A P C E F G V L N>

      220      230      240      250      260      270      280
* * * * *
AGCTTGGCGAATGTCTCTCAGACACCTCAACCAAAAGACACCTGATGCAGCGCTTGGCGAATGGGC
TGGAAACCGCTTACAGGAGATGTGCTGGAGTTGGTTTTCTGTGGGACTACGTGCGGAAACCGCTTACCGG
S L A N V L S Q R L N Q K D T L M Q R L R N G>

      290      300      310      320      330      340      350
* * * * *
TTCAGAATTGTGCCACACACACCCAGCCCTTCTGGGGATCACTGACCAACAGGACACGGCCACCATCTGT
AAGTCTTACACCGGTGTGTGTGGTGGGAAAGCCCTAGTGACTGGTTGTCTGTGCGGTGGTAGACA
L Q N C A T E T Q P F W G S L T N R T R P P S V>

      360      370      380      390      400      410      420
* * * * *
GCAAGTAGCCAAAACCACTCTTTTAAACAGAGGAGCCTGTGATGCTGGCCTGCTATGTGTGGGGCTTC
CGTTATCGGTTTTTGTGAGGAAATGTGCTCCCTCGGACACTACGACCGGACGATACACACCCGGAAG
Q V A K T T P F N T R E P V M L A C Y V W G F>

      430      440      450      460      470      480      490
* * * * *
TATCCAGCAGAAAGTGAATATCAGCTGGAGGAAGAACGGGAAGCTTGTCATGCCTCAGACAGTGGCACA
ATAGGTGCTGTTCTCACTGATGTCAGCTCTCTTCTGGCCCTTCGAACAGTACGGAGTGTGCTCAGCGGTGT
Y P A E V T I T W R K N G K L V M P H S S A H>

      500      510      520      530      540      550      560
* * * * *
AGACTGCCACGCCCAATGGAGACTGGACATACAGACCCCTCTCCCATTTAGCCTTAAACCCCTCTTAAGG
TCTCAGCGGTGGGTTACCTCTGACCTGTATGGCTCGGGAGAGGGTAAATCGGAATTGGGGGAGAAATGCC
K T A Q P N G D W T Y Q T L S H L A L T P S Y G>

```

FIGURE 14

```

      570      580      590      600      610      620      630
      *      *      *      *      *      *      *
GGACACTTACACTGTGTGGTAGAGCACATTGGGGCTCTGAGCCCATCTTCGGGACTGGACACTTGGG
CCTGTGAATGTGGACACACCATCTCGTGTAAACCCGAGGACTCGGGTAGGAAGCCCTGACCTGTGGACCC
D T Y T C V V E H I G A P E P I L R D W T P G>

      640      650      660      670      680      690      700
      *      *      *      *      *      *      *
CTGTCCCCCATGCAGACCCTGAAGGTTTCTGTGTCTGCAGTGACTCTGGGCTGGGCCTCATCATCTTCT
GACAGGGGGTACGTCTGGGACTTCCAAAGACACAGACGTCACTGAGACCCGGACCCGGAGTAGTAGAAGA
L S P M Q T L K V S V S A V T L G L G L I I F>

      710      720      730      740      750      760      770
      *      *      *      *      *      *      *
CTCTTGGTGTGATCAGCTGGCGGAGAGCTGGCCACTCTAGTTACACTCCTCTTCCTGGGTCCAATTATTC
GAGAACCACACTAGTCGACCGCCTCTCGACCGGTGAGATCAATGTGAGGAGAAGGACCCAGGTATATAG
S L G V I S N R R A G H S S Y T P L P G S N Y S>

      780      790
      *      *
AGAAGGATGGCACATTTCCTAG
TCTTCCTACCGTGTAAAGGATC
E G W H I S *>

```

FIGURE 14 Continued

```

      10      20      30      40      50      60      70
      *      *      *      *      *      *      *
ATGGTTTCTGGGTGGGTCCCTGGGTGGTGGCTCTGTAGTGAATCTGACCCAACTGGATTCTCTCCATGA
TACCCAGAGACCCACCCAGGGGACCCACACCGAGACGATCACTTAGACTGGGTTGACCTAAGGAGGTACT
M G S G W V P W V V A L L V N L T Q L D S S M>

      80      90      100      110      120      130      140
      *      *      *      *      *      *      *
CTCAAGGCACAGACTCTCCAGAAGATTTTGTGATTGAGGCAAGGCTGACTGTTACTTCACCAACGGGAC
GAGTTCGGTGTCTGAGAGGTCTTCTAAACACTAAGTCCGTTTCCGACTGACAAATGAAGTGGTTGCCCTG
T G G T D S P E D F V I Q A K A D C Y F T N G T>

      150      160      170      180      190      200      210
      *      *      *      *      *      *      *
AGAAAGGTGTCAGTTTGTGGTTCAGATTTCATCTTTAACTTGGAGGAGTATGTACGTTTCGACAGTGATGTG
TCTTTTCCAGCTCAAAACCCAGTCTAAGTAGAAATTGAACCTCTCTATACATGCAAAAGCTGTCACTACAC
E K V Q F V V R F I F N L E E Y V R F D S D V>

      220      230      240      250      260      270      280
      *      *      *      *      *      *      *
GGGATGTTTGGCATTGACCAAGCTGGGGCAGCCAGATGCTGAGCAGTGGAAACAGCCGGCTGGATCTCT
CCCTACAAACACCGTAACTGCTTCGACCCCGTGGTCTAGCAGTCTGTCACCTTGTGCGGCGACCTAGAGA
G M F V A L T K L G Q P D A E Q W N S R L D L>

      290      300      310      320      330      340      350
      *      *      *      *      *      *      *
TGGAGAGGAGCAGACAGGCCGTGGATGGGCTCTGTAGACACAACACAGGCTGGGGCGACCCCTTCACTGT
ACCTCTCTCTGCTGTCCGGCACCTACCCAGACATCTGTGTGATGTCGACCCGCGTGGGAAGTGACA
L E R S R Q A V D G V C R H N Y R L G A P F T V>

      360      370      380      390      400      410      420
      *      *      *      *      *      *      *
GGGGAGAAAGTGCAACAGAGGTGACAGTGATACCCAGAGAGGACCCACTCTGCACGACATAATCTG
CCCTCTCTTTACGTTGGTCTCCACTGTTCATGGGTCTCTCTCTGGGGTGAGGACGTTGGTCTATTAGAC
G R X V Q P E V T V Y P E R T F L L H Q H N L>

      430      440      450      460      470      480      490
      *      *      *      *      *      *      *
CTGCACTGCTCTGTGACAGGCTTCTATCCAGGGGATATCAAGATCAAGTGGTCTCTGAATGGCAGGAGG
GACGTACGACGACACTGTCCGAAGATAGGTCCCTATAGTCTAGTTCACCAAGGACTTACCCGCTCTCC
L H C S V T G F Y P G D I K I K W F L N G Q E>

      500      510      520      530      540      550      560
      *      *      *      *      *      *      *

```

FIGURE 15

AGAGAGCTGGGGTCATGTCACCTGGCCCTATCAGGAATGGAGACTGGACCTTTTCAGACTGTGTGATGCT  
 TCTCTCGACCCCGAGTACAGGTGACCGGGATAGTCTTACCTCTGACCTGGAAAGTCTGACACCACTACGA  
 E R A G V M S T G P I R N G D W T F Q T V V M L>

\* 570 \* 580 \* 590 \* 600 \* 610 \* 620 \* 630 \*  
 AGAAATGACTCCTGAACTTGGACATGCTACACCTGCCTTGTGATCACTCCAGCCTGCTGAGCCCTGTT  
 TCTTACTGAGGACTTGAACCTGTACAGATGTGGACGGAACAGCTAGTGAGGTGGGACGACTGGGACAA  
 E M T P E L G H V Y T C L V D H S S L L S P V>

\* 640 \* 650 \* 660 \* 670 \* 680 \* 690 \* 700 \*  
 TCTGTGGAGTGGAGAGCTCAGTCTGAATATTCTTGGAGAAAGATGCTGAGTGGCATTGCGCCCTCTCTAC  
 AGACACCTCACCCTCTCGAGTCAGACTTATAAGAACTCTTCTACGACTCACCGTAACGTGGGAAGGATG  
 S V E W R A Q S E Y S W R K M L S G I A A F L>

\* 710 \* 720 \* 730 \* 740 \* 750 \* 760 \* 770 \*  
 TTGGGCTAATCTTCTCTCTGGTGGGAATGTCATCCAGCTAAGGGCTCAGAAAGGATATGTGAGGACGCA  
 AACCCGATTAGAGGAAGAACCAACCCTTAGCAGTAGGTGATTCGCCGAGTCTTTCCTATACACTCCTCGGT  
 L G L I F L L V G I V I Q L R A Q K G Y V R T Q>

\* 780 \* 790 \* 800 \* 810 \* 820 \*  
 GATGTCTGGTAATGAGGTCTCAAGAGCTGTTCTGCTCCTCAGTCATGCTAA  
 CTACAGACCATTACTCCAGATTCTCGACAAAGACGAGGGAGTCAGTACGATT  
 M S G N E V S R A V L L P Q S C \*>

FIGURE 15 CONTINUED

10 20 30 40 50 60 70  
 ATGCTCTGGGGCTCCAGGAGTCTCTCAAAGCTCTGCCTGCACCACTCTTCTCTCTCTCTCTGCTGCTGCTG  
 TACGGACCCCCAGGTCCTCAGGAGGTTCCGAGACGGACGGTGTAGAAAGGAGGAGAAAGGACAGACAGAC  
 M P G G T C T V L Q A L P A T I F L L F L S A>  
 80 90 100 110 120 130 140  
 TCTACCTGGGCGCTCTGGGTGCCAGGCCCTGTGTGATGCACAAGGTCCTCCAGCATCATGTAGTGTGAGCCCTGGG  
 AGATGGACCCGGGACCCAGGTCCTGGGACACTACGTGTTCCAGGTCGTAGTAGTAACCTACCTCGGCCAGC  
 V Y L G P G C Q A L W M H K V P A S L M V S L G>  
 150 160 170 180 190 200 210  
 GGAAGTCCGCCCACTTCCAATGCCGCCGACATAGCAGCAACGCCCAAGTCACCTGGTGGCGGCTCTC  
 CTTCTCGGGGTGAAGTCTACGGGCGTTATGTGTCGTTGTTGGGTTGCAGTGACACCACCGCGGAGAG  
 E D A H F G C P H N S S N N A N V T W R V L>  
 220 230 240 250 260 270 280  
 CATGGCACTACACGTGGCGCCCTCTGAGTCTTGGGCGCGGCGAGGACCCCAATGGTAGCGTAGTCATCC  
 GTACCGTCTGATGTGCACGCGGGAATCAAGAACCGGCGCGCTCTCTGGGGTTACCATTCGCACTAGTAGG  
 H G N Y T W P P E F L G P G E D P N G T C T L I>  
 290 300 310 320 330 340 350  
 AGAATGTGAACAAGAGCCATCGGGGGCATATAGCTGTGTCGGGGTCCAGGAGGGCAACAGAGCTCATACCAACA  
 TCTTACACTTGTTCTCGGTACCCCGGTATATGACAGCGCCAGGTCCTCCGTTGCTCAGTAGTGTGTTGT  
 Q N V N K S H G G I V C R V Q E G N E S Y T Q G>  
 360 370 380 390 400 410 420  
 GTCTCTGGGCACTTACCTCTCGGCTGCGCCAGCGCGCCCCAGGCCCTCTTGGAACCTGGGGGAGGGACCC  
 CAGGACCCCGTGATGGAGGCGCACCGGTCGCGGGGGTCCGGGAAGGACTGTACCCGCTCCGTTG  
 S C G T Y L R V R Q P P R P R F L D M G E G T>  
 430 440 450 460 470 480 490  
 AAGAACCGAATCATCACGCCGAGGGGATCATCTCTGTTCTGCGCGGTGGTGCTCGGACGCTGCTGCTC  
 TCTTGGCTTAGTAGTGTGGCTCCCTCATGTAGGAGGACAGAGCGCCACCCAGGACCCCTCGAGCAGG  
 K N R I I T A T E G I I L L F C A V V P F G T L L>

FIGURE 16



```

      500      510      520      530      540      550      560
*   *   *   *   *   *   *   *   *   *   *   *
TGTT CAGGAACGATGGCAGAACGAGAAGCTCGGGTTGGATGCCGGGGATGAATATGAAGATGAAAACCT
ACAAAGTCCTTTGCTACCGTCTTTGCTCTTCGAGCCCAACCTACGGCCCTACTTATACTTCTACTTTTGA
L F R K R W Q N E K L G L D A G D E Y E D E N L>

      570      580      590      600      610      620      630
*   *   *   *   *   *   *   *   *   *   *   *
TTATGAAGGCCTGAACCTGGACGACTGCTCCATGTATGAGGACATCTCCCGGGGCTCCAGGGCACCTAC
AATACTTCCGACTTGGACCTGCTGACGAGGTACATACTCTGTAGAGGGCCCGGAGGTCCCGTGGATG
Y E G L N L D D C S M Y E D I S R G L Q G T Y>

      640      650      660      670      680      690      700
*   *   *   *   *   *   *   *   *   *   *
CAGGATGTGGGCAGCCTCAACATAGGAGATGTCCAGCTGGAGAAGCCGTGACACCCCTACTCCTGCCAGG
GTCCCTACACCGTCGGAGTTGTATCCTCTACAGGTCGACCTCTTCGGCACTGTGGGATGAGGACGGTCC
Q D V G S L N I G D V Q L E K P *>

```

FIGURE 16 CONTINUED

GAATTCGGCG GTGACC ATG GCC AGG CTG GCG TTG TCT CCT GTG CCC AGC	49
Met Ala Arg Leu Ala Leu Ser Pro Val Pro Ser	
1 5 10	
CAC TGG ATG GTG GCG TTG CTG CTG CTG CTC TCA GCT GAG CCA GTA CCA	97
His Trp Met Val Ala Leu Leu Leu Leu Ser Ala Glu Pro Val Pro	
15 20 25	
GCA GCC AGA TCG GAG GAC CGG TAC CGG AAT CCC AAA GGT AGT GCT TGT	145
Ala Ala Arg Ser Glu Asp Arg Tyr Arg Asn Pro Lys Gly Ser Ala Cys	
30 35 40	
TCG CGG ATC TGG CAG AGC CCA CGT TTC ATA GCC AGG AAA CGG CGC TTC	193
Ser Arg Ile Trp Gln Ser Pro Arg Phe Ile Ala Arg Lys Arg Arg Phe	
45 50 55	
ACG GTG AAA ATG CAC TGC TAC ATG AAC AGC GCC TCC GGC AAT GTG AGC	241
Thr Val Lys Met His Cys Tyr Met Asn Ser Ala Ser Gly Asn Val Ser	
60 65 70 75	
TGG CTC TGG AAG CAG GAG ATG GAC GAG AAT CCC CAG CAG CTG AAG CTG	289
Trp Leu Trp Lys Gln Glu Met Asp Glu Asn Pro Gln Gln Leu Lys Leu	
80 85 90	
GAA AAG GGC CGC ATG GAA GAG TCC CAG AAC GAA TCT CTC GCC ACC CTC	337
Glu Lys Gly Arg Met Glu Glu Ser Gln Asn Glu Ser Leu Ala Thr Leu	
95 100 105	
ACC ATC CAA GGC ATC CGG TTT GAG GAC AAT GGC ATC TAC TTC TGC CAG	385
Thr Ile Gln Gly Ile Arg Phe Glu Asp Asn Gly Ile Tyr Phe Cys Gln	
110 115 120	
CAG AAG TGC AAC AAC ACC TCG GAG GTC TAC CAG GGC TGC GGC ACA GAG	433
Gln Lys Cys Asn Asn Thr Ser Glu Val Tyr Gln Gly Cys Gly Thr Glu	
125 130 135	
CTG CGA GTC ATG GGA TTC AGC ACC TTG GCA CAG CTG AAG CAG AGG AAC	481
Leu Arg Val Met Gly Phe Ser Thr Leu Ala Gln Leu Lys Gln Arg Asn	
140 145 150 155	
ACG CTG AAG GAT GGT ATC ATC ATG ATC CAG ACG CTG CTG ATC ATC CTC	529
Thr Leu Lys Asp Gly Ile Ile Met Ile Gln Thr Leu Leu Ile Ile Leu	
160 165 170	

FIGURE 17

TTC ATC ATC GTG CCT ATC TTC CTG CTG CTG GAC AAG GAT GAC AGC AAG	577
Phe Ile Ile Val Pro Ile Phe Leu Leu Leu Asp Lys Asp Asp Ser Lys	
175 180 185	
GCT GGC ATG GAG GAA GAT CAC ACC TAC GAG GGC CTG GAC ATT GAC CAG	625
Ala Gly Met Glu Glu Asp His Thr Tyr Glu Gly Leu Asp Ile Asp Gln	
190 195 200	
ACA GCC ACC TAT GAG GAC ATA GTG ACG CTG CGG ACA GGG GAA GTG AAG	673
Thr Ala Thr Tyr Glu Asp Ile Val Thr Leu Arg Thr Gly Glu Val Lys	
205 210 215	
TGG TCT GTA GGT GAG CAC CCA GGC CAG GAG TGAGAGCCAG GTCGCCCAT	723
Trp Ser Val Gly Glu His Pro Gly Gln Glu	
220 225 230	

FIGURE 17 CONTINUED




FIGURE 18

```

      10      20      30      40      50      60      70
* * * * *
GACGGATCGGGAGATCTCCGATCCCTATGGTGGACTCTCAGTACAATCTGCTCTGATGCCGATAGTT
CTGCCTAGCCCTCTAGAGGGCTAGGGGATACCACTGAGAGTCATGTTAGACGAGACTACGGCGTATCAA

      80      90      100     110     120     130     140
* * * * *
AAGCCAGTATCTGCTCCCTGCTTGTGTGTGGAGGTCGCTNAGTAGTGGCGGAGCAAAATTTAAGCTACA
TTCCGTCATAGACGAGGGACGAACACACACCTCCAGGAGCTCATCAGCGGCTCGTTTAAATTCGATGT

      150     160     170     180     190     200     210
* * * * *
ACAAGGCAAGGCTTGACCGACAATTGCATGAAGAATCTGCTTAGGGTTAGGGGTTTTCGGCTGCTTCGGC
TGTTCCGTTCCGAACCTGGCTGTTAACTACTTCTTAGACGAATCCCAATCCGCAAAACCGCAGCAAGCCG

      220     230     240     250     260     270     280
* * * * *
ATGTAAGGGCCAGATATACGCGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTC
TACATGCCCCGCTCTATATGCGCACTGTAACATAAAGTATCAATAATTATCATTAGTTAATGCCCCAG

      290     300     310     320     330     340     350
* * * * *
ATTAGTTCATAGCCCATATATGAGGTTCCGGCTTACATAACTACGGTAATGGCCCGCTGGCTGACCG
TAATCAAGTATCGGGTATATACCTCAAGGCGCAATGATTGAATGCCATTACCGGCGGACCGACTGGC

      360     370     380     390     400     410     420
* * * * *
CCCAACGACCCCCGCCATTGACGTCAATAATGACGTAAGTCCCATAGTAACGCCAATAGGGACTTTCC
GGGTTGCTGGGGCGGGTAAGTGCAGTTATTACTGCATACAGGGTATCATTGCGGTTATCCCTGAAAGG

      430     440     450     460     470     480     490
* * * * *
ATTGACGTCAATGGGTGGACTATTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCC
TAAGTCAGTTACCCACTGATAAATGCCATTGACGGGTGAACCGTCATGTAGTTACATAGTATACGG

      500     510     520     530     540     550     560
* * * * *
AAGTACGCCCCCTATTGACGTCAATGAAGTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTA
TTCATGCGGGGATACTGCAGTTACTGCCATTACCGGGCGGACCGTAATACGGGTCAATGTACTGGAAT

      570     580     590     600     610     620     630
* * * * *
TGGGACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATGCTATTACCATGGTGATGCGGTTTGGC
AOCCTGAAGGATGAACCGTCATGTAGATGCATAATCAGTAGCATATGGTACCCTACCGCCAAACCG

      640     650     660     670     680     690     700
* * * * *
AGTACATCAATGGGCGTGATAGCGTTTGAAGTACAGGGGATTTCCAAGTCTCCACCCCATTTGAAGTCAA
TCATGTAGTTACCGCACCTATGCCCAACTGAGTGGCCCTAAAGGTTCAAGGTTGGGGTAACTGCAGTT

```

FIGURE 19

```

      710      720      730      740      750      760      770
      *      *      *      *      *      *      *
TGGGAGTTGTTTGGGCACAAATCAACGGGACTTTCACAAATGTCCTAACAACTCGGCCCATTTGAGG
ACCCCTCAACCAAAACCGTGGTTTTAGTTGCCCTGAAGGTTTTACAGCATTTGTTAGGCGGGGTAATGTC

      780      790      800      810      820      830      840
      *      *      *      *      *      *      *
CAAAATGGCGGTAGGCGGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCTCTGGCTAACTAGAGAACCCA
GTTTACCGGCATCGGCACATGCCACCTCCAGATATATCGCTCGAGAGACCGATTGATCTCTGGGT

      850      860      870      880      890      900      910
      *      *      *      *      *      *      *
CTGCTTACTGGCTTATCGAAATTAATAGACTCAGTATAGGGAGACCCAGCTGGCTAGAGTAAGTACCG
GACGAATGACCGAATAGCTTTAATTATGCTGAGTGATATCCCTCTGGGTTGACCGATCTCAITCATGGC

      920      930      940      950      960      970      980
      *      *      *      *      *      *      *
CCTATAGAGTCTATAGGCCACCCCTTGGCTTCTATGCATGCTTACTGTTTTGGCTTGGGGTCTAT
GGATACTCAGATATCCGGGTGGGGAAACCGAAGAACTAGTACGATATGACAAAACCGAACCCAGATA

      990      1000      1010      1020      1030      1040      1050
      *      *      *      *      *      *      *
ACACCCCGCTTCCTCATGTTATAGGTGATGGTATAGCTTAGCCTATAGGTGTGGGTATTGACCATAT
TGTGGGGGGAAGGAGTACAATATCCACTACCATATCGAATGGATATCCACCCCAATACCTAGTAATA

      1060      1070      1080      1090      1100      1110      1120
      *      *      *      *      *      *      *
TGACCACTCCCTTATGGTGAGGATACTTTCCATTACTAATCCATAACATGGCTCTTTGCCACAACTCTC
ACTGGTGAGGGGATAACCACTGCTATGAAAGGTAATGATTAGGTATGTACCGAGAAACGGTGTGAGAG

      1130      1140      1150      1160      1170      1180      1190
      *      *      *      *      *      *      *
TTTATGGCTATATGCCAATACAGTGTCTTCAGAGACTGACAGGACTCTGTATTTTACAGGATGGGG
AAATACCGATATACGGTTATGTGACAGGAAGTCTGACTGTGCTGAGACATAAAAAATGCTCAACCC

      1200      1210      1220      1230      1240      1250      1260
      *      *      *      *      *      *      *
TCTCATTTATTTTACAAATCACATATACACACCAACCGTCCCAAGTGCCCGCAGTTTTTTATAAACA
AGAGTAAATAATAATGTTTAAAGTGTATATGTTGGTGGCAGGGGTCAAGGGCGTCAAAAAATATTTGT

      1270      1280      1290      1300      1310      1320      1330
      *      *      *      *      *      *      *
TAACGTGGGATCTCCACGCGAATCTCGGGTACGTGTTCCGAGCATGGGCTCTCTCGGTAGCGGGGAG
ATTGACCCCTAGAGGTGGCTTAGAGCCCTACGACAGGCGTTACCCGAGAGGAGCCATCGCGCCTC

      1340      1350      1360      1370      1380      1390      1400
      *      *      *      *      *      *      *
CTTCTACATCGAGCGCCTGCTCCCATGCTCCAGCGACTCATGGTCTGCTGGCAGCTCTCTGCTCTAAC
GAGATGTAGGCTCGGACGAGGGTACGAGAGGTGCTGAGTACACGAGGCGCTGAGGAGCGAGGATTG

```

FIGURE 19 CONTINUED

```

      1410      1420      1430      1440      1450      1460      1470
*      *      *      *      *      *      *
AGTGGAGGCCAGACTTAGGCACAGCAGATGCCACACACCAGTGTGCCGACAAAGCCGTGCCGGTA
TCACCTCCGGTCTGAATCCGTGTCTGTCTACGGGTGGTGGTGCACACGGCGTGTTCGGCACCGCCAT

      1480      1490      1500      1510      1520      1530      1540
*      *      *      *      *      *      *
GGGTATGTCTGAAJATGAGCTCGGGGAGCGGGCTGCACCGCTGACGCAITTTGGAAGACTTAGGGCAG
CCCATAACAGACTTTTACTCGAGCCCTCGCCGAACGTGGCGACTGCGTAAACCTTCTGAATTCGTC

      1550      1560      1570      1580      1590      1600      1610
*      *      *      *      *      *      *
CGGCAGAAGAAGATGCAGGCAGCTGAGTTGTTGTGTTCTGATAGAGTCAAGGTAACCTCCGTTGCGGT
GCCGTCCTTCTACGTCCGTGCATCAACAACACAAGACTATTCTCAGTCTCATTGAGGGCAACGCCA

      1620      1630      1640      1650      1660      1670      1680
*      *      *      *      *      *      *
GCTGTTAACGGTGGAGGGCACTGTAGTCTGAGCAGTACTCGTTGCTGCCCGCGCGCCACAGACATAAT
CGACAATGCCACCTCCGTCACATCAGACTCGTCATGAGCAACGACGCGCGCGGTGCTGTATTA

      1690      1700      1710      1720      1730      1740      1750
*      *      *      *      *      *      *
AGCTGACAGACTAACAGACTTTCTCTTCCATGGGTCTTTCTGCAGGCTAGCCGGCCTGAATTCGGATA
TCGACTGTCTGATTGCTGACAGGAAAGGTACCCAGAAAGACGTCGATCGGCCGGACTTAAGCCTAT

      1760      1770      1780      1790      1800      1810      1820
*      *      *      *      *      *      *
TCCAGCTTGATGAATAAAGATCAGAGCTCTAGTGATCTGTGTGTTGGTTTTTGTGTGCTCGAGCCCT
AGGTTCCGAACCTACTTATTTCTAGTCTCGAGATCACTAGACACACAACCAAAAAACACAGAGCTCGGG

      1830      1840      1850      1860      1870      1880      1890
*      *      *      *      *      *      *
AGCTGGTCTCTTCCGCTCAGAAGCCATAGAGCCACCGCATCCCCAGCATGCCCTGCTATTGTCTTCCA
TGSACCAAGAAAGCGGAGTCTTCGGTATCTCGGTTGGCGTAGGGGTGCTACGGACGATACCAAGAGGT

      1900      1910      1920      1930      1940      1950      1960
*      *      *      *      *      *      *
ATCCTCCCTCTGCTGCTCCTGCCCAACCCACCCCCAGATAGAAATGACACCTACTCAGACAAATCGGAT
TAGGAAGGGGAACGACAGGACGGGTGGGGTGGGGGCTTATCTTACTGTGGATGAGTCTGTATCGGTA

      1970      1980      1990      2000      2010      2020      2030
*      *      *      *      *      *      *
GCAATTCCTCATTTTATTAGGAAGGACAGTGGAGTGGCACCTTCCAGGGTCAAGGAAGGCACGGGG
CGTTAAGGAGTAAATAATCTTCTTCTGTGACCTCACCGTGGAGGTCCAGTCTCTCCGCCCCC

      2040      2050      2060      2070      2080      2090      2100
*      *      *      *      *      *      *
AGGGGCAACACAGATGGCTGGCACTAGAAAGCACAGTGGAGGCTGATCAGCGAGCTCTAGCGGTACC
TCCCGTTGTGTCTACCGACGTTGATCTCGTGTGACTCCGACTAGTCGCTCGAGATCGCATGG

```

FIGURE 19 CONTINUED

2110 2120 2130 2140 2150 2160 2170  
 GGCATTAGTCTATGGCCGACTCTAGATTTTCTCCTGGCGCGCCCTAGATGCACTGCTCGATCGACCTGC  
 CGGTAATCAGATACCGGCTGAGATCTAAAGAGGAGAACGCCGCGGATCTACGTACGAGCTAGCTGGAGC  
 2180 2190 2200 2210 2220 2230 2240  
 AGTTGGACCTGGGAGTGGACACCTGTGGGAGAGAAAGCAAAGTGGATGTCTATTGCTCACTCAAGTGTATGG  
 TCAACCTGGACCTCACTGTGGACACCTCTCTTTCGGTTTCACTACAGTAACAGTGAGTTCCACATAC  
 2250 2260 2270 2280 2290 2300 2310  
 CCAGATCTCAAGCCTGCCACACCTCAAGCTAGCTTGACACAAAGAGATTGCTCTTTCTGACCAGATGGA  
 GGTCTAGAGTTCGGACGGTGTGGAGTTCGATCGAACTGTGTTTTCTAACAGAAAGACTGGTCTACCT  
 2320 2330 2340 2350 2360 2370 2380  
 CGCGGCCACCTCAAAAGGCATCACCGCGGGCCAGGTGAATCAATCCTCCTCGTTTGGGAAACTGAC  
 GCGCCGCTGGGAGTTTCGGTGTGGCGCCCGGTCACCTTATAGTTTAGGAGGAGCAAAACCTTTGACTG  
 2390 2400 2410 2420 2430 2440 2450  
 AATCTTAGCGCAGAAAGTCACTGCCGCTTTTGAGAGGGAGTACTCACCCACAGCTGGCCCTCGCAGACA  
 TTAGAATCGGCTCTTCAGTACGGCGGAAACTCTCCCTCATGAGTGGGTTGTGACCCGGGAGCGTCTGT  
 2460 2470 2480 2490 2500 2510 2520  
 GCGAATTAATCCAGCACACTGGCGGCGTTACTAGTGGATCGAGCTCGCAAGCTAGCTTGGGTCTCCC  
 CGCTTAATTAAGTCTGTGTGACCGCGGCAATGATCACTAGGCTCGAGCGTTCGATCGAACCCAGAGGG  
 2530 2540 2550 2560 2570 2580 2590  
 TATAGTGAGTCGTATTAATTTGATAAGCGAGTAGCAGTGGGTTCTCTAGTTAGCCAGAGAGCTCTGCT  
 ATATCACTCAGCATAATTAAGCTATTTCGTCATTGTCACCCAAAGAGATCAATCGGTCTCTCGAGACGA  
 2600 2610 2620 2630 2640 2650 2660  
 TATATAGACCTCCCAACGATACAGCGCTACCGCCATTGCGTCAATGGGGCGGAGTTGTTACGACATTTT  
 ATATATCTGGAGGGTGGCAGTGGCGGATGGCGGGTAACGAGTACCCCGCTCAACATGCTGTAAAA  
 2670 2680 2690 2700 2710 2720 2730  
 GGAAGTCCCGTTGATTTTGGTGC AAAACAAACTCCATTGACGTCAATGGGTTGGAGACTTGGAAATC  
 CCTTTCAGGGCACTAAACACCGGTTTGTGAGGGTAAGTCAGTTACCCACCTCTGAACCTTTAG  
 2740 2750 2760 2770 2780 2790 2800  
 CCGGTGAGTCAAAACGCTATCCAGCCATTGATGTACTGCCAAACCCGATCAACATGGTAATAGGAT  
 GGGCACTCAGTTTGGCGATAGGTGCGGGTAACATGACGGTTTGGGTAGTGGTACATTAATCGCTA

FIGURE 19 CONTINUED



```

      2810      2820      2830      2840      2850      2860      2870
      *      *      *      *      *      *      *
GACTAATACTAGATGTACTGCCAAGTAGGAAAGTCCCATAAGGTCAATGTACTGGGCATAATGCCAGGCG
CTGATTATGCATCTACATGACGGTTTCCTTCAGGGTATTCCAGTACATGACCCGTATTACGGTCCGC

      2880      2890      2900      2910      2920      2930      2940
      *      *      *      *      *      *      *
GGCCATTATACCGTCATTGACGCTCAATAGGGGGGCTACTTGGCATATGATACACTTGTACTGCGCAAC
CCGTAATGGCAGTAACCTGCAGTTATCCCCCGCATGAACCGTATACTATGTGAACCTACATGACGGTTCA

      2950      2960      2970      2980      2990      3000      3010
      *      *      *      *      *      *      *
GGCGGCTTACCGTAAATAGTCCACCCATTGACGCTCAATGGAAAGTCCCTATTGGCGTTACTATGGCAAC
CCCGTCAATGGCATTATCAGSTGGGTAACTGCAGTTACCTTTACGGGATAACCGCAATGATACCCCTG

      3020      3030      3040      3050      3060      3070      3080
      *      *      *      *      *      *      *
ATACGTCATTATTGACGCTCAATGGCGGGGGTGGTTGGGCGGTGAGCCAGGCGGGCCATTACCGTAAGT
TATGCAGTAATAACTGCAGTTACCCGCGCCCGCAGCAACCCGCACTCGGTCGCCCGGTAAATGGCAITCA

      3090      3100      3110      3120      3130      3140      3150
      *      *      *      *      *      *      *
TATTAACCGCGAACTCCATATATGGGCTATGAACATTAAGCCCGTAATTGATTACTATTAATAACTAG
ATACATTGGCGCTTGAGGTATATACCCGATACTGTATTACTGGGGGATTAACTAATGATAATTATTGATC

      3160      3170      3180      3190      3200      3210      3220
      *      *      *      *      *      *      *
TCAATAATCAATGTCTGCAATTAATGAATGGGCCAACGCGCGGGGAGAGGCGGTTTGGTATTGGGCGGT
AGTTATTAGTTACAGGACGTAATTACTTAGCCGGTTGCGGCGCCCTCTCCGCGCAAAACGATAAACCCGGA

      3230      3240      3250      3260      3270      3280      3290
      *      *      *      *      *      *      *
CTTCGGCTCTCTCGCTCACTGACTCGTGGCTCGGTGGTCTGGCTGCGGCGAGCGGTATCAGCTCACTC
GAAGGCGAAGGAGCGAGTGACTGAGCGACCGGAGCCAGCAAGCCGAGCGGCTCGCCATAGTCGAGTGG

      3300      3310      3320      3330      3340      3350      3360
      *      *      *      *      *      *      *
AAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAAGCGAGGAAAGAACATGTGAGCAAAAGGCCAG
TTCCGCCCAATTATGCCAATAGGTGTCTTAGTCCCTATTGGCTCCTTTCTGTACTACTGTTTTCGGTCTC

      3370      3380      3390      3400      3410      3420      3430
      *      *      *      *      *      *      *
CAAAAGGCGAGGAACCGTAAAGGCGCGGTTGCTGGCGTTTTCCATAGGCTCGCCCCCTGACGAGC
GTTTTGCGGTCTTGGCATTTTCCGGGCGCAAGACCGCAAAAGGTATCCGAGCGGGGGGACTGCTCG

      3440      3450      3460      3470      3480      3490      3500
      *      *      *      *      *      *      *
ATCAACAAAATCGAGCTCAAGTCAGAGGTGGCGAAACCGACAGGACTATAAAGATACAGGGGTTTCC
TAGTGTTTTAGTCTCGAGTTCAGTCTCCACGCTTTGGGCTGTCTGATATTTCTATGCTCCGCAAGG

```

FIGURE 19 CONTINUED

```

      3510      3520      3530      3540      3550      3560      3570
CCCTGGAAAGCTCCCTCGTGCCTCTCTGTTCCGACCTGCCGCTTACCGGATACCTGTCCGGCTTTCTC
GGGACCTTCGAGGGAGCACCGGAGGAGACAGGCTGGGACGGCGAATGGCTATGACAGCGCGGAAGAG

      3580      3590      3600      3610      3620      3630      3640
CCTTCGGGAAGCGTGGCGCTTCTCAATGCTCAGCGTGTAGGTATCTCAGTTCGGTGTAGGTGTTGCT
GGAAAGCCCTTCGACCGCGAAGAGTTACGAGTGGACATCCATAGAGTCAAGCCACATCCAGCAAGCGA

      3650      3660      3670      3680      3690      3700      3710
CCAACTGGGGCTGTGTGACGAACCCCGCTTCAGCCGACGCGCTGGCGCTTATCCGGTAACATGCTCT
GGTTCGACCCGACACACGTGCTTGGGGGGCAAGTCGGGCTGGGACCGGAAATAGGCCAATTGATGAGAGA

      3720      3730      3740      3750      3760      3770      3780
TGAGTCCAAACCGGTAAAGACAGGACTTATCGGCACTGGCAGCAGCCACTGGTAAACAGGATTAGCAGAGCG
ACTCAGGTTGGGCAATTCGTGTGAATAGCGGTGACCGTCTCGGTGACCAATTGCTCTAATCGTCTGCG

      3790      3800      3810      3820      3830      3840      3850
AGGTATGTAGGCGGTGCTACAGAGTCTTGAAGTGGTGGCTTAACACGGCTACACTAGAAAGGACAGTAT
TCCATACATCCGCCACGATGTCTCAAGAAGCTTACACACGGGATTGATGCCGATGTGATCTTCTGTGATA

      3860      3870      3880      3890      3900      3910      3920
TTGGTATCTGCGCTCTGCTGAAGCCGTTACCTTCGAAAAAGAGTTGGTAGCTCTTGATCCGGCAACA
AACCATAGACCGGAGACGACTTCGGTCAATGGAAGCTTTTCTCAACCATCGAGAAGCTAGGCGGTTTGT

      3930      3940      3950      3960      3970      3980      3990
AACCACCGCTGGTAGCGGTGGTTTTTTTGTGTTCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAA
TTGGTGGCGACATCGCCACCAAAAAACAACGTTCTGTGCTAATGCGGCTCTTTTCTTAGAGTT

      4000      4010      4020      4030      4040      4050      4060
GAAGATCCTTTGATCTTTCTACGGGCTCTGACGCTCAGTGGACGAAAACTCAGTTAAGGGATTTTGG
CTTCTAGGAAACTAGAAAAGATGCCAGACTGCGAGTCACTTGCTTTTGAAGTCAATTCCTATAAAGC

      4070      4080      4090      4100      4110      4120      4130
TCTGAACAATAAACTGCTGCTTACATAAACAGTATATCAAGGGGTGTTATGAGCCATATTCAACGGG
AGTACTGTATTATTGACAGCAATGATTGTCATTATGTTCCCAACAATCTCGGTATAGTTGCC

      4140      4150      4160      4170      4180      4190      4200
AAACGTTCTGCTCGAGGCGGOGATTAATAATCCAACTGAGTCTGATTATATGGGTATAAATGGGCTGG
TTTGCAAGCAGCTCGGCGCTAATTTAAGTTGTACTTCGACTAAATATACCAATATTTACCCGAGC

```

FIGURE 19 CONTINUED

4210      4220      4230      4240      4250      4260      4270  
 CGATAATGTCGGGCAATCAGGTGGGACAAATCTATCGATTGTATGGGAAGCCCGATGGCCAGAGTTGTTT  
 GCTATTACAGCCCGTTAGTCCACGCTGTGTAGATAGCTAACATACCCCTTCGGGCTACGCGGTCTCAACAAA  
  
 4280      4290      4300      4310      4320      4330      4340  
 CTGAACATGGCCAAAGGTAGCGTGGCCAATGATGTTACAGATGAGATGGTCAGACTAAACTGGCTGACGG  
 GACTTTGTACCGTTTCATCGCAACGGTTACTACAATGTCTACTCTACAGTCTGATTTGACCGACTGCC  
  
 4350      4360      4370      4380      4390      4400      4410  
 AATTATGCTCTTCGACCATCAAGCATTTTATCGTACTCGTATGATGCAATGTTTACTCACCCTGCG  
 TTAATACGGAGAAGGCTGGTAGTTCGTAAAAATAGGCATGAGGACTACTACGTACCAATGAGTGGTGACG  
  
 4420      4430      4440      4450      4460      4470      4480  
 GATCCCGGGGAAAAACAGCATTCCAGGTATTAGAAATATCTGATTCAGGTGAAAATATTTGTGATGCG  
 CTAGGGGCGCTTTTGTGCTAAGGTCCATAATCTTCTATAGGACTAAGTCCACTTTTATAACCACTACCG  
  
 4490      4500      4510      4520      4530      4540      4550  
 CTGGCAGTGTCTCTGCGCCGGTTCGATTGCGATTCCTGTTTGTAAATGTCCTTTTACAGCGATCGCGTAT  
 GACCGTCACAGGACGGGCGCAACGTAAAGCTAAGGACAAACATTACAGGAAAAATTTGCTAGCGCATAT  
  
 4560      4570      4580      4590      4600      4610      4620  
 TTGCTCTCGCTCAGGCGCAATCAGAAATGAATAACGGTTTGGTTGATGCGAGTGATTTTGAAGCAGGCG  
 AAGCAGAGCGAGTCCGCGTTAGTGTCTTACTTATGCAAAACCACTACGCTCACTAAACTACTGCTCGC  
  
 4630      4640      4650      4660      4670      4680      4690  
 TAATGGCTGGCTGTGTGAACAAGTCTGGAAAGAAATGCATAAACTTTTGCCATTCTCACCGAATTCAGTC  
 ATTACCGACCGACAACCTTGTTACAGACTTTCTTACGTATTGAAAAACGGTAAGAGTGGCCTTAAGTCAG  
  
 4700      4710      4720      4730      4740      4750      4760  
 GTCACTCATGGTGATTCTCACTTGATAACCTTATTTTGACGAGGGGAATTAATAGGTTGTATTGATG  
 CAGTGAGTACCCTAAAGAGTGAACCTATTGGAATAAAACTGCTCCCTTTAATTATCCAACAATACTAC  
  
 4770      4780      4790      4800      4810      4820      4830  
 TTGGACAGTCCGGAATCGCAGACCGATACCCAGGATCTTGCCATCTATGGAATCGCTCGGTGAGTTTTC  
 AACCTGCTCAGCCTTAGCGCTGGCTATGCTAGAACGGTAGGATACCTTGACGGAGCCACTCAAAAG  
  
 4840      4850      4860      4870      4880      4890      4900  
 TCCTTCATTACAGAAACGGCTTTTTCAAAAATATGGTATTGATAATCCTGATATGAATAAATTCAGGTTT  
 AGGAAGTAATGCTTTTGGCGAAAAAGTTTTATACCACTACTATTAGGACTATACTTATTAACTGCAAA

FIGURE 19 CONTINUED

```
      4910      4920      4930      4940      4950      4960      4970
      *      *      *      *      *      *      *
CAITTTGATGCTCGATGAGTTTTCCTAATCGAATTGGTTAATGGTGTGAACACTGGCAGAGCATCATGA
GTAAACTACGAGCTACTCAAAAAGATTAGTCTTAACCAATTAAACCAATTGTGACCGTCTCGTAGTACT

      4980      4990      5000      5010      5020      5030      5040
      *      *      *      *      *      *      *
GCCGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCTCCGAAAAGT
CGCTATGTATAAACTTACATAAAATCTTTTATTGTTTATCCCAAGGCGCGTGTAAAGGGGCTTTTCA

      5050
      *      *
GCCACCTGAAGTC
CGGTGGACTGCAG
```

FIGURE 19 CONTINUED

```

      10          20          30          40          50          60          70
GCTAGCGCCGCCACCATGGGAATGCAGGTGCAGATCCAGAGCCTGTTCTTCTGCTCTCTCTGGGTGCCCG
CGATCGCGGCGGTGTGATCTCTACGTTCACGTCTAGGTCTCGGACAAAGACGAGGAGCAACCCAGGCG
      M G M Q V Q I Q S L L L L L W V P>

      80          90          100          110          120          130          140
GGTCAGAGGACACACCCCTTGGAAGGCCGGAATCTGTATGAAGCCAAGTTCGTGCTGCTGCTGGACCT
CCAGGTCTCTGTGTGGGACACCTTCGCGGCTTAGGACATATTCCGGTTCGAAGCACCAGCGGACTGGGA
G S R G H T L W K A G I L Y K A K F V A A C T T G L>

      150          160          170          180          190          200          210
GAAGSGTCGCGCTTTCCTGCTAGCGAITTCITTCCTAGCGTGAAGCTGACCCCACTGTGCGTGACCCCTG
CTTCGACGGCGAAGGACGAGATCGCTAAGAAAGATCGCACTTCGACTGGGGTGACAGCCACTGGGAC
K A A A F L P S D F P S V K L T P L C V T L>

      220          230          240          250          260          270          280
TATATGGATGACGTGGTGTGCGGAGCCAGCATCATCAACTTCGAAAGCTGGGACTGTCCAGATACGTGG
ATATACCTACTGACCCAGCACTCTCGGTCTAGTAGTAGTGAAGCTTCGACCCCTGACAGGTCTATGACC>
Y M D D V V L G A S I I N F E K L S R Y T>

      290          300          310          320          330          340          350
CTAGGCTGATCCTGAAGAGGCCCTGTGCAGGGCTGTGCACCTGCCAGAGACCAACCGTGTGAGGAGGAC
GATTCGACTAGGACTTCTCTCGACACGTGCGGCAGCGTGGGACGGTGTCTGTGTGGCACCACTCCTCTG>
A R L I L K P F V H G V S T P T T G T V V R T>

      360          370          380          390          400          410
CGGTGCTACTAGGATGCCCTCTGGAAGTGGCTGAGCCTGCTGGTGCCCTTTGTGGGTACC
GCACATGATACCTCAGGACACACCTTCACGAGCTCGGAGACGACCAAGGAAACCCATCATG
V Y Y G V P V W K W L S L L V P F V G T>

```

37 / 42

```

10      20      30      40      50      60      70
* * * * *
GCTAGCGCCGCCACATGGGAATGCAGGTGCAGATCCAGAGCCTGTTCTGCTCCTCTGTGGGTGCCCG
CGATCGCGCGGTGGTACCTTACGTCCACGTCTAGGTCTCGGACAAAGACGAGGAGCACCCACCGGGC
      M G M Q V Q I Q S L F L L L L W V P>

      80      90      100      110      120      130      140
* * * * *
GGTCCAGAGGACACACCTGTGGAAGGCCGAATCCTGTATAAGGCCAAGTTCGTGGCTGCCTGGACCCCT
CCAGGTCTCTGTGTGGGACACCTTCGGGCTTAGGACATATCCGGTTCAGCACCGAAGGACCTGGGA
G S R G H T L W K A G I L Y K A K F V A A W T L>

      150      160      170      180      190      200      210
* * * * *
GAAGGCTGCGCTTTCCTGCCTAGCGATTCTTTCCTAGCGTGAAGCTGACCCCACTGTGCGTGACCCCTG
CTTCCGACGGGAAAGGACGGATCGCTAAAGAAAGGATCGCACTTCGACTGGGGTGACACGCACTGGGAC
X A A A F L P S D F F P S V K L T F L C V T L>

      220      230      240      250      260      270      280
* * * * *
TATATGGATGACGTGGTGTCTGGAGTGGGACTGTCCAGGTACGTGGCTAGGCTGATCCTGAAGGAGCCTG
ATATACCTACTGCACCAAGACCTTCACCTTGACAGSTCCATGCACCGATCCGACTAGGACTTCTCGGAC
Y M D D V V L G V G L S R Y V A R L I L K E P>

      290      300      310      320      330      340      350
* * * * *
TGCACGCGGTGTCCACCTGCCAGAGACCAACCGTGGTGAGGAGGACCGTGACTATGGAGTGCTGTGTG
ACGTCCCGCACAGGTGGGACGGTCTCTGGTGGCACCACTCTCTGGCACATGATACCTCACGGACACAC
V H G V S T L P E T T V V R R T V Y Y G V P V W>

      360      370      380      390
* * * * *
GAAGTGGCTGAGCCTGCTGGTGCCTTTGTGTGAGGTACC
CTTCACCGACTCGGACGACCAACGGGAAACACACTCCATGG
X W L S L L V P F V >>

```

FIGURE 21

Figure 22




Figure 23






Figure 24





Figure 25



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/85</b>		A3	(11) International Publication Number: <b>WO 99/58658</b>
			(43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/US99/10646		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 13 May 1999 (13.05.99)		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 69/078,904 13 May 1998 (13.05.98) US 60/085,751 15 May 1998 (15.05.98) US		(88) Date of publication of the international search report: 20 April 2000 (20.04.00)	
(71) Applicant: EPIMMUNE, INC. [US/US]; Suite 100, 5820 Nancy Ridge Drive, San Diego, CA 92121 (US).			
(72) Inventors: FIKES, John, D.; 6474 Lipmann Street, San Diego, CA 92122 (US). HERMANSON, Gary, G.; 3159 Via de Caballo, Encinitas, CA 92024 (US). SETTE, Alessandro; 5551 Linda Rosa Avenue, La Jolla, CA 92037 (US). ISHIOKA, Glenn, Y.; Apartment J5, 725 South Nardo Avenue, Solana Beach, CA 92075 (US). LIVINGSTON, Brian; 13555 Chaco Court, San Diego, CA 92129 (US). CHESNUT, Robert, W.; 1473 Kings Cross Drive, Cardiff-by-the-Sea, CA 92007 (US).			
(74) Agents: PARENT, Annette, S. et al.; Townsend and Townsend and Crew LLP, 8th floor, Two Embarcadero Center, San Francisco, CA 94111-3834 (US).			
(54) Title: EXPRESSION VECTORS FOR STIMULATING AN IMMUNE RESPONSE AND METHODS OF USING THE SAME			
(57) Abstract  The present invention relates to nucleic acid vaccines encoding multiple CTL and HTL epitopes and MHC targeting sequences.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LJ	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Cote d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
D3	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# INTERNATIONAL SEARCH REPORT

Intern. Appl. No.  
PCT/US 99/10646

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/85

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 35021 A (US HEALTH ;ZAREMBA SAM (US); SCHLOM JEFFREY (US); TSANG KWONG YOK) 25 September 1997 (1997-09-25)	1,3,5, 10,18, 20,22, 27,35, 37,38, 41,48, 49,52, 53,56,64
Y	page 7, line 7 - line 13; claim 36  page 13, line 24 - line 31 ---	2-9, 11-17, 19-26, 28-34, 36-40, 42-64

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"A" document member of the same patent family

Date of the actual completion of the international search

18 February 2000

Date of mailing of the international search report

01 02 00

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2000, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Sprinks, M

## INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US 99/10646

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THOMSON ET AL.: "Targeting a polypeptide protein incorporating multiple class II-restricted viral epitopes to the secretory/endocytic pathway facilitates immune recognition by CD4+ cytotoxic T lymphocytes: a novel approach to vaccine design." JOURNAL OF VIROLOGY (ONLINE), vol. 72, no. 3, March 1998 (1998-03), pages 2246-2252, XP002130251 cited in the application	1,2,13, 18,19, 30,35, 36,44
Y	abstract; figures 1-5; table 1	2-17, 19-34, 36-64
Y	WO 95 07707 A (CYTEL CORP) 23 March 1995 (1995-03-23) page 5, line 10	8,9,25, 26,39, 40,54,55
Y	WO 95 22317 A (CYTEL CORP) 24 August 1995 (1995-08-24)  page 17 page 23 page 25 page 26 page 28	10-12, 27-29, 41-43, 56-58
X	AN ET AL.: "A multivalent minigene vaccine, containing B-cell, cytotoxic T-lymphocyte, and Th epitopes from several microbes, induces appropriate responses in vivo and confers protection against more than one pathogen." JOURNAL OF VIROLOGY (ONLINE), vol. 71, no. 3, March 1997 (1997-03), pages 2292-2302, XP002130252 cited in the application	48,49
A	abstract	1-47, 50-64
A	WO 97 41227 A (T CELL SCIENCES INC ;THOMAS LAWRENCE J (US)) 6 November 1997 (1997-11-06) page 13 -page 14	1-64
A	US 5 633 234 A (AUGUST J THOMAS ET AL) 27 May 1997 (1997-05-27) cited in the application abstract; claim 15	1-64
	---	
	-/--	

# INTERNATIONAL SEARCH REPORT

Internat'l Application No  
PCT/US 99/10646

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>ULMER ET AL.: "Presentation of an exogenous antigen by major histocompatibility complex class I molecules." EUROPEAN JOURNAL OF IMMUNOLOGY, vol. 24, no. 7, July 1994 (1994-07), pages 1590-1596, XP002130253 the whole document</p> <p>---</p>	<p>13,30, 44,59</p>
P,X	<p>ISHIOKA ET AL.: "Utilization of MHC class I transgenic mice for development of minigene DNA vaccines encoding multiple HLA-restricted CTL epitopes." THE JOURNAL OF IMMUNOLOGY, vol. 162, no. 7, 1 April 1999 (1999-04-01), pages 3915-3925, XP000872894 the whole document</p> <p>-----</p>	<p>1-64</p>

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 99/10646**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 18-47 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/10646

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9735021	A	25-09-1997	AU 2535997 A EP 0888456 A	10-10-1997 07-01-1999
WO 9507707	A	23-03-1995	AU 698962 B AU 7873694 A CN 1135181 A EP 0735893 A JP 9505559 T US 5736142 A	12-11-1998 03-04-1995 06-11-1996 09-10-1996 03-06-1997 07-04-1998
WO 9522317	A	24-08-1995	AU 1847395 A AU 2500499 A CA 2183416 A EP 0804158 A	04-09-1995 24-06-1999 24-08-1995 05-11-1997
WO 9741227	A	06-11-1997	AU 2994697 A CA 2250428 A EP 0914427 A	19-11-1997 06-11-1997 12-05-1999
US 5633234	A	27-05-1997	AT 180835 T CA 2154445 A DE 69418856 D DE 69418856 T EP 0680513 A ES 2132395 T JP 8505878 T WO 9417192 A	15-06-1999 04-08-1994 08-07-1999 29-01-2000 08-11-1995 16-08-1999 25-06-1996 04-08-1994